



Full wwPDB X-ray Structure Validation Report

Oct 5, 2023 – 06:45 AM EDT

PDB ID : 6VT2
Title : Sialic acid binding region of Streptococcus sanguinis SK1 adhesin bound to sTa
Authors : Stubbs, H.E.; Iverson, T.M.
Deposited on : 2020-02-12
Resolution : 1.52 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : **FAILED**
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : **FAILED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.52 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 14469 atoms, of which 6220 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Adhesin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	H	N	O			
1	A	409	6264	1976	3078	553	657	0	16	0
1	E	409	6342	1990	3134	564	654	0	19	0

- Molecule 2 is an oligosaccharide called N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose-(1-3)-2-acetamido-2-deoxy-alpha-D-galactopyranose.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	K	3	46	25	2	19	0	0	0
2	C	3	46	25	2	19	0	0	0
2	D	3	46	25	2	19	0	0	0
2	F	3	46	25	2	19	0	0	0

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Ca		
3	A	4	4	4	0	0
3	E	4	4	4	0	0

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O S 5 4 1	0	0
4	A	1	Total O S 5 4 1	0	0
4	E	1	Total O S 5 4 1	0	0
4	E	1	Total O S 5 4 1	0	0
4	E	1	Total O S 5 4 1	0	0

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	E	1	Total	C	H	O	0	0
			14	3	8	3		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	818	Total	O	0	0
			818	818		
6	E	814	Total	O	0	0
			814	814		

MolProbity and EDS failed to run properly - this section is therefore empty.

3 Data and refinement statistics

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	82.21Å 269.86Å 47.51Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	35.66 – 1.52	Depositor
% Data completeness (in resolution range)	91.2 (35.66-1.52)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.85 (at 1.52Å)	Xtrriage
Refinement program	PHENIX 1.18.2_3874+SVN	Depositor
R, R_{free}	0.177 , 0.193	Depositor
Wilson B-factor (Å ²)	14.4	Xtrriage
Anisotropy	0.323	Xtrriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	14469	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 14.61% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

4 Model quality [i](#)

4.1 Standard geometry [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles [i](#)

4.3.1 Protein backbone [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates [i](#)

12 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	A2G	C	1	2	15,15,15	0.42	0	21,21,21	1.13	2 (9%)
2	GAL	C	2	2	11,11,12	0.60	0	15,15,17	0.88	0
2	SIA	C	3	2	20,20,21	2.02	3 (15%)	24,28,31	1.29	4 (16%)
2	A2G	D	1	2	15,15,15	0.37	0	21,21,21	1.23	2 (9%)
2	GAL	D	2	2	11,11,12	0.69	0	15,15,17	0.87	0
2	SIA	D	3	2	20,20,21	2.16	3 (15%)	24,28,31	1.43	3 (12%)
2	A2G	F	1	2	15,15,15	0.45	0	21,21,21	1.24	3 (14%)
2	GAL	F	2	2	11,11,12	0.59	0	15,15,17	0.88	0
2	SIA	F	3	2	20,20,21	2.01	2 (10%)	24,28,31	1.33	4 (16%)
2	A2G	K	1	2	15,15,15	0.42	0	21,21,21	1.34	2 (9%)
2	GAL	K	2	2	11,11,12	0.60	0	15,15,17	0.93	0
2	SIA	K	3	2	20,20,21	2.08	2 (10%)	24,28,31	1.40	3 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	A2G	C	1	2	-	0/6/26/26	0/1/1/1
2	GAL	C	2	2	-	0/2/19/22	0/1/1/1
2	SIA	C	3	2	-	0/18/34/38	0/1/1/1
2	A2G	D	1	2	-	0/6/26/26	0/1/1/1
2	GAL	D	2	2	-	0/2/19/22	0/1/1/1
2	SIA	D	3	2	-	0/18/34/38	0/1/1/1
2	A2G	F	1	2	-	0/6/26/26	0/1/1/1
2	GAL	F	2	2	-	0/2/19/22	0/1/1/1
2	SIA	F	3	2	-	0/18/34/38	0/1/1/1
2	A2G	K	1	2	-	0/6/26/26	0/1/1/1
2	GAL	K	2	2	-	0/2/19/22	0/1/1/1
2	SIA	K	3	2	-	0/18/34/38	0/1/1/1

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	3	SIA	C2-C1	8.41	1.59	1.52
2	K	3	SIA	C2-C1	8.04	1.59	1.52
2	F	3	SIA	C2-C1	7.95	1.59	1.52
2	C	3	SIA	C2-C1	7.87	1.59	1.52
2	K	3	SIA	O6-C2	2.74	1.47	1.43

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	3	SIA	O6-C2	2.38	1.46	1.43
2	F	3	SIA	O6-C2	2.37	1.46	1.43
2	D	3	SIA	C7-C6	2.28	1.55	1.53
2	C	3	SIA	O6-C2	2.15	1.46	1.43
2	C	3	SIA	C7-C6	2.05	1.55	1.53

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	K	1	A2G	C1-C2-N2	-4.85	105.11	110.73
2	D	1	A2G	C1-C2-N2	-3.94	106.17	110.73
2	D	3	SIA	O1A-C1-C2	-3.79	113.61	122.57
2	K	3	SIA	O1A-C1-C2	-3.64	113.96	122.57
2	F	1	A2G	C1-C2-N2	-3.53	106.64	110.73
2	C	1	A2G	C1-C2-N2	-3.41	106.78	110.73
2	D	1	A2G	C3-C2-N2	-3.10	104.76	110.62
2	F	3	SIA	O6-C2-C3	-2.99	106.35	110.46
2	C	1	A2G	C3-C2-N2	-2.90	105.14	110.62
2	K	3	SIA	O6-C2-C3	-2.79	106.62	110.46
2	C	3	SIA	O6-C2-C3	-2.61	106.86	110.46
2	D	3	SIA	O6-C2-C3	-2.61	106.87	110.46
2	F	1	A2G	C3-C2-N2	-2.60	105.70	110.62
2	F	3	SIA	O1A-C1-C2	-2.52	116.61	122.57
2	F	3	SIA	O1B-C1-O1A	2.34	129.41	124.09
2	C	3	SIA	O1B-C1-O1A	2.34	129.40	124.09
2	C	3	SIA	C4-C3-C2	2.31	113.95	109.81
2	K	3	SIA	C6-O6-C2	2.31	116.28	111.34
2	C	3	SIA	O1A-C1-C2	-2.29	117.15	122.57
2	F	3	SIA	C6-O6-C2	2.25	116.15	111.34
2	F	1	A2G	C6-C5-C4	-2.14	108.00	113.00
2	D	3	SIA	O1B-C1-O1A	2.09	128.83	124.09
2	K	1	A2G	C2-N2-C7	-2.03	118.25	123.18


There are no chirality outliers.


There are no torsion outliers.


There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.









4.6 Ligand geometry [i](#)

Of 14 ligands modelled in this entry, 8 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	SO4	E	1408	-	4,4,4	0.56	0	6,6,6	0.59	0
4	SO4	A	705	-	4,4,4	0.31	0	6,6,6	0.74	0
4	SO4	A	706	-	4,4,4	0.95	0	6,6,6	0.47	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	GOL	E	1401	-	5,5,5	1.44	0	5,5,5	0.75	0
4	SO4	E	1407	-	4,4,4	0.15	0	6,6,6	0.21	0
4	SO4	E	1406	-	4,4,4	0.46	0	6,6,6	0.48	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	E	1401	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	E	1401	GOL	C1-C2-C3-O3
5	E	1401	GOL	O2-C2-C3-O3

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	706	SO4	0	1

4.7 Other polymers [i](#)

There are no such residues in this entry.

4.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

5 Fit of model and data

5.1 Protein, DNA and RNA chains

EDS failed to run properly - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains

EDS failed to run properly - this section is therefore empty.

5.3 Carbohydrates

EDS failed to run properly - this section is therefore empty.

5.4 Ligands

EDS failed to run properly - this section is therefore empty.

5.5 Other polymers

EDS failed to run properly - this section is therefore empty.