



# Full wwPDB X-ray Structure Validation Report ⓘ

May 29, 2020 – 04:28 pm BST

PDB ID : 2VWS  
Title : Crystal structure of YfaU, a metal ion dependent class II aldolase from Escherichia coli K12  
Authors : Rea, D.; Rakus, J.F.; Gerlt, J.A.; Fulop, V.; Bugg, T.D.H.; Roper, D.I.  
Deposited on : 2008-06-26  
Resolution : 1.39 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.39 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1714 (1.40-1.40)
Clashscore	141614	1812 (1.40-1.40)
Ramachandran outliers	138981	1763 (1.40-1.40)
Sidechain outliers	138945	1762 (1.40-1.40)
RSRZ outliers	127900	1674 (1.40-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	267	
1	B	267	
1	C	267	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	PO4	C	1257	-	X	-	-

## 2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 6845 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called YFAU, 2-KETO-3-DEOXY SUGAR ALDOLASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	256	Total 1951	C 1233	N 338	O 371	S 9	0	0	0
1	B	256	Total 1952	C 1233	N 338	O 371	S 10	0	1	0
1	C	256	Total 1951	C 1233	N 338	O 371	S 9	0	0	0

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	P		
2	A	1	Total 5	O 4	P 1	0	0
2	A	1	Total 5	O 4	P 1	0	0
2	B	1	Total 5	O 4	P 1	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	C	1	Total	O	P	0	0
			5	4	1		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	C	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	320	Total	O	0	0
			320	320		
4	B	321	Total	O	0	0
			321	321		
4	C	318	Total	O	0	0
			318	318		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: YFAU, 2-KETO-3-DEOXY SUGAR ALDOLASE



- Molecule 1: YFAU, 2-KETO-3-DEOXY SUGAR ALDOLASE



- Molecule 1: YFAU, 2-KETO-3-DEOXY SUGAR ALDOLASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	105.20Å 136.80Å 123.20Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	59.76 – 1.39 59.80 – 1.39	Depositor EDS
% Data completeness (in resolution range)	99.5 (59.76-1.39) 99.5 (59.80-1.39)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.64 (at 1.39Å)	Xtrriage
Refinement program	REFMAC 5.1.9999	Depositor
R, $R_{free}$	0.171 , 0.191 0.171 , 0.190	Depositor DCC
$R_{free}$ test set	7041 reflections (3.99%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	18.0	Xtrriage
Anisotropy	0.305	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 50.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	6845	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	23.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.45% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.67	0/1988	0.85	3/2706 (0.1%)
1	B	0.60	0/1994	0.79	2/2714 (0.1%)
1	C	0.63	0/1988	0.83	6/2706 (0.2%)
All	All	0.64	0/5970	0.82	11/8126 (0.1%)

There are no bond length outliers.

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	171	ASP	CB-CG-OD2	6.88	124.49	118.30
1	C	188	ASP	CB-CG-OD2	6.09	123.78	118.30
1	B	144	ASP	CB-CG-OD2	5.92	123.63	118.30
1	A	192	HIS	N-CA-C	5.82	126.72	111.00
1	A	171	ASP	CB-CG-OD2	5.79	123.51	118.30
1	A	188	ASP	CB-CG-OD2	5.72	123.45	118.30
1	C	134	ARG	NE-CZ-NH1	5.58	123.09	120.30
1	C	120	ARG	NE-CZ-NH2	-5.28	117.66	120.30
1	B	171	ASP	CB-CG-OD2	5.25	123.03	118.30
1	C	144	ASP	CB-CG-OD2	5.25	123.03	118.30
1	C	239	ASP	CB-CG-OD1	5.04	122.83	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen



atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1951	0	1950	18	0
1	B	1952	0	1951	18	0
1	C	1951	0	1950	8	0
2	A	10	0	0	0	0
2	B	5	0	0	0	0
2	C	5	0	0	0	0
3	C	12	0	16	0	0
4	A	320	0	0	3	0
4	B	321	0	0	5	0
4	C	318	0	0	4	0
All	All	6845	0	5867	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:105:ALA:HB1	1:B:170:ILE:HD11	1.40	1.01
1:A:105:ALA:HB1	1:A:170:ILE:HD11	1.56	0.87
1:A:122:VAL:HG23	4:C:2071:HOH:O	1.81	0.79
1:B:181:SER:OG	1:B:190:ALA:HA	1.86	0.76
1:B:179:ASP:HB2	4:B:2230:HOH:O	1.85	0.75
1:B:13:LEU:HD12	4:B:2027:HOH:O	1.95	0.67
1:B:80:LYS:HG2	4:B:2131:HOH:O	2.02	0.60
1:A:187:PRO:HD2	1:A:189:ASN:HD21	1.68	0.58
1:C:158:LEU:HD11	1:C:180:LEU:HD11	1.85	0.58
1:B:153:GLU:OE1	1:B:153:GLU:HA	2.04	0.57
1:A:241:MET:HG2	4:B:2180:HOH:O	2.05	0.56
1:A:153:GLU:HB3	4:A:2221:HOH:O	2.04	0.56
1:B:222:MET:CE	1:B:226[B]:CYS:SG	2.95	0.54
1:A:189:ASN:ND2	1:A:189:ASN:H	2.06	0.54
1:B:80:LYS:HB2	1:B:81:PRO:HD3	1.90	0.54
1:C:158:LEU:CD1	1:C:180:LEU:HD11	2.39	0.53
1:C:205:ARG:HG3	4:C:2253:HOH:O	2.09	0.52
1:A:102:ALA:HB2	1:A:163:GLU:HB3	1.93	0.51
1:B:47:GLY:O	1:B:76:VAL:HG22	2.10	0.51
1:B:222:MET:HE3	1:B:226[B]:CYS:SG	2.51	0.51
1:C:152:VAL:CG2	1:C:175:ILE:HG12	2.41	0.50

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:115:PRO:HG2	4:C:2071:HOH:O	2.09	0.50
1:B:105:ALA:CB	1:B:170:ILE:HD11	2.28	0.49
1:B:124:ALA:HB1	1:B:129:ALA:HB3	1.96	0.48
1:C:47:GLY:O	1:C:76:VAL:HG22	2.14	0.48
1:A:124:ALA:HB1	1:A:129:ALA:HB3	1.96	0.48
1:A:176:GLY:HA3	4:A:2223:HOH:O	2.14	0.48
1:A:182:ALA:HA	1:A:187:PRO:HA	1.95	0.48
1:A:152:VAL:HG13	1:A:157:ALA:HB1	1.95	0.47
1:A:3:ALA:HB1	1:A:136:GLU:HG2	1.96	0.46
1:B:222:MET:HE2	1:B:226[B]:CYS:SG	2.55	0.46
1:C:80:LYS:HE3	1:C:119:GLU:HB3	1.98	0.46
1:C:76:VAL:HG23	4:C:2114:HOH:O	2.18	0.43
1:A:189:ASN:N	1:A:189:ASN:ND2	2.66	0.43
1:A:187:PRO:O	1:A:188:ASP:HB2	2.18	0.43
1:B:181:SER:HB3	1:B:186:TYR:O	2.19	0.43
1:B:80:LYS:HB2	1:B:81:PRO:CD	2.49	0.43
1:A:211:LYS:NZ	4:A:2276:HOH:O	2.41	0.42
1:B:249:GLN:NE2	4:B:2307:HOH:O	2.40	0.42
1:A:158:LEU:HD11	1:A:198:ILE:HG21	2.01	0.42
1:B:110:SER:OG	1:B:119:GLU:HG3	2.20	0.41
1:B:131:ARG:O	1:B:134:ARG:HD3	2.21	0.41
1:A:152:VAL:HG13	1:A:157:ALA:CB	2.51	0.41
1:C:152:VAL:HG23	1:C:175:ILE:HG12	2.02	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	254/267 (95%)	250 (98%)	2 (1%)	2 (1%)	19 4
1	B	255/267 (96%)	251 (98%)	4 (2%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	254/267 (95%)	246 (97%)	5 (2%)	3 (1%)	13	1
All	All	763/801 (95%)	747 (98%)	11 (1%)	5 (1%)	22	5

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	188	ASP
1	C	190	ALA
1	C	123	GLY
1	C	191	GLY
1	A	192	HIS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	203/211 (96%)	200 (98%)	3 (2%)	65	37
1	B	204/211 (97%)	199 (98%)	5 (2%)	47	14
1	C	203/211 (96%)	201 (99%)	2 (1%)	76	53
All	All	610/633 (96%)	600 (98%)	10 (2%)	62	33

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	23	TRP
1	A	155	LYS
1	A	189	ASN
1	B	23	TRP
1	B	119	GLU
1	B	155	LYS
1	B	179	ASP
1	B	188	ASP
1	C	23	TRP
1	C	225	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	189	ASN
1	B	189	ASN
1	B	249	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	PO4	C	1257	-	4,4,4	1.77	1 (25%)	6,6,6	2.59	3 (50%)
2	PO4	A	1258	-	4,4,4	0.79	0	6,6,6	0.42	0
3	GOL	C	1258	-	5,5,5	0.20	0	5,5,5	0.66	0
2	PO4	B	1257	-	4,4,4	1.98	1 (25%)	6,6,6	2.34	2 (33%)
3	GOL	C	1259	-	5,5,5	0.42	0	5,5,5	0.45	0
2	PO4	A	1257	-	4,4,4	1.60	1 (25%)	6,6,6	2.05	2 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	C	1259	-	-	0/4/4/4	-
3	GOL	C	1258	-	-	2/4/4/4	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1257	PO4	P-O1	3.76	1.59	1.50
2	C	1257	PO4	P-O1	3.32	1.58	1.50
2	A	1257	PO4	P-O1	3.16	1.58	1.50

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	1257	PO4	O2-P-O1	-4.10	95.88	110.89
2	C	1257	PO4	O4-P-O1	4.05	125.71	110.89
2	B	1257	PO4	O4-P-O1	3.92	125.23	110.89
2	B	1257	PO4	O2-P-O1	-3.60	97.73	110.89
2	A	1257	PO4	O4-P-O1	3.37	123.22	110.89
2	A	1257	PO4	O2-P-O1	-3.23	99.09	110.89
2	C	1257	PO4	O3-P-O2	2.28	115.30	107.97

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	1258	GOL	C1-C2-C3-O3
3	C	1258	GOL	O2-C2-C3-O3

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	256/267 (95%)	1.06	35 (13%) 3 2	13, 19, 31, 50	0
1	B	256/267 (95%)	0.88	35 (13%) 3 2	13, 20, 34, 40	0
1	C	256/267 (95%)	0.88	30 (11%) 4 4	14, 19, 29, 53	0
All	All	768/801 (95%)	0.94	100 (13%) 3 3	13, 20, 33, 53	0

All (100) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	190	ALA	33.3
1	C	190	ALA	18.0
1	A	189	ASN	15.5
1	C	189	ASN	11.9
1	B	117	TYR	11.4
1	C	187	PRO	9.5
1	A	191	GLY	9.3
1	A	256	SER	9.0
1	B	256	SER	8.7
1	C	256	SER	8.2
1	C	188	ASP	8.2
1	A	187	PRO	8.1
1	A	117	TYR	7.8
1	A	188	ASP	7.7
1	A	1	MET	7.7
1	C	191	GLY	6.8
1	A	116	PRO	6.4
1	A	192	HIS	6.4
1	B	187	PRO	5.9
1	A	186	TYR	5.5
1	B	1	MET	5.2
1	C	2	ASN	5.1
1	B	116	PRO	4.8

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	B	57	LEU	4.2
1	A	126	VAL	4.1
1	A	54	ILE	4.1
1	C	4	LEU	4.1
1	B	3	ALA	4.1
1	B	137	ASN	4.0
1	C	1	MET	3.9
1	C	229	TRP	3.6
1	B	188	ASP	3.6
1	C	117	TYR	3.5
1	A	2	ASN	3.5
1	B	190	ALA	3.3
1	A	125	SER	3.2
1	A	136	GLU	3.2
1	C	54	ILE	3.1
1	A	16	GLY	3.1
1	B	255	LYS	3.1
1	B	189	ASN	3.1
1	A	89	ILE	3.0
1	C	73	ILE	3.0
1	B	89	ILE	2.9
1	B	136	GLU	2.8
1	B	54	ILE	2.8
1	B	2	ASN	2.8
1	A	221	ASP	2.7
1	C	57	LEU	2.7
1	C	221	ASP	2.7
1	C	53	THR	2.7
1	B	4	LEU	2.7
1	C	14	ARG	2.7
1	A	4	LEU	2.6
1	C	86	VAL	2.6
1	C	27	THR	2.6
1	C	205	ARG	2.5
1	B	186	TYR	2.5
1	C	89	ILE	2.5
1	A	86	VAL	2.4
1	C	195	VAL	2.4
1	C	29	ALA	2.4
1	A	156	THR	2.4
1	B	73	ILE	2.4
1	B	61	LEU	2.4

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	53	THR	2.4
1	C	228	ALA	2.3
1	A	45	ILE	2.3
1	B	119	GLU	2.3
1	C	136	GLU	2.3
1	B	24	LEU	2.3
1	C	137	ASN	2.3
1	B	125	SER	2.3
1	B	197	ARG	2.3
1	A	53	THR	2.3
1	A	3	ALA	2.3
1	B	14	ARG	2.3
1	A	57	LEU	2.3
1	A	64	VAL	2.3
1	A	115	PRO	2.2
1	B	63	ALA	2.2
1	B	45	ILE	2.2
1	B	44	LEU	2.2
1	B	64	VAL	2.2
1	B	122	VAL	2.2
1	A	166	ASP	2.2
1	C	61	LEU	2.2
1	C	3	ALA	2.2
1	A	83	ILE	2.2
1	A	193	PRO	2.2
1	A	61	LEU	2.1
1	A	50	ALA	2.1
1	C	44	LEU	2.1
1	B	27	THR	2.1
1	B	72	VAL	2.1
1	C	58	TYR	2.1
1	B	50	ALA	2.1
1	A	162	ASP	2.0
1	B	47	GLY	2.0
1	A	158	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	PO4	A	1258	5/5	0.62	0.29	65,66,67,68	0
2	PO4	A	1257	5/5	0.68	0.27	22,26,29,30	0
2	PO4	C	1257	5/5	0.70	0.28	22,24,30,31	0
2	PO4	B	1257	5/5	0.71	0.24	21,25,29,32	0
3	GOL	C	1258	6/6	0.82	0.18	25,36,38,39	0
3	GOL	C	1259	6/6	0.84	0.16	22,24,25,26	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.