



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 15, 2024 – 08:12 AM EST

PDB ID : 8W0D
Title : Crystal structure of Acetyl-CoA synthetase 2 from *Candida albicans* in complex with an isopropyl AMP ester inhibitor
Authors : Seattle Structural Genomics Center for Infectious Disease; Seattle Structural Genomics Center for Infectious Disease (SSGCID)
Deposited on : 2024-02-13
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	686	 92% 5%
1	B	686	 9% 88% 8%
1	C	686	 72% 5% 24%

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 14611 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Acetyl-coenzyme A synthetase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	666	5154	3288	877	975	14	0	0	0
1	B	631	4912	3142	834	922	14	0	0	0
1	C	524	4117	2643	692	769	13	0	0	0

There are 51 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	initiating methionine	UNP Q8N3JN3
A	2	HIS	-	expression tag	UNP Q8N3JN3
A	3	HIS	-	expression tag	UNP Q8N3JN3
A	4	HIS	-	expression tag	UNP Q8N3JN3
A	5	HIS	-	expression tag	UNP Q8N3JN3
A	6	HIS	-	expression tag	UNP Q8N3JN3
A	7	HIS	-	expression tag	UNP Q8N3JN3
A	8	HIS	-	expression tag	UNP Q8N3JN3
A	9	HIS	-	expression tag	UNP Q8N3JN3
A	10	GLU	-	expression tag	UNP Q8N3JN3
A	11	ASN	-	expression tag	UNP Q8N3JN3
A	12	LEU	-	expression tag	UNP Q8N3JN3
A	13	TYR	-	expression tag	UNP Q8N3JN3
A	14	PHE	-	expression tag	UNP Q8N3JN3
A	15	GLN	-	expression tag	UNP Q8N3JN3
A	16	GLY	-	expression tag	UNP Q8N3JN3
A	403	ALA	VAL	engineered mutation	UNP Q8N3JN3
B	1	MET	-	initiating methionine	UNP Q8N3JN3
B	2	HIS	-	expression tag	UNP Q8N3JN3
B	3	HIS	-	expression tag	UNP Q8N3JN3
B	4	HIS	-	expression tag	UNP Q8N3JN3
B	5	HIS	-	expression tag	UNP Q8N3JN3
B	6	HIS	-	expression tag	UNP Q8N3JN3

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	7	HIS	-	expression tag	UNP Q8N3N3
B	8	HIS	-	expression tag	UNP Q8N3N3
B	9	HIS	-	expression tag	UNP Q8N3N3
B	10	GLU	-	expression tag	UNP Q8N3N3
B	11	ASN	-	expression tag	UNP Q8N3N3
B	12	LEU	-	expression tag	UNP Q8N3N3
B	13	TYR	-	expression tag	UNP Q8N3N3
B	14	PHE	-	expression tag	UNP Q8N3N3
B	15	GLN	-	expression tag	UNP Q8N3N3
B	16	GLY	-	expression tag	UNP Q8N3N3
B	403	ALA	VAL	engineered mutation	UNP Q8N3N3
C	1	MET	-	initiating methionine	UNP Q8N3N3
C	2	HIS	-	expression tag	UNP Q8N3N3
C	3	HIS	-	expression tag	UNP Q8N3N3
C	4	HIS	-	expression tag	UNP Q8N3N3
C	5	HIS	-	expression tag	UNP Q8N3N3
C	6	HIS	-	expression tag	UNP Q8N3N3
C	7	HIS	-	expression tag	UNP Q8N3N3
C	8	HIS	-	expression tag	UNP Q8N3N3
C	9	HIS	-	expression tag	UNP Q8N3N3
C	10	GLU	-	expression tag	UNP Q8N3N3
C	11	ASN	-	expression tag	UNP Q8N3N3
C	12	LEU	-	expression tag	UNP Q8N3N3
C	13	TYR	-	expression tag	UNP Q8N3N3
C	14	PHE	-	expression tag	UNP Q8N3N3
C	15	GLN	-	expression tag	UNP Q8N3N3
C	16	GLY	-	expression tag	UNP Q8N3N3
C	403	ALA	VAL	engineered mutation	UNP Q8N3N3

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is 5'-O-{(R)-hydroxy[(propan-2-yl)oxy]phosphoryl}adenosine (three-letter code: YHQ) (formula: C₁₃H₂₀N₅O₇P) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			26	13	5	7	1		
3	B	1	Total	C	N	O	P	0	0
			26	13	5	7	1		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	C	1	Total	C	N	O	P	0	0
			26	13	5	7	1		

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Na	0	0
			1	1		
4	C	1	Total	Na	0	0
			1	1		

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	S	0	0
			5	4	1		
5	A	1	Total	O	S	0	0
			5	4	1		
5	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 6 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	2	Total	Cl	0	0
			2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	1	Total 1	O 1	0	0
6	C	2	Total 2	O 2	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	159	Total 159	O 159	0	0
7	B	139	Total 139	O 139	0	0
7	C	24	Total 24	O 24	0	0


3 Residue-property plots i

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Acetyl-coenzyme A synthetase 2



- Molecule 1: Acetyl-coenzyme A synthetase 2



- Molecule 1: Acetyl-coenzyme A synthetase 2



ALA
PRO
LYS
SER
VAL
ILE
ILE
VAL
GLN
ASP
LEU
PRO
LYS
THR
ARG
SER
SER
LYS
ILE
MET
ARG
ARG
ILE
LEU
ARG
LYS
VAL
SER
SER
ASN
GLU
ALA
ASP
GLN
LEU
GLY
ASP
ILE
SER
THR
LEU
SER
SER
ASN
PRO
GLN
SER
SER
VAL
GLU
GLY
ILE
ILE
SER
ALA
PHE
GLY
ALA
GLN
PHE
GLY
LYS

LYS

4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, α , β , γ	139.49Å 139.49Å 544.12Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	49.61 – 2.70 49.61 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.9 (49.61-2.70) 100.0 (49.61-2.70)	Depositor EDS
R_{merge}	0.24	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.88 (at 2.69Å)	Xtrriage
Refinement program	PHENIX (dev_5233: ???)	Depositor
R, R_{free}	0.184 , 0.226 0.186 , 0.227	Depositor DCC
R_{free} test set	4335 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	54.8	Xtrriage
Anisotropy	0.576	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 38.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	14611	wwPDB-VP
Average B, all atoms (Å ²)	64.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.07% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: YHQ, NA, GOL, SO4, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.30	0/5284	0.52	0/7186
1	B	0.29	0/5037	0.51	0/6847
1	C	0.27	0/4236	0.48	0/5768
All	All	0.29	0/14557	0.50	0/19801

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5154	0	5036	17	0
1	B	4912	0	4792	17	0
1	C	4117	0	3981	17	0
2	A	6	0	8	0	0
3	A	26	0	0	0	0
3	B	26	0	0	0	0
3	C	26	0	0	1	0
4	A	1	0	0	0	0
4	C	1	0	0	0	0
5	A	10	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	5	0	0	0	0
6	A	2	0	0	0	0
6	B	1	0	0	0	0
6	C	2	0	0	0	0
7	A	159	0	0	0	0
7	B	139	0	0	0	0
7	C	24	0	0	0	0
All	All	14611	0	13817	50	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:495:VAL:HG12	1:C:548:VAL:HG11	1.79	0.63
1:A:579:VAL:HG13	1:A:591:VAL:HG13	1.87	0.56
1:C:169:TYR:CZ	1:C:202:ILE:HD11	2.41	0.56
1:A:298:THR:HG21	1:A:515:LYS:NZ	2.22	0.55
1:A:143:ASP:OD1	1:A:146:ARG:NH2	2.40	0.54
1:B:632:ILE:O	1:B:682:GLN:NE2	2.39	0.54
1:C:445:GLN:NE2	1:C:530:THR:O	2.41	0.53
1:A:493:GLU:OE1	1:A:545:ARG:NH1	2.38	0.53
1:C:169:TYR:CE1	1:C:202:ILE:HD11	2.44	0.52
1:B:25:VAL:HG22	1:B:535:ALA:HB1	1.91	0.51
1:C:131:ASP:OD1	1:C:223:GLY:N	2.43	0.50
1:A:579:VAL:CG1	1:A:591:VAL:HG13	2.42	0.50
1:B:130:ALA:O	1:B:371:ARG:NH2	2.45	0.50
1:C:505:MET:O	1:C:507:ARG:NH1	2.44	0.49
1:C:467:ALA:O	1:C:468:SER:OG	2.23	0.49
1:B:218:ASP:OD1	1:B:219:GLU:N	2.44	0.49
1:B:110:TYR:CD1	1:B:114:ASP:HB2	2.49	0.47
1:B:420:ILE:O	1:B:420:ILE:HG23	2.15	0.47
1:B:651:LYS:NZ	1:B:656:GLU:OE1	2.41	0.47
1:A:438:HIS:CD2	1:A:457:ALA:HA	2.51	0.46
1:C:420:ILE:O	1:C:420:ILE:HG23	2.15	0.46
1:C:509:VAL:HG23	1:C:516:TYR:HB2	1.97	0.46
1:B:102:LEU:H	1:B:102:LEU:HD12	1.81	0.46
1:B:614:VAL:HG13	1:B:626:ALA:HB1	1.99	0.45
1:A:386:VAL:HG22	1:A:387:ALA:N	2.32	0.45
1:B:169:TYR:CZ	1:B:202:ILE:HD11	2.52	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:102:LEU:HD22	1:B:118:PHE:CZ	2.53	0.44
1:A:420:ILE:O	1:A:420:ILE:HG23	2.18	0.43
1:C:218:ASP:OD1	1:C:219:GLU:N	2.50	0.43
1:A:522:ASN:N	1:A:523:PRO:CD	2.81	0.43
1:B:479:ILE:HD11	1:B:527:TYR:CG	2.54	0.43
1:C:480:ASP:O	1:C:484:GLY:N	2.49	0.43
1:C:479:ILE:HB	1:C:495:VAL:HG23	2.01	0.42
1:A:169:TYR:CZ	1:A:202:ILE:HD11	2.54	0.42
1:B:522:ASN:N	1:B:523:PRO:CD	2.82	0.42
1:A:177:ILE:HD12	1:A:177:ILE:N	2.33	0.42
1:A:413:LEU:HB3	1:A:428:TYR:CZ	2.54	0.42
1:A:610:ARG:NH2	1:A:682:GLN:O	2.49	0.42
1:B:509:VAL:HG23	1:B:516:TYR:HB2	2.02	0.42
1:A:447:GLU:HG2	1:A:520:TYR:CZ	2.54	0.42
1:B:386:VAL:HG22	1:B:387:ALA:N	2.34	0.42
1:C:413:LEU:HB3	1:C:428:TYR:CZ	2.55	0.42
1:B:347:LEU:C	1:B:347:LEU:HD23	2.40	0.42
1:C:522:ASN:N	1:C:523:PRO:CD	2.84	0.41
1:C:25:VAL:HG22	1:C:535:ALA:HB1	2.01	0.41
1:A:102:LEU:HD12	1:A:102:LEU:H	1.86	0.41
1:B:386:VAL:HG22	1:B:387:ALA:H	1.86	0.41
1:C:532:ASP:OD2	3:C:701:YHQ:O1	2.39	0.41
1:C:326:HIS:N	1:C:329:ASP:OD2	2.50	0.41
1:A:444:TRP:HB3	1:A:451:HIS:HA	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	664/686 (97%)	642 (97%)	20 (3%)	2 (0%)	41 66

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	B	621/686 (90%)	603 (97%)	17 (3%)	1 (0%)	47 73
1	C	520/686 (76%)	501 (96%)	18 (4%)	1 (0%)	47 73
All	All	1805/2058 (88%)	1746 (97%)	55 (3%)	4 (0%)	47 73

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	684	GLY
1	A	601	ASN
1	C	468	SER
1	A	684	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	542/568 (95%)	535 (99%)	7 (1%)	69 87
1	B	516/568 (91%)	514 (100%)	2 (0%)	91 97
1	C	432/568 (76%)	428 (99%)	4 (1%)	78 92
All	All	1490/1704 (87%)	1477 (99%)	13 (1%)	78 92

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	93	LEU
1	A	114	ASP
1	A	120	ASN
1	A	300	THR
1	A	434	LYS
1	A	615	LEU
1	A	643	LYS
1	B	114	ASP
1	B	120	ASN
1	C	93	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	114	ASP
1	C	120	ASN
1	C	161	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	445	GLN
1	C	343	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 14 ligands modelled in this entry, 7 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	YHQ	B	701	-	25,28,28	0.64	0	27,42,42	0.85	1 (3%)
5	SO4	A	704	-	4,4,4	0.62	0	6,6,6	0.32	0
5	SO4	A	705	-	4,4,4	0.62	0	6,6,6	0.09	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	YHQ	A	702	-	25,28,28	0.65	0	27,42,42	0.77	1 (3%)
3	YHQ	C	701	-	25,28,28	0.64	0	27,42,42	0.80	1 (3%)
5	SO4	B	702	-	4,4,4	0.62	0	6,6,6	0.20	0
2	GOL	A	701	-	5,5,5	0.32	0	5,5,5	0.26	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	YHQ	A	702	-	-	1/11/31/31	0/3/3/3
3	YHQ	C	701	-	-	0/11/31/31	0/3/3/3
3	YHQ	B	701	-	-	0/11/31/31	0/3/3/3
2	GOL	A	701	-	-	0/4/4/4	-

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	702	YHQ	C5-C8-N5	2.40	124.00	120.35
3	B	701	YHQ	C5-C8-N5	2.38	123.97	120.35
3	C	701	YHQ	C5-C8-N5	2.23	123.74	120.35

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	702	YHQ	C13-C11-O6-P1


There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	701	YHQ	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is

within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	666/686 (97%)	-0.21	7 (1%) 80 82	38, 53, 89, 149	0
1	B	631/686 (91%)	0.21	64 (10%) 7 5	39, 55, 130, 170	0
1	C	524/686 (76%)	-0.16	8 (1%) 73 76	48, 71, 104, 132	0
All	All	1821/2058 (88%)	-0.05	79 (4%) 35 33	38, 59, 104, 170	0

All (79) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	672	VAL	8.0
1	B	676	ILE	7.5
1	B	679	PHE	6.6
1	B	595	VAL	6.1
1	B	632	ILE	6.1
1	B	631	ILE	6.0
1	B	684	GLY	5.6
1	B	630	VAL	5.5
1	B	675	ILE	5.5
1	B	683	PHE	5.3
1	B	678	ALA	5.2
1	B	656	GLU	5.1
1	B	681	ALA	5.0
1	B	572	LYS	5.0
1	B	674	GLY	4.9
1	B	574	VAL	4.9
1	B	645	MET	4.7
1	B	576	GLU	4.7
1	B	653	SER	4.6
1	B	633	VAL	4.6
1	B	680	GLY	4.5
1	B	596	ALA	4.5
1	B	597	LEU	4.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	671	SER	4.4
1	B	573	LYS	4.3
1	B	569	ILE	4.3
1	B	646	ARG	4.2
1	B	655	ASN	4.1
1	B	677	SER	4.0
1	B	648	ILE	3.8
1	B	567	ALA	3.8
1	B	652	VAL	3.8
1	B	593	ALA	3.7
1	B	594	TYR	3.7
1	B	685	LYS	3.6
1	B	673	GLU	3.5
1	B	654	SER	3.4
1	B	651	LYS	3.4
1	B	682	GLN	3.3
1	B	610	ARG	3.3
1	B	669	PRO	3.3
1	B	570	GLU	3.3
1	B	578	ALA	3.3
1	B	577	ALA	3.2
1	A	607	GLU	3.2
1	B	629	SER	3.2
1	B	566	ALA	3.2
1	A	599	GLU	3.2
1	B	575	SER	3.1
1	A	606	SER	2.9
1	B	670	GLN	2.9
1	B	613	LEU	2.9
1	B	668	ASN	2.7
1	A	601	ASN	2.7
1	C	301	PRO	2.6
1	B	647	ARG	2.6
1	B	580	VAL	2.6
1	A	604	GLU	2.6
1	B	571	ASP	2.5
1	C	24	VAL	2.5
1	C	42	GLU	2.5
1	B	650	ARG	2.4
1	C	482	VAL	2.4
1	A	600	GLY	2.4
1	B	649	LEU	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	602	SER	2.4
1	B	568	LEU	2.3
1	C	28	ALA	2.3
1	C	432	VAL	2.3
1	B	585	ASP	2.2
1	B	579	VAL	2.2
1	C	431	PHE	2.2
1	B	564	ILE	2.2
1	B	562	ALA	2.2
1	C	390	ALA	2.2
1	B	586	ILE	2.2
1	B	565	GLU	2.1
1	B	553	ASN	2.1
1	B	563	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GOL	A	701	6/6	0.72	0.24	57,59,65,70	0
6	CL	A	707	1/1	0.73	0.26	75,75,75,75	0
6	CL	C	703	1/1	0.77	0.21	84,84,84,84	0
5	SO4	B	702	5/5	0.86	0.20	96,109,116,117	0
6	CL	B	703	1/1	0.91	0.12	90,90,90,90	0
4	NA	A	703	1/1	0.91	0.29	50,50,50,50	0
5	SO4	A	704	5/5	0.92	0.16	71,71,85,90	0
5	SO4	A	705	5/5	0.92	0.19	110,112,118,131	0
4	NA	C	702	1/1	0.93	0.19	54,54,54,54	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
6	CL	A	706	1/1	0.94	0.22	81,81,81,81	0
6	CL	C	704	1/1	0.95	0.08	82,82,82,82	0
3	YHQ	C	701	26/26	0.96	0.14	59,79,84,89	0
3	YHQ	B	701	26/26	0.98	0.15	49,53,60,62	0
3	YHQ	A	702	26/26	0.99	0.18	39,43,48,52	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around YHQ B 701:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.