

Full wwPDB NMR Structure Validation Report (i)

May 7, 2024 – 12:56 pm BST

PDB ID : 1W4F

Title: Peripheral-subunit from mesophilic, thermophilic and hyperthermophilic bac-

teria fold by ultrafast, apparently two-state transitions

Authors: Ferguson, N.; Sharpe, T.D.; Schartau, P.J.; Allen, M.D.; Johnson, C.M.; Fer-

sht, A.R.

Deposited on : 2004-07-23

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity: 4.02b-467

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

wwPDB-RCI : v 1n 11 5 13 A (Berjanski et al., 2005)

PANAV : Wang et al. (2010)

wwPDB-ShiftChecker : v1.2

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36.2

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $SOLUTION\ NMR$

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$rac{ m NMR~archive}{ m (\#Entries)}$
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Q	uality of chain			
1	A	47	40%	36%	6%	13%	•



2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 14 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues					
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model		
1	A:130-A:168 (39)	0.15	14		

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 4 clusters and 1 single-model cluster was found.

Cluster number	Models
1	1, 3, 4, 5, 7, 12, 13, 14, 17
2	2, 6, 9, 10
3	8, 15, 16, 20
4	18, 19
Single-model clusters	11



3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 723 atoms, of which 372 are hydrogens and 0 are deuteriums.

 \bullet Molecule 1 is a protein called DIHYDROLIPOYLLYSINE-RESIDUE ACETYLTRANSFERASE.

Mol	Chain	Residues		Atoms				Trace	
1	Λ	45	Total	С	Н	N	О	S	0
	A	45	723	217	372	71	62	1	U

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	166	TRP	PHE	engineered mutation	UNP P11961




4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

• Molecule 1: DIHYDROLIPOYLLYSINE-RESIDUE ACETYLTRANSFERASE




4.2 Scores per residue for each member of the ensemble


Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1

• Molecule 1: DIHYDROLIPOYLLYSINE-RESIDUE ACETYLTRANSFERASE




4.2.2 Score per residue for model 2






4.2.3 Score per residue for model 3

• Molecule 1: DIHYDROLIPOYLLYSINE-RESIDUE ACETYLTRANSFERASE




4.2.4 Score per residue for model 4

• Molecule 1: DIHYDROLIPOYLLYSINE-RESIDUE ACETYLTRANSFERASE




4.2.5 Score per residue for model 5

• Molecule 1: DIHYDROLIPOYLLYSINE-RESIDUE ACETYLTRANSFERASE




4.2.6 Score per residue for model 6

• Molecule 1: DIHYDROLIPOYLLYSINE-RESIDUE ACETYLTRANSFERASE




4.2.7 Score per residue for model 7






4.2.8 Score per residue for model 8

• Molecule 1: DIHYDROLIPOYLLYSINE-RESIDUE ACETYLTRANSFERASE




4.2.9 Score per residue for model 9

• Molecule 1: DIHYDROLIPOYLLYSINE-RESIDUE ACETYLTRANSFERASE




4.2.10 Score per residue for model 10

• Molecule 1: DIHYDROLIPOYLLYSINE-RESIDUE ACETYLTRANSFERASE




4.2.11 Score per residue for model 11

• Molecule 1: DIHYDROLIPOYLLYSINE-RESIDUE ACETYLTRANSFERASE




4.2.12 Score per residue for model 12






4.2.13 Score per residue for model 13

• Molecule 1: DIHYDROLIPOYLLYSINE-RESIDUE ACETYLTRANSFERASE




4.2.14 Score per residue for model 14 (medoid)

• Molecule 1: DIHYDROLIPOYLLYSINE-RESIDUE ACETYLTRANSFERASE




4.2.15 Score per residue for model 15

• Molecule 1: DIHYDROLIPOYLLYSINE-RESIDUE ACETYLTRANSFERASE




4.2.16 Score per residue for model 16

• Molecule 1: DIHYDROLIPOYLLYSINE-RESIDUE ACETYLTRANSFERASE




4.2.17 Score per residue for model 17





4.2.18 Score per residue for model 18

• Molecule 1: DIHYDROLIPOYLLYSINE-RESIDUE ACETYLTRANSFERASE




4.2.19 Score per residue for model 19

• Molecule 1: DIHYDROLIPOYLLYSINE-RESIDUE ACETYLTRANSFERASE



4.2.20 Score per residue for model 20





5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: ?.

Of the 20 calculated structures, 20 were deposited, based on the following criterion: NO VIOLATIONS > 0.25.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CNS	refinement	
ANSIG	structure solution	
Azara	structure solution	
CNS	structure solution	

No chemical shift data was provided.



6 Model quality (i)

6.1 Standard geometry (i)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	305	323	323	13±2
All	All	6100	6460	6460	269

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 21.

All unique clashes are listed below, sorted by their clash magnitude.

Atom 1	Atom-2	Clash(Å)	Distance(Å)	Models		
Atom-1	Atom-2	Clash(A)	Distance(A)	Worst	Total	
1:A:139:ALA:HB2	1:A:163:ILE:HD12	0.96	1.33	16	20	
1:A:146:ILE:HG23	1:A:158:VAL:HG21	0.74	1.57	10	13	
1:A:142:LYS:O	1:A:167:LEU:HD21	0.72	1.84	16	19	
1:A:146:ILE:HG23	1:A:158:VAL:CG2	0.72	2.15	10	13	
1:A:147:ARG:HG3	1:A:148:LEU:HD23	0.64	1.69	8	1	
1:A:130:ILE:HG22	1:A:157:ARG:HA	0.63	1.71	9	12	
1:A:135:VAL:HG11	1:A:158:VAL:HG12	0.62	1.70	4	18	
1:A:139:ALA:HB2	1:A:163:ILE:CD1	0.60	2.26	2	20	
1:A:130:ILE:HG22	1:A:157:ARG:CB	0.53	2.34	8	7	
1:A:142:LYS:N	1:A:142:LYS:HD2	0.52	2.20	12	3	
1:A:138:TYR:O	1:A:142:LYS:HG2	0.52	2.05	7	2	
1:A:131:ALA:HB3	1:A:136:ARG:HG3	0.51	1.83	12	3	
1:A:131:ALA:HB3	1:A:136:ARG:HG2	0.49	1.84	5	1	
1:A:139:ALA:CB	1:A:163:ILE:HD12	0.49	2.24	16	4	

Continued on next page...



Continued from previous page...

Atom 1	Atom 2	Cleab (Å)	Distance (Å)	Mod	dels
Atom-1	Atom-2	$\operatorname{Clash}(ext{\AA})$	$\operatorname{Distance}(\operatorname{\AA})$	Worst	Total
1:A:142:LYS:HD2	1:A:142:LYS:N	0.49	2.22	11	1
1:A:144:VAL:HG23	1:A:167:LEU:HD21	0.47	1.85	7	6
1:A:138:TYR:CE2	1:A:163:ILE:HB	0.47	2.44	7	20
1:A:142:LYS:HB3	1:A:167:LEU:HD11	0.46	1.87	18	17
1:A:138:TYR:CD2	1:A:163:ILE:HG21	0.46	2.45	7	15
1:A:136:ARG:O	1:A:140:ARG:HB2	0.46	2.10	20	4
1:A:130:ILE:HG22	1:A:157:ARG:CG	0.45	2.41	8	1
1:A:151:GLY:O	1:A:156:GLY:HA2	0.45	2.12	19	3
1:A:153:GLY:O	1:A:155:ASN:N	0.44	2.51	16	7
1:A:132:MET:HB2	1:A:135:VAL:HG23	0.43	1.90	11	9
1:A:150:GLN:CG	1:A:150:GLN:O	0.43	2.67	9	2
1:A:158:VAL:HG11	1:A:163:ILE:CD1	0.42	2.44	5	4
1:A:136:ARG:O	1:A:140:ARG:CB	0.42	2.67	11	6
1:A:148:LEU:O	1:A:166:TRP:CH2	0.42	2.73	5	19
1:A:139:ALA:HB1	1:A:144:VAL:O	0.42	2.15	1	1
1:A:148:LEU:O	1:A:166:TRP:CZ3	0.41	2.73	16	9
1:A:130:ILE:H	1:A:130:ILE:HD12	0.41	1.76	19	1
1:A:135:VAL:HG13	1:A:163:ILE:CG1	0.41	2.46	4	1
1:A:149:VAL:HG11	1:A:162:ASP:HB3	0.40	1.92	20	1
1:A:144:VAL:HG21	1:A:167:LEU:HG	0.40	1.92	6	4
1:A:148:LEU:HB3	1:A:166:TRP:CZ2	0.40	2.51	8	1
1:A:130:ILE:HG22	1:A:157:ARG:CA	0.40	2.46	19	1

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Pe	rce	ntiles
1	A	39/47 (83%)	36±0 (93±1%)	2±0 (4±1%)	1±0 (3±0%)		8	44
All	All	780/940 (83%)	729 (93%)	31 (4%)	20 (3%)		8	44

All 1 unique Ramachandran outliers are listed below.

Mol	Chain	Res	Type	Models (Total)
1	A	154	LYS	20



6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	31/36 (86%)	24±2 (78±6%)	7±2 (22±6%)	3	31
All	All	620/720 (86%)	486 (78%)	134 (22%)	3	31

All 16 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	152	THR	20
1	A	148	LEU	18
1	A	154	LYS	17
1	A	167	LEU	13
1	A	137	LYS	11
1	A	150	GLN	10
1	A	140	ARG	9
1	A	132	MET	8
1	A	160	LYS	7
1	A	155	ASN	5
1	A	142	LYS	4
1	A	147	ARG	3
1	A	134	SER	3
1	A	136	ARG	2
1	A	157	ARG	2
1	A	161	GLU	2

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



6.5 Carbohydrates (i)

There are no monosaccharides in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



7 Chemical shift validation (i)

No chemical shift data were provided

