

Full wwPDB X-ray Structure Validation Report (i)

Jun 27, 2024 – 01:09 PM JST

PDB ID	:	8W4I
Title	:	Crystal structure of EndoSz mutant D234M in space group P21
Authors	:	Guan, H.H.; Lin, C.C.; Hsieh, Y.C.; Chen, C.J.
Deposited on	:	2023-08-24
Resolution	:	2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R_{free}	130704	1957 (2.90-2.90)
Clashscore	141614	2172 (2.90-2.90)
Ramachandran outliers	138981	2115 (2.90-2.90)
Sidechain outliers	138945	2117 (2.90-2.90)
RSRZ outliers	127900	1906 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain			
1	А	992	6%	19%	•	12%



8W4I

2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 6903 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called glycoside hydrolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	А	876	Total 6902	C 4375	N 1152	O 1360	S 15	80	0	0

• Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	1	Total Ca 1 1	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: glycoside hydrolase



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	51.41Å 101.72Å 129.24Å	Deperitor
a, b, c, α , β , γ	90.00° 90.37° 90.00°	Depositor
$\mathbf{P}_{\text{assolution}}(\hat{\mathbf{A}})$	28.30 - 2.90	Depositor
Resolution (A)	28.30 - 2.90	EDS
% Data completeness	99.7 (28.30-2.90)	Depositor
(in resolution range)	99.6 (28.30-2.90)	EDS
R _{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.73 (at 2.90Å)	Xtriage
Refinement program	PHENIX 1.18.2_3874	Depositor
D D	0.229 , 0.273	Depositor
$\mathbf{n}, \mathbf{n}_{free}$	0.231 , 0.270	DCC
R_{free} test set	2002 reflections $(6.82%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	61.3	Xtriage
Anisotropy	0.395	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.33, 54.6	EDS
L-test for twinning ²	$< L >=0.50, < L^2>=0.33$	Xtriage
Estimated twinning fraction	0.031 for h,-k,-l	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	6903	wwPDB-VP
Average B, all atoms $(Å^2)$	69.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.19% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bo	nd lengths	Bond angles		
	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	0.37	1/7035~(0.0%)	0.61	1/9501~(0.0%)	

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	Observed(Å)	Ideal(Å)
1	А	134	LYS	CD-CE	5.22	1.64	1.51

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	$Observed(^{o})$	$Ideal(^{o})$
1	А	156	LYS	CD-CE-NZ	-5.06	100.06	111.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	6902	0	6887	128	0
2	А	1	0	0	0	0
All	All	6903	0	6887	128	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.



A 1 -		Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:A:453:PHE:HB2	1:A:459:ARG:HG3	1.64	0.78
1:A:364:VAL:HG11	1:A:412:LYS:HE3	1.65	0.77
1:A:704:THR:HG21	1:A:706:LYS:HE3	1.67	0.76
1:A:852:LEU:HD12	1:A:853:GLU:HG3	1.68	0.75
1:A:797:THR:HG22	1:A:798:SER:H	1.51	0.73
1:A:827:ALA:O	1:A:831:LYS:NZ	2.24	0.71
1:A:316:ASP:OD2	1:A:321:LYS:NZ	2.26	0.69
1:A:481:ARG:HA	1:A:503:GLU:HB2	1.74	0.68
1:A:113:PRO:HB2	1:A:439:MET:HE2	1.75	0.68
1:A:782:LYS:NZ	1:A:792:THR:O	2.25	0.66
1:A:302:LEU:HD22	1:A:343:MET:HB2	1.78	0.66
1:A:134:LYS:O	1:A:134:LYS:HG2	1.94	0.66
1:A:282:THR:HG22	1:A:304:GLN:HB2	1.79	0.64
1:A:299:ASP:O	1:A:341:GLN:HG2	1.97	0.64
1:A:829:TYR:HE2	1:A:869:GLN:HA	1.63	0.63
1:A:650:ALA:O	1:A:728:SER:HA	1.99	0.62
1:A:410:HIS:HD2	1:A:411:PRO:O	1.82	0.62
1:A:686:ILE:O	1:A:689:HIS:ND1	2.32	0.60
1:A:891:LYS:HG3	1:A:892:SER:H	1.66	0.60
1:A:615:THR:HG22	1:A:620:VAL:HB	1.85	0.59
1:A:248:SER:HB3	1:A:251:GLY:H	1.68	0.58
1:A:302:LEU:HB3	1:A:399:PHE:CE1	2.38	0.58
1:A:390:LYS:NZ	1:A:529:LEU:O	2.37	0.57
1:A:597:LEU:HD23	1:A:603:ASN:HB3	1.86	0.57
1:A:220:ASP:HA	1:A:224:TYR:HB2	1.85	0.57
1:A:291:ILE:O	1:A:295:ALA:N	2.36	0.57
1:A:319:ASN:HB2	1:A:321:LYS:NZ	2.19	0.57
1:A:484:ASN:O	1:A:487:ILE:HG12	2.05	0.57
1:A:715:SER:OG	1:A:731:GLU:OE2	2.23	0.56
1:A:173:LEU:HD23	1:A:176:GLN:NE2	2.21	0.56
1:A:795:LEU:HD11	1:A:896:TYR:HB3	1.87	0.56
1:A:779:ASN:HB3	1:A:795:LEU:O	2.06	0.56
1:A:125:LYS:O	1:A:129:PRO:HG3	2.06	0.56
1:A:541:ARG:HG3	1:A:542:LYS:H	1.71	0.56
1:A:387:TRP:O	1:A:395:LYS:HE3	2.07	0.55
1:A:788:LEU:HD23	1:A:901:GLU:HG2	1.89	0.55
1:A:848:VAL:HG22	1:A:849:LYS:H	1.72	0.55
1:A:959:ASP:OD2	1:A:959:ASP:N	2.40	0.55
1:A:397:GLY:HA2	1:A:439:MET:HE1	1.89	0.54
1:A:657:GLN:HG2	1:A:737:PHE:HE1	1.72	0.54

All (128) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



		Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:A:186:TRP:HB3	1:A:239:SER:O	2.08	0.53
1:A:235:ILE:HD11	1:A:290:LEU:HD23	1.90	0.53
1:A:818:PHE:CE1	1:A:871:PHE:HB3	2.43	0.53
1:A:951:LEU:HD23	1:A:961:ILE:HD13	1.91	0.53
1:A:453:PHE:HB2	1:A:459:ARG:CG	2.38	0.53
1:A:165:LEU:HA	1:A:169:HIS:HB2	1.89	0.53
1:A:301:LEU:HD23	1:A:337:ILE:HD13	1.91	0.52
1:A:913:ALA:O	1:A:917:THR:HG23	2.09	0.52
1:A:869:GLN:H	1:A:869:GLN:CD	2.12	0.52
1:A:709:LYS:HE3	1:A:738:PHE:HB2	1.91	0.52
1:A:709:LYS:NZ	1:A:743:SER:HB2	2.26	0.51
1:A:512:LYS:HG3	1:A:563:GLU:HB3	1.91	0.51
1:A:697:ASP:OD2	1:A:700:ALA:N	2.43	0.51
1:A:483:ASP:OD1	1:A:483:ASP:N	2.38	0.51
1:A:507:LEU:H	1:A:558:LEU:HD12	1.76	0.50
1:A:672:ILE:HD12	1:A:698:TYR:HE2	1.77	0.50
1:A:724:ASP:OD2	1:A:724:ASP:N	2.44	0.50
1:A:166:ALA:HB2	1:A:226:TYR:HB3	1.94	0.50
1:A:314:GLY:N	1:A:323:VAL:O	2.45	0.49
1:A:643:LYS:HG2	1:A:748:HIS:HE1	1.77	0.49
1:A:773:GLY:O	1:A:854:LYS:NZ	2.45	0.49
1:A:788:LEU:CD2	1:A:901:GLU:HG2	2.42	0.49
1:A:832:GLU:CD	1:A:890:LYS:HD3	2.32	0.49
1:A:182:ARG:HB2	1:A:228:LEU:HD12	1.94	0.49
1:A:674:SER:HB3	1:A:677:ASP:OD2	2.11	0.49
1:A:606:ILE:O	1:A:610:MET:HG3	2.13	0.49
1:A:819:PHE:O	1:A:900:PRO:HD2	2.13	0.48
1:A:675:GLU:HG3	1:A:698:TYR:CE1	2.49	0.48
1:A:863:ASP:OD1	1:A:864:TYR:N	2.46	0.48
1:A:653:THR:HG22	1:A:724:ASP:HB2	1.95	0.48
1:A:667:ASN:OD1	1:A:668:GLN:HG2	2.14	0.47
1:A:463:ILE:HG13	1:A:464:ALA:N	2.29	0.47
1:A:627:PHE:CZ	1:A:686:ILE:HB	2.49	0.47
1:A:845:SER:O	1:A:852:LEU:HD23	2.15	0.47
1:A:319:ASN:HB2	1:A:321:LYS:HZ2	1.79	0.47
1:A:456:LYS:O	1:A:460:GLU:HG3	2.14	0.47
1:A:902:LEU:HD21	1:A:904:ILE:HD11	1.97	0.47
1:A:731:GLU:HB3	1:A:733:TYR:CE1	2.50	0.46
1:A:709:LYS:HZ1	1:A:743:SER:HB2	1.81	0.46
1:A:99:MET:HE1	1:A:163:GLN:HA	1.97	0.46
1:A:173:LEU:HB3	1:A:178:THR:HB	1.97	0.46

Continued from previous page...



	Interatomic Clash						
Atom-1	Atom-2	distance (Å)	overlap (Å)				
1:A:794:THR:OG1	1:A:900:PRO:HA	2.16	0.46				
1:A:539:ASP:OD1	1:A:540:ASP:N	2.48	0.46				
1:A:852:LEU:CD1	1:A:853:GLU:HG3	2.44	0.46				
1:A:829:TYR:CE2	1:A:869:GLN:HA	2.45	0.46				
1:A:541:ARG:HG3	1:A:542:LYS:N	2.31	0.46				
1:A:256:ILE:HG22	1:A:293:ARG:HD2	1.99	0.45				
1:A:279:MET:HE1	1:A:291:ILE:HA	1.99	0.45				
1:A:672:ILE:HG13	1:A:672:ILE:O	2.16	0.45				
1:A:256:ILE:HA	1:A:290:LEU:HD11	1.99	0.45				
1:A:643:LYS:HG2	1:A:748:HIS:CE1	2.52	0.45				
1:A:657:GLN:HG2	1:A:737:PHE:CE1	2.52	0.44				
1:A:287:LYS:HE2	1:A:287:LYS:HB3	1.37	0.44				
1:A:526:LYS:HB2	1:A:526:LYS:HE3	1.80	0.44				
1:A:893:GLN:OE1	1:A:893:GLN:HA	2.18	0.43				
1:A:389:PRO:O	1:A:475:ARG:HD3	2.19	0.43				
1:A:466:VAL:HG22	1:A:476:PHE:CD2	2.52	0.43				
1:A:507:LEU:HB2	1:A:558:LEU:HD11	1.98	0.43				
1:A:542:LYS:O	1:A:544:GLU:N	2.46	0.43				
1:A:292:GLU:HG3	1:A:336:TYR:HE1	1.84	0.43				
1:A:360:TYR:H	1:A:410:HIS:CE1	2.36	0.43				
1:A:392:GLY:H	1:A:477:ASN:HB2	1.82	0.43				
1:A:832:GLU:OE1	1:A:890:LYS:HD3	2.19	0.43				
1:A:376:ILE:HD13	1:A:431:VAL:HG13	2.01	0.43				
1:A:118:TYR:HB2	1:A:401:TYR:HA	2.00	0.43				
1:A:374:SER:HA	1:A:431:VAL:HG23	1.99	0.43				
1:A:824:ILE:HD13	1:A:824:ILE:N	2.33	0.43				
1:A:673:ASN:OD1	1:A:673:ASN:N	2.48	0.42				
1:A:704:THR:HG22	1:A:706:LYS:H	1.85	0.42				
1:A:187:ARG:HG3	1:A:239:SER:O	2.19	0.42				
1:A:312:LYS:HE2	1:A:535:THR:O	2.20	0.42				
1:A:389:PRO:HG2	1:A:393:GLY:O	2.20	0.42				
1:A:448:ILE:HD13	1:A:473:LEU:HD21	2.02	0.42				
1:A:675:GLU:CD	1:A:675:GLU:H	2.23	0.42				
1:A:891:LYS:HG3	1:A:892:SER:N	2.34	0.42				
1:A:286:ASP:OD2	1:A:286:ASP:N	2.32	0.41				
1:A:826:LYS:HD2	1:A:827:ALA:CB	2.50	0.41				
1:A:435:LEU:O	1:A:439:MET:HG3	2.20	0.41				
1:A:420:ASP:OD1	1:A:420:ASP:N	2.52	0.41				
1:A:797:THR:HG22	1:A:798:SER:N	2.28	0.41				
1:A:227:ASN:OD1	1:A:275:ARG:NH2	2.53	0.41				
1:A:861:VAL:HG12	1:A:875:LEU:HD22	2.02	0.41				

Continued from previous page...



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)	
1:A:283:TYR:CG	1:A:287:LYS:O	2.73	0.41	
1:A:595:LEU:O	1:A:632:PRO:HG3	2.21	0.41	
1:A:120:ARG:HD2	1:A:122:TRP:CZ2	2.56	0.41	
1:A:911:GLU:HG2	1:A:914:THR:HB	2.02	0.40	
1:A:143:LYS:HE2	1:A:143:LYS:HB3	1.85	0.40	
1:A:602:GLU:HA	1:A:602:GLU:OE1	2.21	0.40	

Continued from previous page...

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percenti	les
1	А	874/992~(88%)	806 (92%)	63~(7%)	5 (1%)	25 58	8

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	А	878	ILE
1	А	132	LYS
1	А	599	ALA
1	А	821	ASP
1	А	822	SER

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	А	753/847~(89%)	741 (98%)	12 (2%)	62 86	

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	134	LYS
1	А	284	MET
1	А	286	ASP
1	А	287	LYS
1	А	604	ARG
1	А	626	VAL
1	А	707	ASP
1	А	822	SER
1	А	823	LYS
1	А	876	ASN
1	А	893	GLN
1	А	959	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	410	HIS
1	А	876	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.



There are no bond length outliers. There are no bond angle outliers. There are no chirality outliers. There are no torsion outliers. There are no ring outliers. No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	867/992~(87%)	0.29	58 (6%) 17 13	31, 69, 110, 160	1 (0%)

All (58) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	А	617	HIS	6.0
1	А	653	THR	6.0
1	А	416	PRO	5.9
1	А	844	ASP	5.2
1	А	972	ILE	5.0
1	А	419	PRO	4.8
1	А	414	ASN	4.4
1	А	823	LYS	3.7
1	А	841	HIS	3.7
1	А	845	SER	3.7
1	А	867	GLU	3.7
1	А	848	VAL	3.5
1	А	543	GLU	3.4
1	А	651	ASN	3.3
1	А	852	LEU	3.3
1	А	755	GLY	3.2
1	А	134	LYS	3.2
1	А	918	THR	3.2
1	А	728	SER	3.2
1	А	743	SER	3.2
1	А	925	LEU	3.1
1	А	650	ALA	3.1
1	А	415	GLY	3.0
1	А	618	GLY	3.0
1	А	973	ASP	3.0
1	А	541	ARG	2.9
1	А	542	LYS	2.9



Mol	Chain	Res	Type	RSRZ
1	А	798	SER	2.8
1	А	971	TYR	2.8
1	А	413	LYS	2.8
1	А	130	ALA	2.8
1	А	545	ALA	2.8
1	А	850	ASP	2.8
1	А	846	SER	2.7
1	А	824	ILE	2.7
1	А	742	ASN	2.7
1	А	540	ASP	2.6
1	А	744	THR	2.6
1	А	621	SER	2.5
1	А	418	THR	2.5
1	А	924	GLU	2.5
1	А	616	LYS	2.5
1	А	937	LEU	2.5
1	А	129	PRO	2.4
1	А	131	GLU	2.4
1	А	638	ASP	2.4
1	А	741	ILE	2.3
1	А	908	ARG	2.3
1	А	451	LYS	2.3
1	А	935	GLU	2.2
1	А	450	GLN	2.2
1	А	538	ASN	2.2
1	А	654	ILE	2.1
1	А	117	GLY	2.1
1	А	544	GLU	2.1
1	А	840	GLY	2.0
1	А	620	VAL	2.0
1	А	795	LEU	2.0

Continued from previous page...

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.



6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathrm{\AA}^2)$	Q<0.9
2	CA	А	1101	1/1	0.89	0.04	90,90,90,90	0

6.5 Other polymers (i)

There are no such residues in this entry.

