



# Full wwPDB X-ray Structure Validation Report ⓘ

May 14, 2020 – 10:14 pm BST

PDB ID : 1W5Q  
Title : Stepwise introduction of zinc binding site into porphobilinogen synthase of *Pseudomonas aeruginosa* (mutations A129C, D131C, D139C, P132E, K229R)  
Authors : Frere, F.; Reents, H.; Schubert, W.-D.; Heinz, D.W.; Jahn, D.  
Deposited on : 2004-08-09  
Resolution : 1.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1714 (1.40-1.40)
Clashscore	141614	1812 (1.40-1.40)
Ramachandran outliers	138981	1763 (1.40-1.40)
Sidechain outliers	138945	1762 (1.40-1.40)
RSRZ outliers	127900	1674 (1.40-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	337	 3% 81% 13% • 5%
1	B	337	 3% 82% 11% • 5%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	FMT	A	1337	-	-	X	-

## 2 Entry composition i

There are 6 unique types of molecules in this entry. The entry contains 6026 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DELTA-AMINOLEVULINIC ACID DEHYDRATASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	321	2704	1688	486	512	18	0	24	1
1	B	319	2688	1679	484	507	18	0	24	1

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	129	CYS	ALA	engineered mutation	UNP Q59643
B	129	CYS	ALA	engineered mutation	UNP Q59643
A	131	CYS	ASP	engineered mutation	UNP Q59643
B	131	CYS	ASP	engineered mutation	UNP Q59643
A	132	GLU	PRO	engineered mutation	UNP Q59643
B	132	GLU	PRO	engineered mutation	UNP Q59643
A	139	CYS	ASP	engineered mutation	UNP Q59643
B	139	CYS	ASP	engineered mutation	UNP Q59643
A	139	CYS	ASP	engineered mutation	UNP Q59643
B	139	CYS	ASP	engineered mutation	UNP Q59643
A	199	VAL	ILE	SEE REMARK 999	UNP Q59643
B	199	VAL	ILE	SEE REMARK 999	UNP Q59643
A	229	ARG	LYS	engineered mutation	UNP Q59643
B	229	ARG	LYS	engineered mutation	UNP Q59643

- Molecule 2 is FORMIC ACID (three-letter code: FMT) (formula: CH<sub>2</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 3 1 2	0	0
2	B	1	Total C O 3 1 2	0	0

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	2	Total Mg 2 2	0	0
3	A	3	Total Mg 3 3	0	0

- Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total Zn 1 1	0	0
4	A	1	Total Zn 1 1	0	0

- Molecule 5 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total Na 1 1	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Na	0	0
			1	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	304	Total	O	0	0
			304	304		
6	B	315	Total	O	0	0
			315	315		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DELTA-AMINOLEVULINIC ACID DEHYDRATASE



- Molecule 1: DELTA-AMINOLEVULINIC ACID DEHYDRATASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 4 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	125.67Å 125.67Å 85.52Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	87.71 – 1.40 19.35 – 1.40	Depositor EDS
% Data completeness (in resolution range)	97.2 (87.71-1.40) 97.3 (19.35-1.40)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.29 (at 1.40Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.147 , 0.173 0.159 , 0.184	Depositor DCC
$R_{free}$ test set	6524 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	14.5	Xtrriage
Anisotropy	0.232	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.42 , 70.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.35$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	6026	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	19.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.45% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: NA, MG, FMT, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	9.31	9/2749 (0.3%)	3.46	19/3720 (0.5%)
1	B	1.22	7/2733 (0.3%)	1.16	15/3699 (0.4%)
All	All	6.65	16/5482 (0.3%)	2.58	34/7419 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	1
All	All	0	3

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	335	ARG	CZ-NH2	484.52	7.62	1.33
1	A	273	LYS	CE-NZ	8.79	1.71	1.49
1	B	324	TYR	CE2-CZ	-6.72	1.29	1.38
1	A	289	TYR	CE2-CZ	-6.59	1.29	1.38
1	A	202	TYR	CE2-CZ	-5.91	1.30	1.38
1	B	47	GLY	N-CA	5.76	1.54	1.46
1	A	289	TYR	CG-CD1	-5.56	1.31	1.39
1	B	148	VAL	CB-CG1	-5.48	1.41	1.52
1	B	211	TYR	CE2-CZ	-5.44	1.31	1.38
1	A	283	TYR	CE2-CZ	-5.40	1.31	1.38
1	A	273	LYS	CD-CE	-5.36	1.37	1.51
1	A	210	TYR	CD2-CE2	-5.20	1.31	1.39
1	B	335	ARG	C-N	-5.20	1.23	1.33
1	B	283	TYR	CE1-CZ	-5.19	1.31	1.38

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	324	TYR	CE2-CZ	-5.18	1.31	1.38
1	B	211	TYR	CG-CD1	-5.01	1.32	1.39

All (34) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	335	ARG	NE-CZ-NH2	-198.84	20.88	120.30
1	A	335	ARG	NH1-CZ-NH2	-15.69	102.14	119.40
1	A	229	ARG	NE-CZ-NH1	9.86	125.23	120.30
1	A	25	ARG	NE-CZ-NH1	9.81	125.21	120.30
1	A	229	ARG	NE-CZ-NH2	-9.74	115.43	120.30
1	B	25	ARG	NE-CZ-NH1	8.17	124.39	120.30
1	A	51	ARG	NE-CZ-NH1	7.96	124.28	120.30
1	B	66	ASP	CB-CG-OD1	7.81	125.33	118.30
1	A	97	ASP	CB-CG-OD1	6.77	124.40	118.30
1	A	51	ARG	NE-CZ-NH2	-6.31	117.14	120.30
1	A	144	ASP	CB-CG-OD1	6.25	123.92	118.30
1	B	270[A]	ARG	NE-CZ-NH2	-6.04	117.28	120.30
1	B	270[B]	ARG	NE-CZ-NH2	-6.04	117.28	120.30
1	B	202	TYR	CB-CG-CD2	-5.93	117.44	121.00
1	B	277[A]	ARG	NE-CZ-NH2	-5.90	117.35	120.30
1	B	277[B]	ARG	NE-CZ-NH2	-5.90	117.35	120.30
1	A	202	TYR	CB-CG-CD2	-5.83	117.50	121.00
1	B	202	TYR	CB-CG-CD1	5.72	124.43	121.00
1	B	20	ARG	NE-CZ-NH1	5.64	123.12	120.30
1	A	20	ARG	NE-CZ-NH1	5.48	123.04	120.30
1	A	263[A]	MET	CG-SD-CE	-5.48	91.43	100.20
1	A	263[B]	MET	CG-SD-CE	-5.48	91.43	100.20
1	B	97	ASP	CB-CG-OD1	5.44	123.20	118.30
1	B	235	ASP	CB-CG-OD1	5.43	123.19	118.30
1	B	110	ARG	NE-CZ-NH2	5.35	122.97	120.30
1	A	291[A]	MET	CG-SD-CE	5.33	108.73	100.20
1	A	291[B]	MET	CG-SD-CE	5.33	108.73	100.20
1	A	20	ARG	NE-CZ-NH2	-5.31	117.64	120.30
1	B	198	ARG	NE-CZ-NH2	-5.30	117.65	120.30
1	B	26	ARG	NE-CZ-NH2	-5.28	117.66	120.30
1	B	51	ARG	NE-CZ-NH1	-5.27	117.66	120.30
1	A	17	ARG	NE-CZ-NH1	-5.07	117.76	120.30
1	A	270[A]	ARG	NE-CZ-NH2	-5.05	117.78	120.30
1	A	270[B]	ARG	NE-CZ-NH2	-5.05	117.78	120.30

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	19	ARG	Sidechain
1	A	335	ARG	Sidechain
1	B	19	ARG	Sidechain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2704	0	2645	37	0
1	B	2688	0	2640	31	0
2	A	3	0	1	5	0
2	B	3	0	1	0	0
3	A	3	0	0	0	0
3	B	2	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	A	304	0	0	13	0
6	B	315	0	0	15	0
All	All	6026	0	5287	63	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (63) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:273:LYS:NZ	1:A:273:LYS:CE	1.71	1.50
1:B:275[B]:GLU:HG2	6:B:2285:HOH:O	1.40	1.20
1:A:263[A]:MET:HG2	1:B:263[A]:MET:HG2	1.35	1.07
1:B:50[B]:GLN:OE1	6:B:2090:HOH:O	1.70	1.05
1:A:228:ASN:CA	6:A:2241:HOH:O	2.05	1.03
1:B:275[A]:GLU:OE2	6:B:2284:HOH:O	1.77	1.03
1:A:263[B]:MET:HE2	1:B:263[B]:MET:HE2	1.39	1.01

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:159[B]:ARG:NH1	6:A:2204:HOH:O	1.94	0.98
1:A:113[B]:ARG:NH2	6:A:2155:HOH:O	1.82	0.96
1:B:328[A]:GLN:OE1	6:B:2311:HOH:O	1.82	0.95
1:A:228:ASN:HA	6:A:2241:HOH:O	1.63	0.95
1:B:50[A]:GLN:OE1	6:B:2091:HOH:O	1.82	0.94
1:A:263[B]:MET:HE2	1:B:263[B]:MET:CE	1.99	0.92
1:A:263[B]:MET:CE	1:B:263[B]:MET:HE2	2.01	0.88
1:A:6:ALA:N	6:A:2002:HOH:O	2.10	0.85
1:B:275[B]:GLU:CG	6:B:2285:HOH:O	2.08	0.82
1:B:228:ASN:N	6:B:2252:HOH:O	2.11	0.82
1:B:50[B]:GLN:NE2	6:B:2090:HOH:O	2.11	0.81
1:A:113[B]:ARG:NE	6:A:2155:HOH:O	2.12	0.81
1:A:228:ASN:N	6:A:2241:HOH:O	2.07	0.80
1:B:228:ASN:N	6:B:2253:HOH:O	2.14	0.80
1:A:285:VAL:HB	2:A:1337:FMT:H	1.68	0.74
1:B:6:ALA:N	6:B:2001:HOH:O	2.25	0.69
1:A:285:VAL:HB	2:A:1337:FMT:C	2.22	0.69
1:B:50[B]:GLN:CD	6:B:2090:HOH:O	2.14	0.69
1:A:258[B]:MET:SD	1:A:322:LEU:HD13	2.34	0.68
1:B:258[B]:MET:SD	1:B:322:LEU:HD13	2.34	0.68
1:B:293[B]:MET:HE2	1:B:293[B]:MET:HA	1.78	0.66
1:A:74[A]:GLU:OE1	1:A:327:LYS:NZ	2.21	0.65
1:A:263[B]:MET:CE	1:B:263[B]:MET:CE	2.70	0.65
1:A:136[A]:HIS:HD2	1:A:138:GLN:H	1.44	0.63
1:A:136[A]:HIS:HE1	6:A:2136:HOH:O	1.81	0.62
1:A:110[B]:ARG:NH1	6:A:2154:HOH:O	2.35	0.59
1:A:286:SER:H	2:A:1337:FMT:C	2.16	0.58
1:A:273:LYS:NZ	1:A:273:LYS:CD	2.59	0.58
1:B:132:GLU:HG2	6:B:2196:HOH:O	2.04	0.57
1:A:89:THR:OG1	1:A:132[A]:GLU:HG3	2.05	0.56
1:A:143:ASP:HB3	1:A:149:LEU:HG	1.88	0.55
1:B:42[B]:VAL:HG11	1:B:68:LEU:HD13	1.89	0.54
1:B:50[A]:GLN:CD	6:B:2091:HOH:O	2.40	0.54
1:A:42[B]:VAL:HG11	1:A:68:LEU:CD1	2.38	0.52
1:B:229[A]:ARG:NE	6:B:2254:HOH:O	2.42	0.52
1:A:211:TYR:CE1	1:A:260:LYS:HE2	2.45	0.51
1:B:211:TYR:O	1:B:212:GLY:C	2.48	0.50
1:A:42[B]:VAL:HG11	1:A:68:LEU:HD13	1.95	0.49
1:B:213:PRO:HB2	1:B:286:SER:HB2	1.95	0.49
1:A:166[A]:GLU:OE1	6:A:2208:HOH:O	2.20	0.49
1:A:45:LEU:HD21	1:A:62[B]:ARG:HD3	1.96	0.48

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:42[A]:VAL:O	1:B:42[A]:VAL:HG23	2.14	0.48
1:A:113[B]:ARG:CZ	6:A:2155:HOH:O	2.26	0.47
1:A:285:VAL:CB	2:A:1337:FMT:H	2.42	0.47
1:B:289:TYR:CZ	1:B:293[A]:MET:HG3	2.52	0.45
1:A:35[B]:VAL:HG23	6:A:2070:HOH:O	2.16	0.45
1:B:73[B]:GLU:HG2	1:B:119:PHE:HZ	1.82	0.45
1:B:109:GLN:O	1:B:113[B]:ARG:HG3	2.16	0.44
1:B:229[A]:ARG:HG3	6:B:2255:HOH:O	2.17	0.44
1:A:32:VAL:HG12	6:A:2021:HOH:O	2.17	0.43
1:A:213:PRO:HB2	1:A:286:SER:HB2	2.02	0.42
1:A:143:ASP:HB3	1:A:149:LEU:CG	2.50	0.42
1:A:228:ASN:HD22	1:A:228:ASN:N	2.18	0.42
1:B:263[A]:MET:HB3	1:B:264:PRO:HD3	2.01	0.41
1:A:214:PHE:HB2	2:A:1337:FMT:O1	2.20	0.40
1:B:73[B]:GLU:CG	1:B:118[B]:ARG:HH21	2.35	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	340/337 (101%)	335 (98%)	5 (2%)	0	100	100
1	B	339/337 (101%)	330 (97%)	8 (2%)	1 (0%)	41	18
All	All	679/674 (101%)	665 (98%)	13 (2%)	1 (0%)	47	23

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	212	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	282/270 (104%)	277 (98%)	5 (2%)	59	28
1	B	281/270 (104%)	275 (98%)	6 (2%)	53	21
All	All	563/540 (104%)	552 (98%)	11 (2%)	60	23

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	181	ARG
1	A	198	ARG
1	A	228	ASN
1	A	283	TYR
1	A	335	ARG
1	B	50[A]	GLN
1	B	50[B]	GLN
1	B	181	ARG
1	B	229[A]	ARG
1	B	229[B]	ARG
1	B	283	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	298	ASN
1	B	298	ASN
1	B	332	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 9 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	FMT	A	1337	-	0,2,2	0.00	-	0,1,1	0.00	-
2	FMT	B	1337	-	0,2,2	0.00	-	0,1,1	0.00	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1337	FMT	5	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	335:ARG	C	336:GLY	N	8.36



## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	321/337 (95%)	-0.14	11 (3%) 45 44	10, 15, 31, 81	2 (0%)
1	B	319/337 (94%)	-0.16	11 (3%) 45 44	10, 15, 30, 87	4 (1%)
All	All	640/674 (94%)	-0.15	22 (3%) 45 44	10, 15, 31, 87	6 (0%)

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	336	GLY	14.6
1	B	214	PHE	11.3
1	A	336	GLY	10.8
1	A	214	PHE	10.5
1	B	335	ARG	7.0
1	A	212	GLY	6.3
1	A	216	ASP	6.1
1	B	215	ARG	5.9
1	A	215	ARG	4.7
1	B	144	ASP	4.4
1	B	212	GLY	4.2
1	B	213	PRO	3.7
1	A	217	ALA	3.4
1	A	7	ASN	3.3
1	A	335	ARG	3.2
1	A	213	PRO	2.9
1	B	334	ARG	2.9
1	A	334	ARG	2.6
1	B	145	ASP	2.4
1	B	7	ASN	2.3
1	B	6	ALA	2.2
1	A	6	ALA	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	MG	A	1340	1/1	0.92	0.08	17,17,17,17	1
3	MG	B	1339	1/1	0.94	0.18	20,20,20,20	1
3	MG	A	1341	1/1	0.95	0.20	11,11,11,11	1
2	FMT	B	1337	3/3	0.95	0.11	27,27,27,32	0
5	NA	A	1342	1/1	0.96	0.10	18,18,18,18	0
5	NA	B	1342	1/1	0.97	0.05	18,18,18,18	0
4	ZN	B	1340	1/1	0.98	0.04	15,15,15,15	1
2	FMT	A	1337	3/3	0.98	0.07	22,22,30,37	0
4	ZN	A	1339	1/1	0.99	0.04	15,15,15,15	1
3	MG	B	1338	1/1	1.00	0.07	8,8,8,8	1
3	MG	A	1338	1/1	1.00	0.03	10,10,10,10	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.