



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2018 – 09:08 pm GMT

PDB ID : 5W5U  
Title : Crystal structure of the A/Puerto Rico/8/1934 (H1N1) influenza virus hemagglutinin in complex with cyclic peptide CP141037 (P4)  
Authors : Wilson, I.A.; Kadam, R.U.  
Deposited on : 2017-06-15  
Resolution : 2.46 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.3 (157068), CSD as539be (2018)  
Xtriage (Phenix) : 1.13  
EDS : trunk30967  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk30967


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.46 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	1259 (2.48-2.44)
Clashscore	122126	1323 (2.48-2.44)
Ramachandran outliers	120053	1314 (2.48-2.44)
Sidechain outliers	120020	1314 (2.48-2.44)
RSRZ outliers	108989	1238 (2.48-2.44)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	326	
2	B	176	
3	D	12	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	NAG	A	401	-	-	-	X
4	NAG	A	403	-	-	-	X

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 4172 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hemagglutinin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	322	2542	1603	443	483	13	0	0	0


- Molecule 2 is a protein called Hemagglutinin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	171	1380	866	235	272	7	0	0	0

- Molecule 3 is a protein called ACE-PH8-ORN-MLE-GLU-TYR-PHE-GLU-TRP-LEU-SER-BAL.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	D	12	107	76	13	18	0	0	0

- Molecule 4 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	B	1	Total	C	N	O	0	0
			14	8	1	5		
4	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Cl	0	0
			1	1		
5	A	1	Total	Cl	0	0
			1	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	30	Total	O	0	0
			30	30		
6	B	26	Total	O	0	0
			26	26		
6	D	1	Total	O	0	0
			1	1		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Hemagglutinin



- Molecule 2: Hemagglutinin



- Molecule 3: ACE-PH8-ORN-MLE-GLU-TYR-PHE-GLU-TRP-LEU-SER-BAL



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 21 3	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	161.36Å 161.36Å 161.36Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	31.65 – 2.46 31.65 – 2.46	Depositor EDS
% Data completeness (in resolution range)	98.4 (31.65-2.46) 98.4 (31.65-2.46)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.08	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.34 (at 2.45Å)	Xtrriage
Refinement program	PHENIX 1.8.4_1496	Depositor
R, $R_{free}$	0.210 , 0.251 0.215 , 0.251	Depositor DCC
$R_{free}$ test set	1273 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	57.5	Xtrriage
Anisotropy	0.000	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 37.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.024 for -l,-k,-h	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	4172	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	68.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.05% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, ACE, MLE, CL, ORN, PH8, BAL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.28	0/2606	0.41	0/3544
2	B	0.29	0/1407	0.38	0/1891
3	D	3.16	12/72 (16.7%)	1.79	2/97 (2.1%)
All	All	0.50	12/4085 (0.3%)	0.46	2/5532 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	D	0	1

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	6	TYR	CB-CG	-9.04	1.38	1.51
3	D	9	TRP	CD2-CE2	-8.48	1.31	1.41
3	D	9	TRP	CG-CD2	-7.44	1.31	1.43
3	D	7	PHE	CB-CG	-7.41	1.38	1.51
3	D	10	LEU	CA-C	-7.13	1.34	1.52
3	D	8	GLU	CA-C	-6.62	1.35	1.52
3	D	6	TYR	CA-C	-6.52	1.35	1.52
3	D	11	SER	CA-C	-6.15	1.36	1.52
3	D	5	GLU	CA-C	-6.04	1.37	1.52
3	D	9	TRP	CA-C	-5.91	1.37	1.52
3	D	7	PHE	CA-C	-5.71	1.38	1.52
3	D	9	TRP	CD2-CE3	-5.69	1.31	1.40

All (2) bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	6	TYR	CB-CG-CD2	-6.17	117.30	121.00
3	D	6	TYR	CB-CG-CD1	6.09	124.66	121.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	D	11	SER	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2542	0	2470	18	0
2	B	1380	0	1309	10	0
3	D	107	0	95	6	0
4	A	56	0	52	0	0
4	B	28	0	25	1	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	A	30	0	0	0	0
6	B	26	0	0	1	0
6	D	1	0	0	0	0
All	All	4172	0	3951	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:50:ASN:O	2:B:54:THR:HG22	1.50	1.10
2:B:50:ASN:O	2:B:54:THR:CG2	2.08	1.01
3:D:2:PH8:CD1	3:D:2:PH8:HBA	2.05	0.85
1:A:125:PRO:O	1:A:126:SER:HB2	1.87	0.73
3:D:2:PH8:HD1	3:D:2:PH8:HBA	1.75	0.68

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:76:CYS:O	1:A:149:ARG:NH2	2.27	0.67
1:A:125(B):GLU:O	1:A:125(C):SER:HB2	1.98	0.64
2:B:147:GLU:OE2	4:B:202:NAG:N2	2.35	0.60
2:B:50:ASN:O	2:B:54:THR:HG23	2.00	0.58
1:A:310:ARG:NH1	2:B:90:ASP:OD1	2.31	0.55
1:A:180:TRP:HB3	1:A:254:PRO:HG3	1.90	0.52
1:A:119:GLU:HB2	1:A:259:ALA:HB3	1.92	0.51
1:A:216:GLU:O	1:A:220:ARG:NH2	2.42	0.51
3:D:2:PH8:CD1	3:D:2:PH8:CB	2.85	0.50
1:A:103:ILE:HG13	1:A:233:TYR:CE2	2.48	0.48
1:A:15:ILE:HD11	2:B:122:VAL:HG21	1.98	0.46
1:A:83:ARG:HB2	1:A:118:PHE:CE2	2.52	0.45
2:B:150:GLU:HG2	2:B:153:ARG:HH12	1.82	0.45
1:A:43:LEU:HB2	1:A:314:LEU:HB2	1.99	0.44
1:A:146:SER:OG	1:A:147:PHE:N	2.50	0.44
1:A:117:SER:OG	1:A:261:SER:HB2	2.18	0.43
2:B:128:ASN:OD1	2:B:159:TYR:OH	2.30	0.43
3:D:10:LEU:HD23	3:D:10:LEU:HA	1.81	0.42
3:D:3:ORN:HA	3:D:4:MLE:HN1	1.80	0.42
3:D:9:TRP:O	3:D:9:TRP:HE3	2.01	0.42
1:A:214:THR:HA	1:A:215:PRO:HD3	1.93	0.42
2:B:128:ASN:ND2	6:B:303:HOH:O	2.50	0.42
1:A:128:PRO:O	1:A:130:HIS:ND1	2.48	0.41
1:A:316:MET:HB2	1:A:316:MET:HE2	1.92	0.41
1:A:18:HIS:HB2	2:B:21:TRP:HA	2.03	0.41
1:A:276:GLU:H	1:A:276:GLU:HG3	1.75	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [\(i\)](#)

### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	320/326 (98%)	299 (93%)	19 (6%)	2 (1%)	27	32
2	B	169/176 (96%)	165 (98%)	4 (2%)	0	100	100
3	D	7/12 (58%)	7 (100%)	0	0	100	100
All	All	496/514 (96%)	471 (95%)	23 (5%)	2 (0%)	36	45

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	125(C)	SER
1	A	125(A)	LYS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	285/289 (99%)	277 (97%)	8 (3%)	47	59
2	B	147/151 (97%)	144 (98%)	3 (2%)	58	71
3	D	7/7 (100%)	7 (100%)	0	100	100
All	All	439/447 (98%)	428 (98%)	11 (2%)	50	63

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	92	ASN
1	A	102	PHE
1	A	125(C)	SER
1	A	126	SER
1	A	155	THR
1	A	156	GLU
1	A	261	SER
1	A	276	GLU
2	B	22	TYR
2	B	54	THR
2	B	128	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	BAL	D	12	3	4,4,5	1.75	1 (25%)	3,3,5	1.24	0
3	PH8	D	2	3	13,13,14	1.52	2 (15%)	12,15,17	0.85	0
3	ORN	D	3	3	7,7,8	3.87	2 (28%)	4,7,9	1.81	2 (50%)
3	MLE	D	4	3	8,8,9	3.39	1 (12%)	7,9,11	1.12	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	BAL	D	12	3	-	0/1/2/3	0/0/0/0
3	PH8	D	2	3	-	0/6/8/10	0/1/1/1
3	ORN	D	3	3	-	0/4/6/8	0/0/0/0
3	MLE	D	4	3	-	0/4/8/10	0/0/0/0

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	3	ORN	CA-C	-9.71	1.37	1.50
3	D	4	MLE	CA-C	-9.55	1.37	1.50
3	D	2	PH8	CJ-CG	-4.88	1.37	1.51

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	12	BAL	CA-C	-3.44	1.40	1.49
3	D	3	ORN	CB-CA	-3.05	1.49	1.53
3	D	2	PH8	CB-CA	-2.06	1.50	1.53

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	3	ORN	O-C-CA	-2.64	117.70	124.96
3	D	3	ORN	CB-CA-C	-2.23	107.59	111.85

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	2	PH8	3	0
3	D	3	ORN	1	0
3	D	4	MLE	1	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 2 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	NAG	A	401	1	14,14,15	1.05	0	17,19,21	1.29	2 (11%)
4	NAG	A	402	1	14,14,15	0.60	0	17,19,21	1.12	1 (5%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	NAG	A	403	1	14,14,15	0.27	0	17,19,21	0.68	0
4	NAG	A	404	1	14,14,15	1.06	1 (7%)	17,19,21	1.55	3 (17%)
4	NAG	B	201	2,4	14,14,15	0.62	0	17,19,21	1.63	4 (23%)
4	NAG	B	202	4	14,14,15	0.59	0	17,19,21	1.85	4 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	401	1	-	0/6/23/26	0/1/1/1
4	NAG	A	402	1	-	0/6/23/26	0/1/1/1
4	NAG	A	403	1	-	0/6/23/26	0/1/1/1
4	NAG	A	404	1	-	0/6/23/26	0/1/1/1
4	NAG	B	201	2,4	-	0/6/23/26	0/1/1/1
4	NAG	B	202	4	-	0/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	404	NAG	C2-N2	-2.13	1.42	1.46

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	201	NAG	C6-C5-C4	-4.71	101.86	112.99
4	B	202	NAG	C2-N2-C7	-4.59	116.25	122.94
4	B	202	NAG	C4-C3-C2	-3.04	106.56	111.02
4	A	404	NAG	C6-C5-C4	-2.85	106.25	112.99
4	A	401	NAG	O5-C1-C2	-2.66	107.85	111.52
4	A	404	NAG	O5-C1-C2	-2.66	107.85	111.52
4	B	202	NAG	C3-C4-C5	-2.54	105.69	110.24
4	B	201	NAG	O5-C1-C2	-2.43	108.17	111.52
4	B	201	NAG	O5-C5-C4	-2.37	105.05	110.83
4	A	402	NAG	O5-C1-C2	-2.29	108.37	111.52
4	B	201	NAG	O5-C5-C6	-2.11	103.80	107.15
4	A	401	NAG	C1-O5-C5	2.94	116.23	112.19
4	A	404	NAG	C1-O5-C5	3.19	116.57	112.19
4	B	202	NAG	O5-C5-C6	3.50	112.69	107.15

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	202	NAG	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	322/326 (98%)	0.01	9 (2%) 53 49	42, 65, 110, 147	0
2	B	171/176 (97%)	-0.10	3 (1%) 68 64	37, 58, 90, 123	0
3	D	7/12 (58%)	-0.80	0 100 100	55, 65, 74, 87	0
All	All	500/514 (97%)	-0.04	12 (2%) 59 55	37, 62, 107, 147	0

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	325	SER	9.4
2	B	61	ILE	4.3
2	B	63	PHE	4.2
1	A	92	ASN	3.8
1	A	125(B)	GLU	3.5
2	B	168	LEU	3.5
1	A	219	GLU	2.7
1	A	264	PHE	2.2
1	A	83(A)	SER	2.1
1	A	94	GLU	2.1
1	A	198	GLU	2.1
1	A	19	ALA	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	BAL	D	12	5/6	0.92	0.23	83,83,85,87	0

*Continued on next page...*



Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	PH8	D	2	13/14	0.95	0.24	55,61,77,77	0
3	ORN	D	3	8/9	0.95	0.09	53,55,69,77	0
3	MLE	D	4	9/10	0.97	0.17	42,53,62,63	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	NAG	A	402	14/15	0.75	0.32	101,114,120,121	0
4	NAG	A	403	14/15	0.76	0.51	133,141,149,150	0
4	NAG	B	202	14/15	0.78	0.37	123,128,133,134	0
4	NAG	A	401	14/15	0.78	0.54	136,146,157,161	0
4	NAG	B	201	14/15	0.81	0.28	103,105,114,123	0
4	NAG	A	404	14/15	0.81	0.38	113,124,132,133	0
5	CL	A	405	1/1	0.92	0.16	83,83,83,83	0
5	CL	B	203	1/1	0.98	0.07	65,65,65,65	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.