



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 14, 2018 – 02:29 am GMT

PDB ID : 4W63  
Title : TORPEDO CALIFORNICA ACETYLCHOLINESTERASE IN COMPLEX WITH A TACRINE-BENZOFURAN HYBRID INHIBITOR  
Authors : Pesaresi, A.; Samez, S.; Lamba, D.  
Deposited on : 2014-08-20  
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.3 (157068), CSD as539be (2018)  
Xtrriage (Phenix) : 1.13  
EDS : trunk31020  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk31020


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	2792 (2.80-2.80)
Clashscore	122126	3209 (2.80-2.80)
Ramachandran outliers	120053	3158 (2.80-2.80)
Sidechain outliers	120020	3160 (2.80-2.80)
RSRZ outliers	108989	2726 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	534	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	A	604	X	-	-	-

## 2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 4447 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Acetylcholinesterase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	534	4263	2731	724	786	22	39	0	0

- Molecule 2 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	14	8	1	5	0	0
2	A	1	14	8	1	5	0	0
2	A	1	14	8	1	5	0	0
2	A	1	14	8	1	5	0	0


- Molecule 3 is N-(1-benzofuran-2-ylmethyl)-N'-(1,2,3,4-tetrahydroacridin-9-yl)heptane-1,7-diamine (three-letter code: HTB) (formula: C<sub>29</sub>H<sub>35</sub>N<sub>3</sub>O).



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Acetylcholinesterase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	112.22Å 112.22Å 138.72Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	45.90 – 2.80 45.86 – 2.80	Depositor EDS
% Data completeness (in resolution range)	89.6 (45.90-2.80) 89.7 (45.86-2.80)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.16 (at 2.81Å)	Xtrriage
Refinement program	REFMAC 5.7.0032	Depositor
R, $R_{free}$	0.192 , 0.260 0.197 , 0.254	Depositor DCC
$R_{free}$ test set	1132 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	62.3	Xtrriage
Anisotropy	0.183	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 39.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.022 for -h,-k,l	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	4447	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	60.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.17% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, HTB

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.76	6/4387 (0.1%)	0.88	6/5955 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	286	SER	CA-CB	10.71	1.69	1.52
1	A	511	LYS	CG-CD	-10.08	1.18	1.52
1	A	454	LYS	CD-CE	-9.55	1.27	1.51
1	A	498	LYS	CG-CD	-8.42	1.23	1.52
1	A	192	LYS	CA-CB	-8.27	1.35	1.53
1	A	413	LYS	CG-CD	-5.11	1.35	1.52

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	486	HIS	CB-CA-C	-10.96	88.48	110.40
1	A	488	GLN	CB-CG-CD	6.99	129.78	111.60
1	A	518	VAL	CB-CA-C	-6.01	99.98	111.40
1	A	498	LYS	CB-CG-CD	5.76	126.57	111.60
1	A	511	LYS	CB-CG-CD	5.70	126.43	111.60
1	A	288	PHE	CB-CA-C	5.40	121.20	110.40

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	288	PHE	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4263	0	4106	74	0
2	A	56	0	51	1	0
3	A	33	0	35	7	0
4	A	95	0	0	16	0
All	All	4447	0	4192	80	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (80) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:288:PHE:HD1	1:A:289:ARG:HG2	1.45	0.81
1:A:175:MET:SD	4:A:719:HOH:O	2.41	0.77
1:A:287:ILE:HG12	1:A:331:PHE:CE2	2.22	0.75
2:A:603:NAG:H3	4:A:788:HOH:O	1.95	0.67
1:A:284:PHE:HA	1:A:285:ASP:HB3	1.80	0.63
3:A:605:HTB:H12	3:A:605:HTB:H15	1.82	0.61
1:A:127:LEU:HD12	1:A:130:TYR:CE2	2.36	0.61
3:A:605:HTB:H15	3:A:605:HTB:CAH	2.31	0.60
1:A:286:SER:O	4:A:701:HOH:O	2.17	0.60
1:A:321:LEU:HD11	1:A:408:VAL:HG22	1.84	0.59
1:A:221:ARG:HD3	1:A:480:GLY:HA2	1.85	0.59
1:A:284:PHE:HA	1:A:285:ASP:CB	2.33	0.58
3:A:605:HTB:H12	3:A:605:HTB:CAR	2.33	0.58
1:A:278:GLU:O	1:A:281:VAL:HG22	2.04	0.58
1:A:288:PHE:CD1	1:A:289:ARG:HG2	2.32	0.57
1:A:27:ILE:HD11	1:A:133:LYS:HB2	1.86	0.57

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:366:LEU:HD12	1:A:535:THR:HG21	1.85	0.56
1:A:200:SER:OG	3:A:605:HTB:NAW	2.39	0.55
1:A:520:MET:O	1:A:523:PHE:HB3	2.07	0.55
1:A:244:ARG:NH1	1:A:291:SER:O	2.34	0.54
1:A:88:ARG:HD2	4:A:706:HOH:O	2.09	0.53
1:A:515:ARG:HB3	1:A:518:VAL:HG22	1.91	0.52
1:A:453:VAL:HB	1:A:456:LEU:HD12	1.92	0.52
1:A:208:MET:HG3	1:A:229:PRO:HB3	1.91	0.52
1:A:432:TRP:HB3	1:A:433:PRO:HD2	1.93	0.51
1:A:276:ASP:O	1:A:279:TRP:NE1	2.38	0.51
1:A:45:PHE:O	1:A:164:ALA:HA	2.11	0.51
1:A:287:ILE:HG12	1:A:331:PHE:CZ	2.46	0.50
1:A:485:PRO:HA	4:A:734:HOH:O	2.12	0.50
1:A:255:ASN:HB3	1:A:261:GLU:HB3	1.93	0.49
1:A:199:GLU:HA	1:A:225:GLN:O	2.12	0.49
1:A:175:MET:CE	4:A:719:HOH:O	2.60	0.49
1:A:253:ASN:HA	4:A:754:HOH:O	2.11	0.49
1:A:475:THR:HG22	1:A:481:ASN:O	2.12	0.49
1:A:452:LEU:HD22	1:A:463:GLU:HG2	1.95	0.48
1:A:22:VAL:HG11	1:A:137:TYR:CB	2.43	0.48
3:A:605:HTB:H18	3:A:605:HTB:CBC	2.44	0.48
1:A:285:ASP:N	1:A:285:ASP:OD1	2.48	0.47
1:A:285:ASP:HB2	4:A:701:HOH:O	2.14	0.47
1:A:16:MET:HB2	1:A:57:VAL:HG13	1.97	0.47
3:A:605:HTB:CAR	3:A:605:HTB:CAH	2.93	0.47
1:A:48:PRO:HB2	4:A:719:HOH:O	2.15	0.46
1:A:233:TRP:CZ2	3:A:605:HTB:H35	2.50	0.46
1:A:197:PHE:HB2	1:A:223:ILE:HB	1.98	0.45
1:A:355:GLY:HA3	1:A:391:LEU:HD21	1.97	0.45
1:A:35:PHE:HD1	1:A:175:MET:HG2	1.80	0.45
1:A:109:THR:OG1	1:A:110:THR:N	2.49	0.45
1:A:228:SER:OG	1:A:230:ASN:OD1	2.34	0.45
1:A:333:LEU:HB2	1:A:436:MET:HE3	1.97	0.45
1:A:193:THR:O	1:A:193:THR:HG22	2.16	0.45
1:A:48:PRO:C	4:A:719:HOH:O	2.55	0.45
1:A:115:ILE:O	1:A:202:GLY:HA3	2.17	0.44
1:A:518:VAL:O	1:A:522:VAL:HG23	2.17	0.44
1:A:88:ARG:NE	4:A:706:HOH:O	2.49	0.44
1:A:515:ARG:CB	1:A:518:VAL:HG22	2.48	0.44
1:A:24:SER:C	4:A:711:HOH:O	2.56	0.43
1:A:75:PHE:CE1	1:A:341:LYS:HD3	2.53	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:177:LEU:HD21	1:A:196:ILE:HG22	2.01	0.43
1:A:248:LEU:HD12	1:A:277:VAL:HG23	2.01	0.43
1:A:69:GLN:HG3	4:A:770:HOH:O	2.17	0.43
1:A:197:PHE:CB	1:A:223:ILE:HB	2.49	0.43
1:A:251:ASN:OD1	1:A:280:ASN:ND2	2.52	0.43
1:A:98:ASN:O	1:A:144:VAL:HA	2.18	0.42
1:A:253:ASN:HB3	4:A:757:HOH:O	2.19	0.42
1:A:78:PHE:O	1:A:82:GLU:HB2	2.18	0.42
1:A:19:ARG:NH2	1:A:26:HIS:HB2	2.33	0.42
1:A:347:ILE:HG22	1:A:348:SER:O	2.19	0.42
1:A:502:PHE:CE1	1:A:516:LEU:HB2	2.55	0.42
1:A:287:ILE:CG1	1:A:335:GLY:HA3	2.50	0.41
1:A:240:GLU:O	1:A:244:ARG:HG3	2.21	0.41
1:A:339:PHE:CE2	1:A:391:LEU:HD23	2.55	0.41
1:A:505:LEU:HA	1:A:505:LEU:HD12	1.87	0.41
1:A:154:GLY:O	1:A:168:VAL:HG22	2.20	0.41
1:A:326:ASP:HB2	1:A:438:VAL:O	2.20	0.41
1:A:249:GLY:O	1:A:250:ARG:C	2.58	0.41
1:A:421:TYR:HB2	1:A:505:LEU:HD22	2.02	0.41
1:A:341:LYS:O	1:A:433:PRO:HG3	2.21	0.41
1:A:88:ARG:CD	4:A:706:HOH:O	2.66	0.40
1:A:132:GLY:HA3	1:A:143:LEU:HD13	2.03	0.40
1:A:175:MET:HE3	4:A:719:HOH:O	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	532/534 (100%)	481 (90%)	47 (9%)	4 (1%)	<b>21</b>   53

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	284	PHE
1	A	380	ASP
1	A	285	ASP
1	A	486	HIS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	467/467 (100%)	441 (94%)	26 (6%)	23 54

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	ASP
1	A	3	HIS
1	A	7	LEU
1	A	18	THR
1	A	25	SER
1	A	69	GLN
1	A	74	GLN
1	A	143	LEU
1	A	156	LEU
1	A	192	LYS
1	A	197	PHE
1	A	228	SER
1	A	285	ASP
1	A	287	ILE
1	A	295	VAL
1	A	299	GLU
1	A	321	LEU
1	A	349	ARG
1	A	350	GLU
1	A	353	MET
1	A	366	LEU
1	A	426	ARG
1	A	473	TRP

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	490	SER
1	A	511	LYS
1	A	518	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	74	GLN
1	A	251	ASN
1	A	280	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	A	601	1	14,14,15	0.71	0	17,19,21	2.61	3 (17%)
2	NAG	A	602	1,2	14,14,15	0.59	0	17,19,21	2.20	10 (58%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	A	603	2	14,14,15	0.88	1 (7%)	17,19,21	1.87	3 (17%)
2	NAG	A	604	1	14,14,15	1.06	1 (7%)	17,19,21	1.77	4 (23%)
3	HTB	A	605	-	33,37,37	2.07	8 (24%)	36,49,49	1.76	6 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	601	1	-	0/6/23/26	0/1/1/1
2	NAG	A	602	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	603	2	-	0/6/23/26	0/1/1/1
2	NAG	A	604	1	1/1/5/7	0/6/23/26	0/1/1/1
3	HTB	A	605	-	-	0/11/20/20	0/5/5/5

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	605	HTB	CAT-CBB	-5.98	1.41	1.51
3	A	605	HTB	CAS-CBA	-5.60	1.42	1.50
3	A	605	HTB	CAE-CBD	-3.98	1.32	1.42
3	A	605	HTB	CAI-CAZ	-3.05	1.33	1.39
3	A	605	HTB	CBE-NAV	-2.75	1.32	1.37
3	A	605	HTB	CBG-CBE	-2.43	1.38	1.42
2	A	603	NAG	C2-N2	-2.30	1.42	1.46
3	A	605	HTB	CAI-CBD	-2.24	1.33	1.41
3	A	605	HTB	CAU-CAZ	3.10	1.55	1.51
2	A	604	NAG	C1-C2	3.26	1.56	1.52

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	602	NAG	O5-C1-C2	-5.14	104.42	111.52
2	A	603	NAG	O5-C1-C2	-4.49	105.33	111.52
2	A	601	NAG	O5-C1-C2	-3.79	106.29	111.52
3	A	605	HTB	CAO-CAS-CBA	-3.74	107.22	113.55
2	A	604	NAG	O5-C5-C4	-3.37	102.64	110.83
3	A	605	HTB	CAL-CAN-CAR	-2.44	101.83	113.65
3	A	605	HTB	CAD-CAB-CAF	-2.44	117.03	120.44
2	A	602	NAG	O7-C7-C8	-2.36	117.80	122.07

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	602	NAG	O5-C5-C4	-2.26	105.33	110.83
2	A	601	NAG	C2-N2-C7	-2.24	119.67	122.94
2	A	602	NAG	C3-C4-C5	-2.15	106.39	110.24
2	A	602	NAG	O4-C4-C3	-2.13	105.37	110.34
2	A	604	NAG	O5-C1-C2	-2.01	108.75	111.52
2	A	602	NAG	C2-N2-C7	2.13	126.05	122.94
2	A	603	NAG	C4-C3-C2	2.39	114.53	111.02
2	A	602	NAG	O7-C7-N2	2.43	126.53	121.94
2	A	602	NAG	C6-C5-C4	2.69	119.35	112.99
2	A	602	NAG	C4-C3-C2	2.73	115.01	111.02
2	A	602	NAG	C1-O5-C5	2.81	116.06	112.19
2	A	603	NAG	C1-O5-C5	2.90	116.17	112.19
2	A	604	NAG	C2-N2-C7	3.08	127.43	122.94
2	A	604	NAG	C1-O5-C5	3.20	116.59	112.19
3	A	605	HTB	CAS-CBA-CBB	3.48	124.93	121.50
3	A	605	HTB	CAI-CBD-CBF	4.16	109.90	106.27
3	A	605	HTB	CBA-NAV-CBE	4.25	122.65	117.71
2	A	601	NAG	C1-O5-C5	9.17	124.80	112.19

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	604	NAG	C1

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	603	NAG	1	0
3	A	605	HTB	7	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	534/534 (100%)	-0.24	9 (1%) 70 63	34, 56, 88, 137	12 (2%)

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	2	ASP	7.3
1	A	3	HIS	5.4
1	A	255	ASN	2.9
1	A	535	THR	2.7
1	A	486	HIS	2.6
1	A	55	SER	2.1
1	A	264	HIS	2.1
1	A	257	ASN	2.1
1	A	266	LEU	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	NAG	A	603	14/15	0.78	0.38	77,90,107,118	0
2	NAG	A	604	14/15	0.86	0.29	69,91,107,107	0
2	NAG	A	601	14/15	0.87	0.19	74,99,106,107	0
2	NAG	A	602	14/15	0.92	0.15	56,73,81,89	0
3	HTB	A	605	33/33	0.98	0.19	40,47,55,56	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.