



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 6, 2023 – 08:28 PM EST

PDB ID : 5W6B
Title : Phosphotriesterase variant S1
Authors : Miton, C.M.; Campbell, E.C.; Jackson, C.J.; Tokuriki, N.
Deposited on : 2017-06-16
Resolution : 1.74 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.74 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3764 (1.76-1.72)
Clashscore	141614	3923 (1.76-1.72)
Ramachandran outliers	138981	3878 (1.76-1.72)
Sidechain outliers	138945	3878 (1.76-1.72)
RSRZ outliers	127900	3705 (1.76-1.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	333	
1	G	333	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5587 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phosphotriesterase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	329	2614	1644	470	492	8	0	13	0
1	G	330	2560	1616	455	482	7	0	6	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	254	SER	ARG	conflict	UNP A0A060GZX0
G	254	SER	ARG	conflict	UNP A0A060GZX0

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Zn	0	0
			2	2		
2	G	2	Total	Zn	0	0
			2	2		

- Molecule 3 is CACODYLATE ION (three-letter code: CAC) (formula: C₂H₆AsO₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	As	C	O		
3	A	1	5	1	2	2	0	0
3	G	1	5	1	2	2	0	0

- Molecule 4 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula: C₆H₁₄O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
4	A	1	8	6	2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			8	6	2		
4	G	1	Total	C	O	0	0
			8	6	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	198	Total	O	0	0
			198	198		
5	G	177	Total	O	0	0
			177	177		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Phosphotriesterase



- Molecule 1: Phosphotriesterase



4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	85.66Å 85.79Å 88.17Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.74 – 1.74 30.74 – 1.74	Depositor EDS
% Data completeness (in resolution range)	99.9 (30.74-1.74) 99.9 (30.74-1.74)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.71 (at 1.74Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.164 , 0.198 0.166 , 0.198	Depositor DCC
R_{free} test set	3409 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	17.4	Xtriage
Anisotropy	0.763	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 44.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.012 for -h,l,k 0.015 for -l,-k,-h 0.013 for k,h,-l 0.000 for k,l,h 0.000 for l,h,k	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5587	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.43% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CAC, KCX, MPD, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.39	0/2647	0.58	0/3593
1	G	0.38	0/2593	0.58	0/3519
All	All	0.39	0/5240	0.58	0/7112

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2614	0	2629	16	0
1	G	2560	0	2582	9	0
2	A	2	0	0	0	0
2	G	2	0	0	0	0
3	A	5	0	0	0	0
3	G	5	0	0	0	0
4	A	16	0	28	1	0
4	G	8	0	14	0	0
5	A	198	0	0	1	0
5	G	177	0	0	0	0
All	All	5587	0	5253	25	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:2404:MPD:O4	4:A:2404:MPD:O2	2.10	0.66
1:A:67[B]:ARG:HG2	1:G:159:GLU:HG2	1.81	0.62
1:A:110[B]:VAL:HG23	5:A:2577:HOH:O	2.00	0.61
1:A:361:THR:HG22	1:A:363:ARG:H	1.68	0.59
1:A:35:ASP:N	1:A:35:ASP:OD1	2.36	0.58
1:A:202[B]:THR:HG23	1:A:212:GLN:NE2	2.21	0.55
1:A:241[A]:THR:HG23	1:A:292:TYR:CE2	2.43	0.53
1:A:335:PHE:O	1:A:338:GLU:HG2	2.10	0.52
1:A:241[A]:THR:HG23	1:A:292:TYR:HE2	1.75	0.51
1:G:77:LYS:NZ	1:G:81:GLU:OE2	2.38	0.50
1:G:292:TYR:HA	1:G:294[B]:LYS:HE2	1.94	0.49
1:G:57:HIS:O	1:G:303:LEU:HA	2.15	0.47
1:G:300:ASN:OD1	1:G:328:ILE:HG12	2.15	0.46
1:A:302:TRP:CH2	1:A:321:ASN:HB3	2.50	0.46
1:A:110[B]:VAL:HG21	1:A:158:ILE:HG21	1.98	0.45
1:G:294[B]:LYS:HD3	1:G:294[B]:LYS:H	1.82	0.44
1:G:294[B]:LYS:H	1:G:294[B]:LYS:CD	2.29	0.44
1:A:175:LYS:HA	1:A:211:GLN:HE21	1.83	0.43
1:G:225:ARG:NH1	1:G:357:PHE:O	2.52	0.43
1:A:362:LEU:HA	1:A:362:LEU:HD23	1.84	0.42
1:A:353:ASN:HB2	1:A:354:PRO:HD3	2.01	0.41
1:G:302:TRP:CH2	1:G:321:ASN:HB3	2.56	0.41
1:A:333:ILE:HB	1:A:334:PRO:HD3	2.02	0.41
1:A:57:HIS:O	1:A:303:LEU:HA	2.21	0.41
1:A:175:LYS:HA	1:A:175:LYS:HD3	1.90	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	339/333 (102%)	327 (96%)	11 (3%)	1 (0%)	41	23
1	G	333/333 (100%)	322 (97%)	10 (3%)	1 (0%)	41	23
All	All	672/666 (101%)	649 (97%)	21 (3%)	2 (0%)	41	23

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	G	362	LEU
1	A	176	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	277/266 (104%)	270 (98%)	7 (2%)	47	24
1	G	270/266 (102%)	265 (98%)	5 (2%)	57	36
All	All	547/532 (103%)	535 (98%)	12 (2%)	53	29

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	51	PHE
1	A	260	ILE
1	A	263	GLU
1	A	293[A]	MET
1	A	293[B]	MET
1	A	299	SER
1	A	303	LEU
1	G	51	PHE
1	G	206	GLN
1	G	299	SER
1	G	303	LEU
1	G	306	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	KCX	G	169	1,2	9,11,12	1.52	1 (11%)	5,12,14	2.18	1 (20%)
1	KCX	A	169	1,2	9,11,12	1.42	2 (22%)	5,12,14	1.98	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	KCX	G	169	1,2	-	0/9/10/12	-
1	KCX	A	169	1,2	-	0/9/10/12	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	G	169	KCX	CX-NZ	3.64	1.41	1.35
1	A	169	KCX	CX-NZ	2.69	1.39	1.35
1	A	169	KCX	OQ1-CX	2.67	1.26	1.21

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	G	169	KCX	OQ1-CX-NZ	-4.77	117.56	124.96
1	A	169	KCX	OQ1-CX-NZ	-4.04	118.69	124.96

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 4 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	MPD	A	2404	-	7,7,7	0.54	0	9,10,10	0.41	0
3	CAC	A	2403	2	0,4,4	-	-	0,6,6	-	-
4	MPD	G	2404	-	7,7,7	0.55	0	9,10,10	0.94	1 (11%)
3	CAC	G	2403	2	0,4,4	-	-	0,6,6	-	-
4	MPD	A	2405	-	7,7,7	0.55	0	9,10,10	0.56	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MPD	A	2404	-	-	3/5/5/5	-
4	MPD	A	2405	-	-	1/5/5/5	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MPD	G	2404	-	-	3/5/5/5	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	G	2404	MPD	CM-C2-C1	-2.60	105.16	110.57

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	2404	MPD	O2-C2-C3-C4
4	A	2405	MPD	O2-C2-C3-C4
4	G	2404	MPD	O2-C2-C3-C4
4	A	2404	MPD	CM-C2-C3-C4
4	G	2404	MPD	C1-C2-C3-C4
4	G	2404	MPD	CM-C2-C3-C4
4	A	2404	MPD	C2-C3-C4-O4

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	2404	MPD	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	328/333 (98%)	0.27	23 (7%) 16 21	10, 20, 41, 64	0
1	G	329/333 (98%)	0.17	9 (2%) 54 60	14, 29, 44, 67	0
All	All	657/666 (98%)	0.22	32 (4%) 29 34	10, 25, 43, 67	0

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	362	LEU	10.1
1	A	260	ILE	7.8
1	A	261	GLY	7.6
1	A	274	ILE	6.9
1	A	264	ASP	6.8
1	A	266	ALA	6.4
1	A	262	LEU	6.3
1	A	263	GLU	6.1
1	G	363	ARG	5.8
1	A	35	ASP	5.1
1	A	363	ARG	4.7
1	A	265	ASN	4.3
1	A	362	LEU	4.2
1	A	270	ALA	3.9
1	A	269	SER	3.8
1	A	268	ALA	3.8
1	A	267	SER	3.3
1	G	361	THR	3.2
1	A	259	ALA	3.0
1	A	272	LEU	2.9
1	A	175	LYS	2.7
1	A	338	GLU	2.7
1	A	173	THR	2.7
1	A	174	GLY	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	G	337	ARG	2.4
1	G	173	THR	2.3
1	G	101	VAL	2.3
1	A	46	ILE	2.2
1	G	175	LYS	2.1
1	A	271	LEU	2.1
1	G	35	ASP	2.0
1	G	252	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	KCX	G	169	12/13	0.95	0.14	19,20,23,23	0
1	KCX	A	169	12/13	0.96	0.11	10,12,14,15	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	MPD	G	2404	8/8	0.48	0.37	34,39,47,50	8
4	MPD	A	2404	8/8	0.58	0.28	34,42,47,50	8
4	MPD	A	2405	8/8	0.79	0.23	32,40,47,48	0
2	ZN	G	2401	1/1	0.99	0.06	22,22,22,22	0
3	CAC	A	2403	5/5	0.99	0.11	14,14,17,18	5
3	CAC	G	2403	5/5	0.99	0.07	22,23,26,30	5
2	ZN	G	2402	1/1	1.00	0.05	23,23,23,23	0
2	ZN	A	2402	1/1	1.00	0.06	13,13,13,13	1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	ZN	A	2401	1/1	1.00	0.06	13,13,13,13	0

6.5 Other polymers [i](#)

There are no such residues in this entry.