



Full wwPDB X-ray Structure Validation Report ⓘ

May 25, 2020 – 08:16 pm BST

PDB ID : 5W6L
Title : Crystal Structure of RRSP, a MARTX Toxin Effector Domain from *Vibrio vulnificus* CMCP6
Authors : Minasov, G.; Wawrzak, Z.; Biancucci, M.; Shuvalova, L.; Dubrovskaya, I.; Satchell, K.J.; Anderson, W.F.; Center for Structural Genomics of Infectious Diseases (CSGID)
Deposited on : 2017-06-16
Resolution : 3.45 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1291 (3.52-3.40)
Clashscore	141614	1372 (3.52-3.40)
Ramachandran outliers	138981	1337 (3.52-3.40)
Sidechain outliers	138945	1338 (3.52-3.40)
RSRZ outliers	127900	1205 (3.52-3.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	518	 3% 90% 6% 5%
1	B	518	 4% 87% 8% 5%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	A	4101	-	-	-	X
3	CL	B	4101	-	-	-	X
4	SO4	B	4102	-	-	-	X
4	SO4	B	4103	-	-	-	X

2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 7694 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RTX repeat-containing cytotoxin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	Se			
1	A	494	3835	2404	667	752	12	0	0	0
1	B	492	3827	2396	667	752	12	0	1	0

There are 52 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	3572	MSE	-	initiating methionine	UNP A0A1V8MQR3
A	3573	HIS	-	expression tag	UNP A0A1V8MQR3
A	3574	HIS	-	expression tag	UNP A0A1V8MQR3
A	3575	HIS	-	expression tag	UNP A0A1V8MQR3
A	3576	HIS	-	expression tag	UNP A0A1V8MQR3
A	3577	HIS	-	expression tag	UNP A0A1V8MQR3
A	3578	HIS	-	expression tag	UNP A0A1V8MQR3
A	3579	SER	-	expression tag	UNP A0A1V8MQR3
A	3580	SER	-	expression tag	UNP A0A1V8MQR3
A	3581	GLY	-	expression tag	UNP A0A1V8MQR3
A	3582	VAL	-	expression tag	UNP A0A1V8MQR3
A	3583	ASP	-	expression tag	UNP A0A1V8MQR3
A	3584	LEU	-	expression tag	UNP A0A1V8MQR3
A	3585	GLY	-	expression tag	UNP A0A1V8MQR3
A	3586	THR	-	expression tag	UNP A0A1V8MQR3
A	3587	GLU	-	expression tag	UNP A0A1V8MQR3
A	3588	ASN	-	expression tag	UNP A0A1V8MQR3
A	3589	LEU	-	expression tag	UNP A0A1V8MQR3
A	3590	TYR	-	expression tag	UNP A0A1V8MQR3
A	3591	PHE	-	expression tag	UNP A0A1V8MQR3
A	3592	GLN	-	expression tag	UNP A0A1V8MQR3
A	3593	SER	-	expression tag	UNP A0A1V8MQR3
A	3594	ASN	-	expression tag	UNP A0A1V8MQR3
A	3595	ALA	-	expression tag	UNP A0A1V8MQR3
A	3695	ILE	LEU	conflict	UNP A0A1V8MQR3

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	4087	GLY	ASP	conflict	UNP A0A1V8MQR3
B	3572	MSE	-	initiating methionine	UNP A0A1V8MQR3
B	3573	HIS	-	expression tag	UNP A0A1V8MQR3
B	3574	HIS	-	expression tag	UNP A0A1V8MQR3
B	3575	HIS	-	expression tag	UNP A0A1V8MQR3
B	3576	HIS	-	expression tag	UNP A0A1V8MQR3
B	3577	HIS	-	expression tag	UNP A0A1V8MQR3
B	3578	HIS	-	expression tag	UNP A0A1V8MQR3
B	3579	SER	-	expression tag	UNP A0A1V8MQR3
B	3580	SER	-	expression tag	UNP A0A1V8MQR3
B	3581	GLY	-	expression tag	UNP A0A1V8MQR3
B	3582	VAL	-	expression tag	UNP A0A1V8MQR3
B	3583	ASP	-	expression tag	UNP A0A1V8MQR3
B	3584	LEU	-	expression tag	UNP A0A1V8MQR3
B	3585	GLY	-	expression tag	UNP A0A1V8MQR3
B	3586	THR	-	expression tag	UNP A0A1V8MQR3
B	3587	GLU	-	expression tag	UNP A0A1V8MQR3
B	3588	ASN	-	expression tag	UNP A0A1V8MQR3
B	3589	LEU	-	expression tag	UNP A0A1V8MQR3
B	3590	TYR	-	expression tag	UNP A0A1V8MQR3
B	3591	PHE	-	expression tag	UNP A0A1V8MQR3
B	3592	GLN	-	expression tag	UNP A0A1V8MQR3
B	3593	SER	-	expression tag	UNP A0A1V8MQR3
B	3594	ASN	-	expression tag	UNP A0A1V8MQR3
B	3595	ALA	-	expression tag	UNP A0A1V8MQR3
B	3695	ILE	LEU	conflict	UNP A0A1V8MQR3
B	4087	GLY	ASP	conflict	UNP A0A1V8MQR3

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Cl	0	0
			1	1		

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	11	Total	O	0	0
			11	11		
5	B	4	Total	O	0	0
			4	4		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: RTX repeat-containing cytotoxin



- Molecule 1: RTX repeat-containing cytotoxin



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 3	Depositor
Cell constants a, b, c, α , β , γ	247.10Å 247.10Å 247.10Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.97 – 3.45 29.97 – 3.45	Depositor EDS
% Data completeness (in resolution range)	98.6 (29.97-3.45) 98.8 (29.97-3.45)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	0.11	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.72 (at 3.47Å)	Xtrriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.228 , 0.249 0.232 , 0.257	Depositor DCC
R_{free} test set	1562 reflections (4.79%)	wwPDB-VP
Wilson B-factor (Å ²)	101.9	Xtrriage
Anisotropy	0.000	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 60.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.000 for -l,-k,-h	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	7694	wwPDB-VP
Average B, all atoms (Å ²)	124.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.66% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, SO4, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.30	0/3888	0.52	0/5232
1	B	0.31	0/3878	0.52	2/5217 (0.0%)
All	All	0.31	0/7766	0.52	2/10449 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	3866	GLY	N-CA-C	7.14	130.96	113.10
1	B	3865	SER	N-CA-CB	-5.96	101.56	110.50

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3835	0	3787	16	0
1	B	3827	0	3777	22	0
2	A	6	0	8	0	0
3	B	1	0	0	0	0
4	B	10	0	0	0	0
5	A	11	0	0	0	0
5	B	4	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	7694	0	7572	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:3674:MSE:HE1	1:B:3699:ALA:HB3	1.79	0.65
1:A:3702:GLU:HA	1:A:3706:LYS:O	2.02	0.59
1:B:3696:PHE:CE2	1:B:3700:ILE:HD11	2.38	0.59
1:A:3896:LEU:HD11	1:A:4023:VAL:HG13	1.85	0.59
1:A:3748:LEU:HD11	1:A:3780:VAL:HG12	1.85	0.58
1:A:3687:THR:HG21	1:A:3726:PRO:HG3	1.88	0.54
1:A:3748:LEU:N	1:A:3748:LEU:HD12	2.23	0.54
1:B:3782:LEU:HD21	1:B:3789:LEU:HD11	1.90	0.54
1:B:3943:ARG:O	1:B:3947:THR:HG22	2.08	0.54
1:B:4082:LYS:HG3	1:B:4084:THR:HG23	1.90	0.53
1:B:3879:ALA:HB1	1:B:3884:GLU:HB3	1.91	0.53
1:B:3991:VAL:HG23	1:B:3994:THR:HG23	1.92	0.52
1:B:3848:ILE:HG21	1:B:3999:MSE:HE2	1.91	0.51
1:A:3906:VAL:HG12	1:A:3906:VAL:O	2.11	0.50
1:B:3906:VAL:HG12	1:B:3906:VAL:O	2.12	0.50
1:B:3898:ILE:HD12	1:B:3898:ILE:N	2.27	0.49
1:B:3748:LEU:N	1:B:3748:LEU:HD12	2.28	0.48
1:B:3674:MSE:HE1	1:B:3699:ALA:CB	2.42	0.48
1:A:3925:THR:O	1:A:3925:THR:HG22	2.16	0.46
1:A:3928:GLY:C	1:A:3929:LEU:HD12	2.36	0.46
1:A:3802:PHE:CE2	1:A:3806:LYS:HD2	2.51	0.46
1:B:3872:LEU:HD22	1:B:4063:VAL:HG21	1.97	0.45
1:A:3876:LEU:HD12	1:A:3877:ALA:N	2.32	0.45
1:B:3844:GLN:HE21	1:B:3985:SER:CB	2.30	0.44
1:B:3991:VAL:CG2	1:B:3994:THR:HG23	2.48	0.44
1:A:3691:LEU:HD11	1:A:3842:MSE:HE2	1.99	0.43
1:A:3844:GLN:HE21	1:A:3985:SER:CB	2.31	0.43
1:B:3929:LEU:H	1:B:3981:LEU:HD13	1.82	0.43
1:B:3996:HIS:CD2	1:B:4037:ILE:HG22	2.53	0.43
1:B:3920:LYS:HB2	1:B:3976:MSE:CE	2.49	0.42
1:B:3848:ILE:HG21	1:B:3999:MSE:CE	2.49	0.42
1:B:3896:LEU:HD13	1:B:3898:ILE:HD11	1.99	0.42
1:B:3746:VAL:HG21	1:B:3805:LEU:HD11	2.01	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:3674:MSE:HE1	1:A:3699:ALA:HB3	2.02	0.41
1:A:3848:ILE:HG21	1:A:3999:MSE:HE2	2.03	0.41
1:A:3712:ASP:OD2	1:A:4082:LYS:HG3	2.20	0.41
1:A:3724:LEU:HD12	1:A:3724:LEU:N	2.36	0.41
1:B:3717:TRP:CE2	1:B:3845:ILE:HD11	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	492/518 (95%)	467 (95%)	25 (5%)	0	100	100
1	B	489/518 (94%)	460 (94%)	29 (6%)	0	100	100
All	All	981/1036 (95%)	927 (94%)	54 (6%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	406/415 (98%)	403 (99%)	3 (1%)	84	93
1	B	406/415 (98%)	403 (99%)	3 (1%)	84	93
All	All	812/830 (98%)	806 (99%)	6 (1%)	84	93

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3629	ARG
1	A	3688	ARG
1	A	4026	TYR
1	B	3607	LYS
1	B	3957	MSE
1	B	4026	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 1 is monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	A	4101	-	5,5,5	0.29	0	5,5,5	0.22	0
4	SO4	B	4103	-	4,4,4	0.33	0	6,6,6	0.05	0
4	SO4	B	4102	-	4,4,4	0.32	0	6,6,6	0.04	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	4101	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	482/518 (93%)	0.05	16 (3%) 46 44	91, 114, 152, 222	0
1	B	480/518 (92%)	0.21	21 (4%) 34 33	93, 124, 185, 220	0
All	All	962/1036 (92%)	0.13	37 (3%) 40 39	91, 118, 178, 222	0

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	3604	VAL	5.5
1	B	3603	LYS	5.1
1	A	3603	LYS	4.5
1	A	3604	VAL	4.4
1	A	3608	PRO	4.4
1	A	3602	ALA	4.4
1	A	3605	PHE	3.9
1	B	3594	ASN	3.7
1	B	3602	ALA	3.7
1	A	3815	ILE	3.5
1	B	3609	ILE	3.5
1	B	3595	ALA	3.3
1	B	3596	GLN	3.3
1	A	3595	ALA	3.2
1	B	3599	LYS	3.2
1	B	3600	GLU	3.2
1	A	3596	GLN	3.1
1	B	3597	GLU	2.8
1	B	3608	PRO	2.8
1	B	3877	ALA	2.8
1	A	3601	ARG	2.8
1	A	3606	ALA	2.8
1	A	3599	LYS	2.8
1	A	3611	ALA	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	4019	GLY	2.7
1	A	3607	LYS	2.5
1	B	3607	LYS	2.5
1	A	3594	ASN	2.4
1	B	3601	ARG	2.4
1	B	3598	LEU	2.4
1	A	3600	GLU	2.4
1	A	4087	GLY	2.4
1	B	4024	ALA	2.3
1	B	4018	ASP	2.2
1	B	3733	GLU	2.2
1	B	3878	SER	2.0
1	B	3625	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	GOL	A	4101	6/6	0.59	0.49	134,134,136,136	0
3	CL	B	4101	1/1	0.62	0.67	132,132,132,132	0
4	SO4	B	4103	5/5	0.69	0.41	222,224,224,226	0
4	SO4	B	4102	5/5	0.75	0.58	186,187,188,190	0

6.5 Other polymers [i](#)

There are no such residues in this entry.