



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 13, 2018 – 03:24 pm GMT

PDB ID : 2W79
Title : Establishing wild-type levels of catalytic activity on natural and artificial (ba)8-barrel protein scaffolds
Authors : Claren, J.; Malisi, C.; Hocker, B.; Sterner, R.
Deposited on : 2008-12-21
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtrriage (Phenix) : 1.13
EDS : trunk31020
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk31020


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	2111 (1.86-1.86)
Clashscore	122126	2258 (1.86-1.86)
Ramachandran outliers	120053	2234 (1.86-1.86)
Sidechain outliers	120020	2234 (1.86-1.86)
RSRZ outliers	108989	2075 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	241	
1	B	241	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 4182 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 1-(5-PHOSPHORIBOSYL)-5-[(5-PHOSPHORIBOSYLAMIN O) METHYLIDENEAMINO] IMIDAZOLE-4-CARBOXAMIDE ISOMERASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	241	1918	1228	327	360	3	0	4	0
1	B	239	1885	1208	321	353	3	0	3	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	75	TYR	HIS	engineered mutation	UNP Q9X0C7
A	111	SER	PHE	engineered mutation	UNP Q9X0C7
A	127	VAL	ASP	engineered mutation	UNP Q9X0C7
A	169	VAL	ASP	engineered mutation	UNP Q9X0C7
B	75	TYR	HIS	engineered mutation	UNP Q9X0C7
B	111	SER	PHE	engineered mutation	UNP Q9X0C7
B	127	VAL	ASP	engineered mutation	UNP Q9X0C7
B	169	VAL	ASP	engineered mutation	UNP Q9X0C7

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Cl	0	0
			1	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	159	Total	O	0	0
			159	159		
3	B	219	Total	O	0	0
			219	219		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 1-(5-PHOSPHORIBOSYL)-5-[(5-PHOSPHORIBOSYLAMINO) METHYLIDENEAMINO] IMIDAZOLE-4-CARBOXAMIDE ISOMERASE



- Molecule 1: 1-(5-PHOSPHORIBOSYL)-5-[(5-PHOSPHORIBOSYLAMINO) METHYLIDENEAMINO] IMIDAZOLE-4-CARBOXAMIDE ISOMERASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	49.00Å 64.10Å 79.20Å 90.00° 94.00° 90.00°	Depositor
Resolution (Å)	33.63 – 1.85 33.63 – 1.85	Depositor EDS
% Data completeness (in resolution range)	100.0 (33.63-1.85) 99.2 (33.63-1.85)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.20 (at 1.85Å)	Xtrriage
Refinement program	REFMAC 5.4.0066	Depositor
R, R_{free}	0.202 , 0.267 0.202 , 0.268	Depositor DCC
R_{free} test set	2082 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	26.6	Xtrriage
Anisotropy	0.243	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 42.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4182	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 47.07 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.0460e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.53	0/1955	0.63	0/2629
1	B	0.58	0/1918	0.65	0/2579
All	All	0.55	0/3873	0.64	0/5208

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1918	0	2014	17	0
1	B	1885	0	1973	9	0
2	B	1	0	0	0	0
3	A	159	0	0	3	0
3	B	219	0	0	4	0
All	All	4182	0	3987	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:129:ARG:NE	1:B:167:GLU:OE2	2.21	0.70
1:A:28:TYR:CE1	1:A:227:LEU:HD23	2.34	0.63
1:B:38:LYS:HD3	3:B:2008:HOH:O	1.99	0.62
1:A:113:LYS:HD2	1:A:116[B]:ARG:HH21	1.66	0.61
1:B:65:VAL:HG23	3:B:2078:HOH:O	2.02	0.60
1:B:160:GLU:HG3	1:B:190:LYS:HB2	1.83	0.58
1:A:38:LYS:O	1:A:42:GLU:HG2	2.04	0.58
1:A:208:LYS:O	1:A:212:GLU:HG3	2.04	0.58
1:A:113:LYS:HD2	1:A:116[B]:ARG:NH2	2.19	0.57
1:A:48:HIS:CE1	1:A:100:ILE:HD13	2.44	0.52
1:A:116[A]:ARG:NH2	3:A:2084:HOH:O	2.36	0.52
1:A:136:LYS:HG3	3:A:2099:HOH:O	2.10	0.51
1:B:85:LEU:HD11	1:B:114:SER:HB3	1.94	0.49
1:B:177:PHE:HB3	1:B:209:VAL:HG21	1.95	0.48
1:A:1:MET:HB2	1:A:1:MET:HE2	1.56	0.48
1:B:2:LEU:HA	3:B:2002:HOH:O	2.13	0.47
1:A:213:THR:HB	1:A:216:LEU:HB2	1.96	0.46
1:A:154:LYS:HD2	1:A:187:ALA:O	2.16	0.46
1:A:57:GLU:O	1:A:58:ASN:HB2	2.16	0.45
1:A:205:THR:O	1:A:209:VAL:HG23	2.17	0.45
1:A:126:LEU:HD21	1:A:146:PRO:HB3	1.97	0.45
1:A:30:LYS:HD3	1:A:35:LEU:HD22	1.99	0.44
1:A:228:GLU:HG3	3:A:2153:HOH:O	2.17	0.44
1:B:120:VAL:O	1:B:122:PRO:HD3	2.19	0.43
1:B:116[A]:ARG:NH2	3:B:2143:HOH:O	2.52	0.41
1:A:208:LYS:HB2	1:A:208:LYS:HE3	1.68	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	243/241 (101%)	237 (98%)	6 (2%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	238/241 (99%)	231 (97%)	7 (3%)	0	100	100
All	All	481/482 (100%)	468 (97%)	13 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	211/208 (101%)	206 (98%)	5 (2%)	52	35
1	B	206/208 (99%)	203 (98%)	3 (2%)	67	56
All	All	417/416 (100%)	409 (98%)	8 (2%)	58	45

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	11	ARG
1	A	106	LEU
1	A	136	LYS
1	A	142	GLU
1	A	178	SER
1	B	83	ARG
1	B	145	ASP
1	B	214	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	241/241 (100%)	0.15	7 (2%) 51 50	14, 24, 33, 45	0
1	B	239/241 (99%)	0.10	8 (3%) 46 45	14, 24, 33, 39	0
All	All	480/482 (99%)	0.12	15 (3%) 49 47	14, 24, 33, 45	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	241	ARG	4.4
1	A	214	ASN	3.9
1	B	144	ILE	3.8
1	B	208	LYS	2.9
1	B	214	ASN	2.8
1	A	211	THR	2.7
1	B	241	ARG	2.5
1	B	213	THR	2.5
1	A	208	LYS	2.3
1	B	211	THR	2.3
1	A	144	ILE	2.3
1	B	212	GLU	2.2
1	A	221	ILE	2.2
1	A	169	VAL	2.1
1	B	221	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	CL	B	1242	1/1	0.99	0.03	32,32,32,32	0

6.5 Other polymers [i](#)

There are no such residues in this entry.