



# wwPDB EM Validation Summary Report ⓘ

May 6, 2024 – 04:18 AM JST

PDB ID : 8W9F  
EMDB ID : EMD-37367  
Title : Cryo-EM structure of the Rpd3S-nucleosome complex from budding yeast in State 3  
Authors : Wang, C.; Zhan, X.  
Deposited on : 2023-09-05  
Resolution : 4.40 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev92  
MolProbity : 4.02b-467  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
MapQ : 1.9.13  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36.2

# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:  
*ELECTRON MICROSCOPY*

The reported resolution of this entry is 4.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1536	
2	E	684	
2	F	684	
3	B	433	
4	C	401	
4	D	401	
4	G	401	
5	a	136	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
5	e	136	68% 27%
6	b	103	74% 23%
6	f	103	77% 20%
7	c	130	78% 18%
7	g	130	78% 18%
8	d	126	70% 25%
8	h	126	71% 25%
9	i	147	100%
10	j	147	99%

## 2 Entry composition [i](#)

There are 11 unique types of molecules in this entry. The entry contains 27868 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transcriptional regulatory protein SIN3.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	606	5066	3253	860	938	15	0	0

- Molecule 2 is a protein called Transcriptional regulatory protein RCO1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	E	353	2884	1837	493	536	18	0	0
2	F	156	1282	822	211	239	10	0	0

- Molecule 3 is a protein called Histone deacetylase RPD3.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
3	B	385	3057	1948	513	571	25	0	0

- Molecule 4 is a protein called Chromatin modification-related protein EAF3.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
4	C	183	1483	950	239	285	9	0	0
4	D	185	1497	959	241	288	9	0	0
4	G	69	570	371	98	97	4	0	0

- Molecule 5 is a protein called Histone H3.1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
5	a	97	801	505	155	137	4	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					AltConf	Trace
5	e	99	Total	C	N	O	S	0	0
			816	514	158	140	4		

- Molecule 6 is a protein called Histone H4.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	b	79	Total	C	N	O	S	0	0
			627	395	121	110	1		
6	f	82	Total	C	N	O	S	0	0
			653	412	127	113	1		

- Molecule 7 is a protein called Histone H2A type 1-B/E.

Mol	Chain	Residues	Atoms				AltConf	Trace
7	c	107	Total	C	N	O	0	0
			817	516	158	143		
7	g	107	Total	C	N	O	0	0
			817	516	158	143		

- Molecule 8 is a protein called Histone H2B type 1-K.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	d	94	Total	C	N	O	S	0	0
			735	461	134	138	2		
8	h	94	Total	C	N	O	S	0	0
			735	461	134	138	2		

- Molecule 9 is a DNA chain called 5-DNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
9	i	147	Total	C	N	O	P	0	0
			3011	1440	546	879	146		

- Molecule 10 is a DNA chain called 3-DNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
10	j	147	Total	C	N	O	P	0	0
			3010	1440	543	881	146		

- Molecule 11 is ZINC ION (three-letter code: ZN) (formula: Zn).


<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>AltConf</b>
11	E	4	Total 4	Zn 4	0
11	B	1	Total 1	Zn 1	0
11	F	2	Total 2	Zn 2	0














• Molecule 3: Histone deacetylase RPD3














• Molecule 7: Histone H2A type 1-B/E




• Molecule 7: Histone H2A type 1-B/E



• Molecule 8: Histone H2B type 1-K




• Molecule 8: Histone H2B type 1-K



• Molecule 9: 5-DNA



• Molecule 10: 3-DNA



## 4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	213127	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	50	Depositor
Minimum defocus (nm)	1800	Depositor
Maximum defocus (nm)	2300	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	0.056	Depositor
Minimum map value	-0.022	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.003	Depositor
Recommended contour level	0.012	Depositor
Map size ( $\text{\AA}$ )	304.36002, 304.36002, 304.36002	wwPDB
Map dimensions	280, 280, 280	wwPDB
Map angles ( $^\circ$ )	90.0, 90.0, 90.0	wwPDB
Pixel spacing ( $\text{\AA}$ )	1.087, 1.087, 1.087	Depositor

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section:  
ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.26	0/5180	0.41	0/6984
2	E	0.55	2/2952 (0.1%)	0.86	12/3978 (0.3%)
2	F	2.28	3/1313 (0.2%)	0.70	5/1763 (0.3%)
3	B	0.43	2/3137 (0.1%)	0.63	4/4246 (0.1%)
4	C	0.27	0/1509	0.48	0/2039
4	D	1.84	6/1524 (0.4%)	0.49	1/2061 (0.0%)
4	G	0.70	0/583	0.97	3/777 (0.4%)
5	a	0.52	0/813	0.63	2/1090 (0.2%)
5	e	0.51	0/828	0.58	1/1109 (0.1%)
6	b	0.50	0/634	0.50	0/848
6	f	0.46	0/660	0.46	0/883
7	c	0.67	2/827 (0.2%)	0.55	1/1116 (0.1%)
7	g	0.40	0/827	0.49	0/1116
8	d	0.46	0/746	0.55	1/1003 (0.1%)
8	h	0.43	0/746	0.50	0/1003
9	i	0.77	0/3378	0.98	0/5212
10	j	0.77	0/3376	0.99	1/5209 (0.0%)
All	All	0.83	15/29033 (0.1%)	0.72	31/40437 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
2	E	0	1
3	B	0	1
All	All	0	3

The worst 5 of 15 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	372	TYR	CB-CG	80.40	2.72	1.51
4	D	236	TRP	CE3-CZ3	52.29	2.27	1.38
4	D	236	TRP	CZ3-CH2	28.56	1.85	1.40
4	D	236	TRP	CE2-CZ2	25.32	1.82	1.39
4	D	236	TRP	CD2-CE2	21.01	1.66	1.41

The worst 5 of 31 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	384	ARG	CB-CA-C	15.82	142.05	110.40
2	F	372	TYR	CA-CB-CG	12.40	136.95	113.40
2	F	372	TYR	CB-CG-CD2	11.70	128.02	121.00
2	F	372	TYR	CD1-CG-CD2	-11.38	105.38	117.90
2	F	372	TYR	CB-CG-CD1	9.19	126.51	121.00

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	1232	TYR	Peptide
3	B	157	ALA	Mainchain
2	E	470	SER	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5066	0	5024	202	0
2	E	2884	0	2831	196	0
2	F	1282	0	1239	65	0
3	B	3057	0	2930	155	0
4	C	1483	0	1510	31	0
4	D	1497	0	1524	74	0
4	G	570	0	563	0	0
5	a	801	0	839	0	0
5	e	816	0	856	0	0
6	b	627	0	663	0	0
6	f	653	0	696	0	0

*Continued on next page...*



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	c	817	0	872	0	0
7	g	817	0	872	0	0
8	d	735	0	756	0	0
8	h	735	0	756	0	0
9	i	3011	0	1662	0	0
10	j	3010	0	1663	0	0
11	B	1	0	0	0	0
11	E	4	0	0	0	0
11	F	2	0	0	0	0
All	All	27868	0	25256	563	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

The worst 5 of 563 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:D:236:TRP:CE2	4:D:236:TRP:CZ2	1.82	1.65
4:D:236:TRP:CH2	4:D:236:TRP:CZ3	1.85	1.59
1:A:922:LYS:CE	2:E:129:TRP:CZ2	1.96	1.48
1:A:922:LYS:HE3	2:E:129:TRP:CE2	1.49	1.44
2:E:379:ILE:HD12	2:E:380:PRO:CD	1.53	1.39

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	598/1536 (39%)	573 (96%)	24 (4%)	1 (0%)	47 81
2	E	343/684 (50%)	293 (85%)	41 (12%)	9 (3%)	5 35

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	F	152/684 (22%)	150 (99%)	2 (1%)	0	100	100
3	B	383/433 (88%)	362 (94%)	18 (5%)	3 (1%)	19	60
4	C	181/401 (45%)	171 (94%)	9 (5%)	1 (1%)	25	65
4	D	183/401 (46%)	179 (98%)	4 (2%)	0	100	100
4	G	63/401 (16%)	62 (98%)	1 (2%)	0	100	100
5	a	95/136 (70%)	95 (100%)	0	0	100	100
5	e	97/136 (71%)	96 (99%)	1 (1%)	0	100	100
6	b	77/103 (75%)	72 (94%)	5 (6%)	0	100	100
6	f	80/103 (78%)	77 (96%)	3 (4%)	0	100	100
7	c	105/130 (81%)	101 (96%)	4 (4%)	0	100	100
7	g	105/130 (81%)	103 (98%)	2 (2%)	0	100	100
8	d	92/126 (73%)	89 (97%)	3 (3%)	0	100	100
8	h	92/126 (73%)	88 (96%)	4 (4%)	0	100	100
All	All	2646/5530 (48%)	2511 (95%)	121 (5%)	14 (0%)	32	68

5 of 14 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	E	171	LEU
2	E	174	ASN
2	E	470	SER
3	B	114	ASP
3	B	115	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	560/1391 (40%)	532 (95%)	28 (5%)	24	51
2	E	332/653 (51%)	293 (88%)	39 (12%)	5	23
2	F	148/653 (23%)	143 (97%)	5 (3%)	37	61

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	B	326/367 (89%)	299 (92%)	27 (8%)	11	37
4	C	172/359 (48%)	159 (92%)	13 (8%)	13	40
4	D	174/359 (48%)	170 (98%)	4 (2%)	50	70
4	G	56/359 (16%)	56 (100%)	0	100	100
5	a	85/111 (77%)	77 (91%)	8 (9%)	8	30
5	e	86/111 (78%)	79 (92%)	7 (8%)	11	37
6	b	64/79 (81%)	61 (95%)	3 (5%)	26	53
6	f	67/79 (85%)	64 (96%)	3 (4%)	27	54
7	c	83/100 (83%)	80 (96%)	3 (4%)	35	60
7	g	83/100 (83%)	78 (94%)	5 (6%)	19	46
8	d	80/105 (76%)	74 (92%)	6 (8%)	13	40
8	h	80/105 (76%)	76 (95%)	4 (5%)	24	51
All	All	2396/4931 (49%)	2241 (94%)	155 (6%)	21	44

5 of 155 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	F	570	GLN
6	f	30	THR
5	a	87	SER
8	d	31	ARG
7	g	76	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 41 such sidechains are listed below:

Mol	Chain	Res	Type
4	D	317	GLN
6	b	93	GLN
4	D	375	ASN
2	F	355	ASN
5	e	125	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 7 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-37367. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

### 6.1 Orthogonal projections [i](#)


#### 6.1.1 Primary map



X




Y




Z


#### 6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

## 6.2 Central slices [i](#)


### 6.2.1 Primary map



X Index: 140




Y Index: 140




Z Index: 140


### 6.2.2 Raw map



X Index: 140



Y Index: 140




Z Index: 140


The images above show central slices of the map in three orthogonal directions.

## 6.3 Largest variance slices [i](#)


### 6.3.1 Primary map



X Index: 154




Y Index: 168




Z Index: 121


### 6.3.2 Raw map



X Index: 154



Y Index: 134




Z Index: 121


The images above show the largest variance slices of the map in three orthogonal directions.

## 6.4 Orthogonal standard-deviation projections (False-color) [i](#)


### 6.4.1 Primary map



X




Y




Z


### 6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.




## 6.5 Orthogonal surface views [i](#)

### 6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.012. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

### 6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


## 6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

## 7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


## 7.2 Volume estimate [\(i\)](#)



The volume at the recommended contour level is 279 nm<sup>3</sup>; this corresponds to an approximate mass of 252 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

### 7.3 Rotationally averaged power spectrum [i](#)




\*Reported resolution corresponds to spatial frequency of 0.227 Å<sup>-1</sup>

## 8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

### 8.1 FSC [i](#)



\*Reported resolution corresponds to spatial frequency of 0.227 Å<sup>-1</sup>

## 8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	4.40	-	-
Author-provided FSC curve	4.35	7.03	4.59
Unmasked-calculated*	6.41	9.30	6.86

\*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 6.41 differs from the reported value 4.4 by more than 10 %

## 9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-37367 and PDB model 8W9F. Per-residue inclusion information can be found in section 3 on page 7.

### 9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.012 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

## 9.2 Q-score mapped to coordinate model [\(i\)](#)



The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


## 9.3 Atom inclusion mapped to coordinate model [\(i\)](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.012).







































## 9.4 Atom inclusion [i](#)



At the recommended contour level, 61% of all backbone atoms, 57% of all non-hydrogen atoms, are inside the map.

## 9.5 Map-model fit summary [i](#)

The table lists the average atom inclusion at the recommended contour level (0.012) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.5710	 0.1560
A	 0.3860	 0.0520
B	 0.6130	 0.0480
C	 0.4590	 0.0390
D	 0.1660	 0.0200
E	 0.2740	 0.0410
F	 0.0950	 0.0210
G	 0.0290	 0.0430
a	 0.7360	 0.3290
b	 0.7980	 0.3710
c	 0.8160	 0.3750
d	 0.8030	 0.3820
e	 0.8340	 0.3790
f	 0.8600	 0.4020
g	 0.7880	 0.3750
h	 0.8230	 0.3800
i	 0.8890	 0.2500
j	 0.9020	 0.2260

