



# Full wwPDB X-ray Structure Validation Report ⓘ

Dec 9, 2023 – 10:10 am GMT

PDB ID : 2WBV  
Title : Canine adenovirus 2 fibre head in complex with sialic acid  
Authors : Seiradake, E.; Henaff, D.; Wodrich, H.; Billet, O.; Perreau, M.; Hippert, C.; Mennechet, F.; Schoehn, G.; Lortat-Jacob, H.; Dreja, H.; Ibanes, S.; Kalatzis, V.; Wang, J.P.; Finberg, R.W.; Cusack, S.; Kremer, E.J.  
Deposited on : 2009-03-05  
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	189	 92% 5% .
1	B	189	 2% 92% 7% .
1	C	189	 93% . .
1	D	189	 % 91% 6% .

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	E	189	 94% . .
1	F	189	 % 93% 5% .
2	G	2	 50% 50%

## 2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 9900 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called FIBER PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	183	1437	913	246	269	9	0	4	0
1	B	186	1435	910	243	273	9	0	1	0
1	C	183	1424	904	242	269	9	0	3	0
1	D	184	1431	912	241	269	9	0	3	0
1	E	182	1418	900	241	268	9	0	3	0
1	F	184	1430	909	244	268	9	0	2	0

- Molecule 2 is an oligosaccharide called N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	G	2	32	17	1	14	0	0	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	C	1	Total C O 6 3 3	0	0
3	D	1	Total C O 6 3 3	0	0
3	E	1	Total C O 6 3 3	0	0
3	F	1	Total C O 6 3 3	0	0

- Molecule 4 is N-acetyl-alpha-neuraminic acid (three-letter code: SIA) (formula: C<sub>11</sub>H<sub>19</sub>NO<sub>9</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			21	11	1	9		
4	B	1	Total	C	N	O	0	0
			21	11	1	9		
4	C	1	Total	C	N	O	0	0
			21	11	1	9		
4	E	1	Total	C	N	O	0	0
			21	11	1	9		
4	F	1	Total	C	N	O	0	0
			21	11	1	9		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	208	Total	O	0	0
			208	208		
5	B	180	Total	O	0	0
			180	180		
5	C	217	Total	O	0	0
			217	217		
5	D	193	Total	O	0	0
			193	193		
5	E	184	Total	O	0	0
			184	184		
5	F	170	Total	O	0	0
			170	170		

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: FIBER PROTEIN

Chain A:  92% 5%




- Molecule 1: FIBER PROTEIN

Chain B:  2% 92% 7%




- Molecule 1: FIBER PROTEIN

Chain C:  93%




- Molecule 1: FIBER PROTEIN

Chain D:  % 91% 6%



- Molecule 1: FIBER PROTEIN

Chain E:  94%



- Molecule 1: FIBER PROTEIN

Chain F:  % 93% 5%



- Molecule 2: N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose

Chain G: 50% 50%





## 4 Data and refinement statistics i

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	215.00Å 61.25Å 143.11Å 90.00° 131.67° 90.00°	Depositor
Resolution (Å)	46.13 – 1.90 46.11 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.8 (46.13-1.90) 99.8 (46.11-1.90)	Depositor EDS
$R_{merge}$	0.15	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.95 (at 1.90Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.157 , 0.195 0.164 , 0.200	Depositor DCC
$R_{free}$ test set	5480 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	11.5	Xtrriage
Anisotropy	0.214	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 45.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.022 for -h-2*1,-k,l	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	9900	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	12.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.40% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, GAL, SIA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.44	0/1488	0.68	2/2026 (0.1%)
1	B	0.43	0/1480	0.62	0/2019
1	C	0.43	0/1469	0.62	0/2003
1	D	0.42	0/1476	0.62	0/2012
1	E	0.43	0/1462	0.63	0/1991
1	F	0.40	0/1473	0.60	0/2007
All	All	0.43	0/8848	0.63	2/12058 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	493[A]	ARG	NE-CZ-NH2	-6.00	117.30	120.30
1	A	493[B]	ARG	NE-CZ-NH2	-6.00	117.30	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1437	0	1430	3	0
1	B	1435	0	1410	7	0
1	C	1424	0	1404	3	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	1431	0	1422	7	0
1	E	1418	0	1401	1	0
1	F	1430	0	1415	5	0
2	G	32	0	28	0	0
3	A	12	0	16	0	0
3	C	6	0	8	2	0
3	D	6	0	8	1	0
3	E	6	0	8	0	0
3	F	6	0	8	0	0
4	A	21	0	18	1	0
4	B	21	0	18	1	0
4	C	21	0	18	0	0
4	E	21	0	18	0	0
4	F	21	0	18	0	0
5	A	208	0	0	0	0
5	B	180	0	0	1	1
5	C	217	0	0	0	0
5	D	193	0	0	0	0
5	E	184	0	0	0	0
5	F	170	0	0	0	0
All	All	9900	0	8648	25	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:361:ALA:HB3	5:B:2003:HOH:O	1.89	0.72
1:A:387:ILE:HD11	1:A:436:ILE:HD13	1.72	0.71
1:A:416[B]:SER:OG	4:A:1545:SIA:H113	1.98	0.63
1:D:387[A]:ILE:HD11	1:D:436:ILE:HD13	1.81	0.60
1:F:417:GLN:HB2	1:F:517:LEU:HD23	1.85	0.58
1:C:370:GLY:H	3:C:1544:GOL:H31	1.69	0.57
1:B:416[B]:SER:OG	4:B:1543:SIA:H113	2.05	0.57
1:D:370:GLY:H	3:D:1543:GOL:C3	2.21	0.54
1:F:387[A]:ILE:HD11	1:F:436:ILE:HD13	1.90	0.53
1:B:358:PRO:N	1:B:359:PRO:HD2	2.26	0.49
1:F:467:ARG:HG3	1:F:510:TYR:CE1	2.48	0.48
1:D:387[A]:ILE:CD1	1:D:436:ILE:HD13	2.45	0.47
1:A:387:ILE:CD1	1:A:436:ILE:HD13	2.43	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:377:ILE:HD12	1:B:382:VAL:HG21	1.98	0.45
1:E:477:ILE:HG22	1:F:531:ASP:OD2	2.16	0.45
1:B:467:ARG:HG3	1:B:510:TYR:CE1	2.52	0.45
1:D:410[A]:ILE:HD13	1:D:527:LEU:HA	1.99	0.45
1:B:422:MET:HG2	1:B:436:ILE:HB	1.98	0.44
1:C:398:VAL:O	1:C:534:THR:HA	2.18	0.44
1:F:387[A]:ILE:CD1	1:F:436:ILE:HD13	2.47	0.43
1:D:483:ASP:O	1:D:487:VAL:HG22	2.19	0.43
1:C:370:GLY:H	3:C:1544:GOL:C3	2.32	0.42
1:D:398:VAL:O	1:D:534:THR:HA	2.19	0.41
1:D:467:ARG:HD2	1:D:509:THR:HG22	2.01	0.41
1:B:361:ALA:N	1:B:362:PRO:CD	2.85	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:B:2001:HOH:O	5:B:2129:HOH:O[4_454]	1.93	0.27

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	186/189 (98%)	176 (95%)	10 (5%)	0	100	100
1	B	186/189 (98%)	178 (96%)	8 (4%)	0	100	100
1	C	184/189 (97%)	176 (96%)	8 (4%)	0	100	100
1	D	185/189 (98%)	176 (95%)	9 (5%)	0	100	100
1	E	183/189 (97%)	175 (96%)	8 (4%)	0	100	100
1	F	184/189 (97%)	177 (96%)	7 (4%)	0	100	100
All	All	1108/1134 (98%)	1058 (96%)	50 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	164/163 (101%)	157 (96%)	7 (4%)	29	19
1	B	164/163 (101%)	162 (99%)	2 (1%)	71	70
1	C	162/163 (99%)	157 (97%)	5 (3%)	40	32
1	D	163/163 (100%)	162 (99%)	1 (1%)	86	87
1	E	162/163 (99%)	159 (98%)	3 (2%)	57	53
1	F	162/163 (99%)	160 (99%)	2 (1%)	71	70
All	All	977/978 (100%)	957 (98%)	20 (2%)	57	51

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	363	ILE
1	A	366	TRP
1	A	369	PRO
1	A	412	SER
1	A	467[A]	ARG
1	A	467[B]	ARG
1	A	521	ARG
1	B	366	TRP
1	B	379	ASP
1	C	363	ILE
1	C	366	TRP
1	C	412[A]	SER
1	C	412[B]	SER
1	C	441	THR
1	D	366	TRP
1	E	366	TRP
1	E	457	SER
1	E	527	LEU
1	F	366	TRP

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type
1	F	453	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	F	426	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	GAL	G	1	2	12,12,12	0.44	0	17,17,17	0.60	0
2	SIA	G	2	2	20,20,21	0.57	0	24,28,31	1.06	2 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GAL	G	1	2	-	0/2/22/22	0/1/1/1
2	SIA	G	2	2	-	0/18/34/38	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	G	2	SIA	O1B-C1-C2	2.04	118.87	113.03
2	G	2	SIA	C4-C5-N5	-2.02	106.38	110.38


There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



## 5.6 Ligand geometry [i](#)

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	GOL	C	1544	-	5,5,5	0.39	0	5,5,5	0.75	0
3	GOL	D	1543	-	5,5,5	0.33	0	5,5,5	0.65	0
3	GOL	E	1543	-	5,5,5	0.38	0	5,5,5	0.39	0
3	GOL	A	1543	-	5,5,5	0.34	0	5,5,5	0.29	0
4	SIA	E	1544	-	21,21,21	0.99	1 (4%)	25,31,31	0.94	1 (4%)
4	SIA	B	1543	-	21,21,21	0.82	1 (4%)	25,31,31	1.06	2 (8%)
3	GOL	F	1543	-	5,5,5	0.40	0	5,5,5	0.43	0
4	SIA	C	1543	-	21,21,21	1.02	2 (9%)	25,31,31	1.16	3 (12%)
3	GOL	A	1544	-	5,5,5	0.38	0	5,5,5	0.27	0
4	SIA	A	1545	-	21,21,21	0.90	1 (4%)	25,31,31	1.15	2 (8%)
4	SIA	F	1544	-	21,21,21	0.86	1 (4%)	25,31,31	1.11	3 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	C	1544	-	-	2/4/4/4	-
3	GOL	D	1543	-	-	4/4/4/4	-
3	GOL	E	1543	-	-	2/4/4/4	-
3	GOL	A	1543	-	-	4/4/4/4	-
4	SIA	E	1544	-	-	2/20/38/38	0/1/1/1
4	SIA	B	1543	-	-	2/20/38/38	0/1/1/1
3	GOL	F	1543	-	-	3/4/4/4	-
4	SIA	C	1543	-	-	1/20/38/38	0/1/1/1
3	GOL	A	1544	-	-	3/4/4/4	-
4	SIA	A	1545	-	-	2/20/38/38	0/1/1/1
4	SIA	F	1544	-	-	0/20/38/38	0/1/1/1

All (6) bond length outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	1543	SIA	O2-C2	3.32	1.43	1.39
4	E	1544	SIA	O2-C2	3.17	1.43	1.39
4	A	1545	SIA	O2-C2	2.70	1.43	1.39
4	B	1543	SIA	O2-C2	2.55	1.42	1.39
4	F	1544	SIA	O2-C2	2.55	1.42	1.39
4	C	1543	SIA	C3-C2	2.27	1.54	1.51

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1545	SIA	O1A-C1-C2	-3.91	117.67	123.59
4	F	1544	SIA	O1A-C1-C2	-3.89	117.69	123.59
4	E	1544	SIA	O1A-C1-C2	-3.35	118.51	123.59
4	C	1543	SIA	O1A-C1-C2	-3.30	118.58	123.59
4	B	1543	SIA	O1A-C1-C2	-3.21	118.72	123.59
4	C	1543	SIA	C6-C5-N5	-2.89	106.11	110.91
4	A	1545	SIA	O2-C2-C1	-2.72	105.22	110.76
4	F	1544	SIA	O2-C2-C1	-2.66	105.34	110.76
4	C	1543	SIA	O6-C6-C5	-2.12	107.70	109.78
4	B	1543	SIA	C4-C5-N5	-2.01	106.40	110.38
4	F	1544	SIA	C4-C5-N5	-2.00	106.42	110.38

There are no chirality outliers.

All (25) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1543	GOL	C1-C2-C3-O3
3	C	1544	GOL	O1-C1-C2-C3
3	D	1543	GOL	O1-C1-C2-C3
3	D	1543	GOL	C1-C2-C3-O3
3	E	1543	GOL	O1-C1-C2-C3
3	F	1543	GOL	O1-C1-C2-C3
3	A	1544	GOL	O1-C1-C2-O2
3	E	1543	GOL	O1-C1-C2-O2
3	A	1543	GOL	O1-C1-C2-C3
3	A	1544	GOL	O1-C1-C2-C3
3	F	1543	GOL	C1-C2-C3-O3
3	D	1543	GOL	O2-C2-C3-O3
3	F	1543	GOL	O1-C1-C2-O2
3	C	1544	GOL	O1-C1-C2-O2
3	D	1543	GOL	O1-C1-C2-O2
3	A	1543	GOL	O2-C2-C3-O3

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Res	Type	Atoms
4	A	1545	SIA	O1B-C1-C2-C3
4	B	1543	SIA	O1B-C1-C2-C3
4	E	1544	SIA	O1B-C1-C2-C3
3	A	1543	GOL	O1-C1-C2-O2
4	A	1545	SIA	O1A-C1-C2-O2
4	B	1543	SIA	O1A-C1-C2-O2
4	C	1543	SIA	O1A-C1-C2-O2
4	E	1544	SIA	O1A-C1-C2-O2
3	A	1544	GOL	C1-C2-C3-O3


There are no ring outliers.


4 monomers are involved in 5 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	1544	GOL	2	0
3	D	1543	GOL	1	0
4	B	1543	SIA	1	0
4	A	1545	SIA	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	183/189 (96%)	-0.63	0 <b>100</b> <b>100</b>	5, 9, 17, 24	0
1	B	186/189 (98%)	-0.61	3 (1%) 72 74	4, 9, 17, 33	0
1	C	183/189 (96%)	-0.69	0 <b>100</b> <b>100</b>	5, 10, 15, 23	0
1	D	184/189 (97%)	-0.55	1 (0%) 91 92	4, 10, 17, 24	0
1	E	182/189 (96%)	-0.64	0 <b>100</b> <b>100</b>	6, 10, 17, 22	0
1	F	184/189 (97%)	-0.51	2 (1%) 80 82	6, 13, 20, 29	1 (0%)
All	All	1102/1134 (97%)	-0.60	6 (0%) 91 92	4, 10, 18, 33	1 (0%)

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	359	PRO	3.8
1	B	455	ARG	3.1
1	D	455	ARG	3.0
1	F	521[A]	ARG	2.6
1	B	357	SER	2.3
1	B	360	ALA	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.


### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	GAL	G	1	12/12	0.91	0.17	23,29,31,33	0
2	SIA	G	2	20/21	0.96	0.09	9,14,19,20	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	GOL	D	1543	6/6	0.74	0.24	39,40,41,42	0
3	GOL	A	1544	6/6	0.77	0.20	35,37,38,38	0
3	GOL	A	1543	6/6	0.77	0.20	53,53,53,53	0
3	GOL	E	1543	6/6	0.80	0.22	48,49,49,50	0
3	GOL	F	1543	6/6	0.81	0.15	42,42,43,44	0
3	GOL	C	1544	6/6	0.87	0.23	38,39,39,40	0

*Continued on next page...*




*Continued from previous page...*


<b>Mol</b>	<b>Type</b>	<b>Chain</b>	<b>Res</b>	<b>Atoms</b>	<b>RSCC</b>	<b>RSR</b>	<b>B-factors(<math>\text{\AA}^2</math>)</b>	<b>Q&lt;0.9</b>
4	SIA	F	1544	21/21	0.93	0.11	16,19,22,22	0
4	SIA	E	1544	21/21	0.94	0.07	11,13,18,18	0
4	SIA	C	1543	21/21	0.94	0.08	10,13,16,18	0
4	SIA	B	1543	21/21	0.95	0.08	11,14,17,17	0
4	SIA	A	1545	21/21	0.96	0.07	9,11,13,14	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around SIA F 1544:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around SIA C 1543:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around SIA B 1543:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers ⓘ

There are no such residues in this entry.