



# Full wwPDB X-ray Structure Validation Report i

Jan 22, 2024 – 08:16 PM JST

PDB ID : 8WBS  
Title : Crystal structure of cis-Epoxysuccinate Hydrolases KlCESH[L]-D48N complexed with sulfate ions  
Authors : Dong, S.; Xuan, J.S.; Feng, Y.G.; Cui, Q.  
Deposited on : 2023-09-10  
Resolution : 2.03 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>  
with specific help available everywhere you see the i symbol.

The types of validation reports are described at  
<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## X-RAY DIFFRACTION


The reported resolution of this entry is 2.03 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R <sub>free</sub>	130704	10434 (2.04-2.00)
Clashscore	141614	11643 (2.04-2.00)
Ramachandran outliers	138981	11493 (2.04-2.00)
Sidechain outliers	138945	11492 (2.04-2.00)
RSRZ outliers	127900	10220 (2.04-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SO4	D	301	-	-	X	-

## 2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 7748 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called (S)-2-haloacid dehalogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	236	Total	C	N	O	S	0	0	0
			1865	1193	321	345	6			
1	B	236	Total	C	N	O	S	0	0	0
			1865	1193	321	345	6			
1	C	237	Total	C	N	O	S	0	0	0
			1866	1194	321	346	5			
1	D	221	Total	C	N	O	S	0	0	0
			1753	1124	296	328	5			

There are 40 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	40	MET	-	initiating methionine	UNP W8PFD2
A	48	ASN	ASP	engineered mutation	UNP W8PFD2
A	275	LEU	-	expression tag	UNP W8PFD2
A	276	GLU	-	expression tag	UNP W8PFD2
A	277	HIS	-	expression tag	UNP W8PFD2
A	278	HIS	-	expression tag	UNP W8PFD2
A	279	HIS	-	expression tag	UNP W8PFD2
A	280	HIS	-	expression tag	UNP W8PFD2
A	281	HIS	-	expression tag	UNP W8PFD2
A	282	HIS	-	expression tag	UNP W8PFD2
B	40	MET	-	initiating methionine	UNP W8PFD2
B	48	ASN	ASP	engineered mutation	UNP W8PFD2
B	275	LEU	-	expression tag	UNP W8PFD2
B	276	GLU	-	expression tag	UNP W8PFD2
B	277	HIS	-	expression tag	UNP W8PFD2
B	278	HIS	-	expression tag	UNP W8PFD2
B	279	HIS	-	expression tag	UNP W8PFD2
B	280	HIS	-	expression tag	UNP W8PFD2
B	281	HIS	-	expression tag	UNP W8PFD2
B	282	HIS	-	expression tag	UNP W8PFD2
C	40	MET	-	initiating methionine	UNP W8PFD2

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
C	48	ASN	ASP	engineered mutation	UNP W8PFD2
C	275	LEU	-	expression tag	UNP W8PFD2
C	276	GLU	-	expression tag	UNP W8PFD2
C	277	HIS	-	expression tag	UNP W8PFD2
C	278	HIS	-	expression tag	UNP W8PFD2
C	279	HIS	-	expression tag	UNP W8PFD2
C	280	HIS	-	expression tag	UNP W8PFD2
C	281	HIS	-	expression tag	UNP W8PFD2
C	282	HIS	-	expression tag	UNP W8PFD2
D	40	MET	-	initiating methionine	UNP W8PFD2
D	48	ASN	ASP	engineered mutation	UNP W8PFD2
D	275	LEU	-	expression tag	UNP W8PFD2
D	276	GLU	-	expression tag	UNP W8PFD2
D	277	HIS	-	expression tag	UNP W8PFD2
D	278	HIS	-	expression tag	UNP W8PFD2
D	279	HIS	-	expression tag	UNP W8PFD2
D	280	HIS	-	expression tag	UNP W8PFD2
D	281	HIS	-	expression tag	UNP W8PFD2
D	282	HIS	-	expression tag	UNP W8PFD2

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	C	1	Total O S 5 4 1	0	0
2	C	1	Total O S 5 4 1	0	0
2	D	1	Total O S 5 4 1	0	0
2	D	1	Total O S 5 4 1	0	0

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Ca 1 1	0	0
3	C	1	Total Ca 1 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	141	Total O 141 141	0	0
4	B	129	Total O 129 129	0	0
4	C	68	Total O 68 68	0	0
4	D	19	Total O 19 19	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: (S)-2-haloacid dehalogenase



- Molecule 1: (S)-2-haloacid dehalogenase




- Molecule 1: (S)-2-haloacid dehalogenase



- Molecule 1: (S)-2-haloacid dehalogenase





## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	119.85Å 94.43Å 99.60Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	76.60 – 2.03 76.60 – 2.03	Depositor EDS
% Data completeness (in resolution range)	99.9 (76.60-2.03) 99.9 (76.60-2.03)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) >$ <sup>1</sup>	2.14 (at 2.03Å)	Xtriage
Refinement program	PHENIX v1.21	Depositor
$R$ , $R_{free}$	0.178 , 0.209 0.179 , 0.210	Depositor DCC
$R_{free}$ test set	3813 reflections (5.18%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	36.6	Xtriage
Anisotropy	0.237	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 61.7	EDS
L-test for twinning <sup>2</sup>	$<  L  > = 0.50$ , $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	7748	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.97% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $< |L| >$ ,  $< L^2 >$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [\(i\)](#)

### 5.1 Standard geometry [\(i\)](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.43	0/1909	0.60	0/2589
1	B	0.42	0/1909	0.58	0/2589
1	C	0.40	0/1911	0.56	0/2594
1	D	0.31	0/1790	0.52	0/2422
All	All	0.39	0/7519	0.56	0/10194

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1865	0	1851	8	0
1	B	1865	0	1851	7	0
1	C	1866	0	1840	21	0
1	D	1753	0	1740	36	0
2	A	10	0	0	0	0
2	B	10	0	0	0	0
2	C	10	0	0	1	0
2	D	10	0	0	6	0
3	A	1	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	C	1	0	0	0	0
4	A	141	0	0	0	1
4	B	129	0	0	0	0
4	C	68	0	0	1	1
4	D	19	0	0	1	0
All	All	7748	0	7282	69	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (69) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:87:GLU:HA	1:D:90:SER:HB3	1.49	0.94
1:D:153:ARG:NH2	4:D:401:HOH:O	2.09	0.84
1:D:48:ASN:ND2	2:D:301:SO4:O3	2.25	0.69
1:D:58:SER:O	1:D:62:ARG:HG2	1.93	0.67
1:D:48:ASN:ND2	2:D:301:SO4:O4	2.27	0.67
1:D:93:ASP:HB3	1:D:97:LYS:HZ1	1.59	0.67
1:C:60:ILE:HD12	1:C:81:THR:HG23	1.76	0.66
1:C:249:LYS:HD2	1:C:249:LYS:H	1.63	0.64
1:B:40:MET:HG3	1:B:213:HIS:HE1	1.62	0.63
1:A:40:MET:HA	1:A:214:GLN:HE21	1.63	0.63
1:C:241:PRO:HA	1:C:250:VAL:HG11	1.80	0.63
1:C:250:VAL:HG21	4:C:401:HOH:O	2.00	0.60
1:D:48:ASN:ND2	2:D:301:SO4:S	2.76	0.59
1:C:79:THR:O	1:C:83:GLN:HG3	2.03	0.57
1:D:50:GLN:NE2	2:D:301:SO4:O1	2.37	0.57
1:D:87:GLU:OE2	1:D:112:ALA:HB1	2.05	0.57
1:C:164:SER:OG	2:C:301:SO4:O3	2.21	0.56
1:D:50:GLN:HA	1:D:54:VAL:HG12	1.88	0.56
1:C:153:ARG:CZ	1:C:276:GLU:HG3	2.37	0.55
1:D:65:GLU:HG3	1:D:77:TRP:CD1	2.42	0.55
1:B:40:MET:HG3	1:B:213:HIS:CE1	2.42	0.54
1:D:250:VAL:HG23	1:D:252:THR:HG23	1.90	0.54
1:D:94:GLN:O	1:D:99:GLU:N	2.40	0.54
1:A:107:ARG:NH2	1:C:249:LYS:HE2	2.24	0.52
1:C:226:GLN:O	1:C:230:ARG:HG2	2.09	0.52
1:D:50:GLN:NE2	2:D:301:SO4:S	2.83	0.52
1:D:56:PHE:CZ	1:D:85:ARG:NH1	2.78	0.52
1:C:96:ILE:HD11	1:C:194:SER:HB3	1.92	0.51

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:221:HIS:NE2	1:C:243:GLU:OE2	2.40	0.50
1:A:105:THR:HG21	1:A:166:GLY:HA2	1.93	0.49
1:C:260:TYR:HB3	1:C:271:MET:HE1	1.94	0.49
1:D:93:ASP:HA	1:D:96:ILE:HB	1.96	0.48
1:C:249:LYS:H	1:C:249:LYS:CD	2.27	0.47
1:C:104:THR:OG1	1:C:107:ARG:HG2	2.15	0.47
1:D:201:TYR:O	1:D:205:LEU:HD23	2.15	0.47
1:D:153:ARG:NE	1:D:153:ARG:HA	2.31	0.46
1:A:221:HIS:NE2	1:A:243:GLU:OE2	2.41	0.46
1:D:67:PHE:CE2	1:D:134:LEU:HD12	2.51	0.45
1:D:91:ARG:HH21	1:D:115:GLY:HA3	1.82	0.45
1:D:76:ASP:O	1:D:80:VAL:HG23	2.17	0.45
1:D:97:LYS:HE2	1:D:97:LYS:HB2	1.77	0.45
1:D:105:THR:HG21	1:D:166:GLY:HA2	2.00	0.44
1:C:260:TYR:CD2	1:C:271:MET:HE3	2.53	0.44
1:D:46:PHE:CE1	1:D:204:ALA:HB1	2.53	0.43
1:C:268:LEU:HA	1:C:271:MET:HE2	2.00	0.43
1:B:87:GLU:OE2	1:B:91:ARG:NE	2.49	0.43
1:C:117:LEU:HD23	1:C:117:LEU:HA	1.78	0.43
1:C:268:LEU:HD12	1:C:271:MET:HE2	2.00	0.43
1:D:85:ARG:NH2	2:D:302:SO4:O2	2.52	0.42
1:D:268:LEU:HA	1:D:271:MET:CE	2.49	0.42
1:D:65:GLU:HG3	1:D:77:TRP:HD1	1.82	0.42
1:D:97:LYS:HE2	1:D:99:GLU:HG3	2.00	0.42
1:A:78:THR:O	1:A:82:GLU:HG3	2.19	0.42
1:B:248:LYS:HA	1:B:248:LYS:HD3	1.87	0.42
1:A:159:ILE:HG23	1:B:172:LEU:HD21	2.02	0.42
1:A:223:TYR:CG	1:A:224:ASP:N	2.87	0.42
1:D:87:GLU:CA	1:D:90:SER:HB3	2.34	0.41
1:A:195:LYS:HG2	1:A:201:TYR:CZ	2.56	0.41
1:D:223:TYR:CG	1:D:224:ASP:N	2.88	0.41
1:C:271:MET:HE2	1:C:271:MET:HB2	1.76	0.41
1:D:147:THR:HG22	1:D:151:LEU:HD22	2.03	0.41
1:D:88:TYR:CE1	1:D:105:THR:HG23	2.55	0.41
1:C:93:ASP:O	1:C:97:LYS:HG2	2.20	0.41
1:C:184:ALA:HB2	1:D:172:LEU:HD12	2.03	0.41
1:D:67:PHE:CD2	1:D:134:LEU:HD12	2.55	0.41
1:B:60:ILE:HD11	1:B:138:TRP:CZ2	2.56	0.41
1:D:262:ALA:HB2	1:D:271:MET:CE	2.51	0.41
1:B:62:ARG:HH11	1:B:62:ARG:HG3	1.87	0.40
1:D:230:ARG:HB3	1:D:230:ARG:NH1	2.36	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:470:HOH:O	4:C:461:HOH:O[3_555]	2.12	0.08

## 5.3 Torsion angles [\(i\)](#)

### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	234/243 (96%)	232 (99%)	2 (1%)	0	100 100
1	B	234/243 (96%)	231 (99%)	3 (1%)	0	100 100
1	C	235/243 (97%)	232 (99%)	3 (1%)	0	100 100
1	D	213/243 (88%)	203 (95%)	10 (5%)	0	100 100
All	All	916/972 (94%)	898 (98%)	18 (2%)	0	100 100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	195/202 (96%)	192 (98%)	3 (2%)	65 68
1	B	195/202 (96%)	193 (99%)	2 (1%)	76 80
1	C	194/202 (96%)	187 (96%)	7 (4%)	35 32
1	D	185/202 (92%)	179 (97%)	6 (3%)	39 37

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	769/808 (95%)	751 (98%)	18 (2%)	50 51

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	151	LEU
1	A	175	SER
1	A	248	LYS
1	B	88	TYR
1	B	151	LEU
1	C	87	GLU
1	C	88	TYR
1	C	90	SER
1	C	116	ILE
1	C	132	ASP
1	C	151	LEU
1	C	249	LYS
1	D	87	GLU
1	D	89	ARG
1	D	90	SER
1	D	91	ARG
1	D	131	ARG
1	D	151	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	213	HIS
1	D	256	GLN

### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [\(i\)](#)

Of 10 ligands modelled in this entry, 2 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	SO4	C	301	-	4,4,4	0.18	0	6,6,6	0.28	0
2	SO4	D	302	-	4,4,4	0.17	0	6,6,6	0.11	0
2	SO4	C	302	-	4,4,4	0.13	0	6,6,6	0.18	0
2	SO4	B	301	-	4,4,4	0.27	0	6,6,6	0.96	0
2	SO4	D	301	-	4,4,4	0.14	0	6,6,6	0.12	0
2	SO4	A	301	-	4,4,4	0.33	0	6,6,6	0.48	0
2	SO4	B	302	-	4,4,4	0.17	0	6,6,6	0.38	0
2	SO4	A	302	-	4,4,4	0.18	0	6,6,6	0.13	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	301	SO4	1	0
2	D	302	SO4	1	0
2	D	301	SO4	5	0

## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	236/243 (97%)	0.61	2 (0%) 86 85	25, 35, 61, 76	0
1	B	236/243 (97%)	0.55	2 (0%) 86 85	25, 37, 59, 73	0
1	C	237/243 (97%)	0.95	38 (16%) 1 1	28, 54, 103, 122	0
1	D	221/243 (90%)	2.07	75 (33%) 0 0	38, 72, 141, 156	0
All	All	930/972 (95%)	1.03	117 (12%) 3 3	25, 46, 116, 156	0

All (117) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	90	SER	12.3
1	D	88	TYR	11.3
1	D	113	LEU	10.6
1	D	81	THR	9.6
1	D	112	ALA	9.6
1	D	86	ALA	9.5
1	D	95	VAL	9.4
1	D	73	PHE	9.2
1	D	79	THR	7.8
1	D	115	GLY	7.6
1	C	96	ILE	7.5
1	D	83	GLN	7.1
1	D	127	ASN	6.9
1	C	129	ALA	6.5
1	D	93	ASP	6.4
1	D	129	ALA	6.4
1	D	75	ALA	6.3
1	D	77	TRP	6.3
1	D	128	SER	6.0
1	D	101	PRO	5.9
1	D	109	TYR	5.9

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	D	87	GLU	5.8
1	D	91	ARG	5.8
1	D	60	ILE	5.6
1	D	85	ARG	5.5
1	D	247	ASN	5.4
1	D	92	LEU	5.3
1	D	74	GLN	5.3
1	C	123	GLY	5.2
1	D	68	SER	5.2
1	C	88	TYR	5.0
1	C	125	SER	5.0
1	D	99	GLU	5.0
1	D	80	VAL	5.0
1	D	84	TRP	4.7
1	D	116	ILE	4.6
1	C	102	TRP	4.5
1	D	126	LEU	4.4
1	D	102	TRP	4.3
1	C	97	LYS	4.3
1	D	76	ASP	4.1
1	D	132	ASP	4.1
1	C	73	PHE	4.1
1	D	66	ALA	4.1
1	D	111	GLU	4.0
1	C	117	LEU	4.0
1	C	118	ALA	3.9
1	C	92	LEU	3.9
1	D	108	ILE	3.8
1	D	67	PHE	3.8
1	D	244	PHE	3.8
1	D	114	ASP	3.8
1	D	96	ILE	3.7
1	D	78	THR	3.7
1	C	86	ALA	3.6
1	C	130	ASP	3.6
1	C	103	THR	3.6
1	D	133	GLU	3.6
1	C	124	ALA	3.5
1	C	74	GLN	3.5
1	D	94	GLN	3.5
1	D	134	LEU	3.4
1	D	138	TRP	3.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	116	ILE	3.3
1	D	103	THR	3.3
1	D	225	LEU	3.2
1	C	249	LYS	3.1
1	C	94	GLN	3.1
1	D	107	ARG	3.1
1	D	223	TYR	3.0
1	D	194	SER	3.0
1	C	122	TRP	3.0
1	A	40	MET	2.9
1	D	97	LYS	2.9
1	D	273	GLY	2.8
1	C	114	ASP	2.8
1	C	90	SER	2.8
1	D	248	LYS	2.7
1	D	63	GLU	2.7
1	C	70	VAL	2.7
1	C	134	LEU	2.7
1	D	130	ASP	2.7
1	C	119	ASN	2.6
1	D	82	GLU	2.6
1	D	131	ARG	2.6
1	D	62	ARG	2.5
1	D	192	ARG	2.5
1	C	77	TRP	2.5
1	D	98	GLY	2.5
1	C	128	SER	2.5
1	C	98	GLY	2.5
1	D	110	ARG	2.5
1	D	222	LYS	2.4
1	D	249	LYS	2.4
1	D	104	THR	2.4
1	C	67	PHE	2.4
1	C	71	ARG	2.4
1	C	110	ARG	2.4
1	D	197	ASP	2.4
1	B	40	MET	2.4
1	C	107	ARG	2.4
1	C	126	LEU	2.3
1	D	44	ALA	2.3
1	D	230	ARG	2.3
1	D	233	PHE	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	90	SER	2.2
1	D	253	LYS	2.2
1	D	137	LEU	2.2
1	C	248	LYS	2.2
1	C	111	GLU	2.1
1	D	56	PHE	2.1
1	C	78	THR	2.1
1	C	199	LYS	2.1
1	D	213	HIS	2.1
1	A	41	GLY	2.1
1	C	91	ARG	2.0
1	D	231	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	SO4	A	302	5/5	0.53	0.38	85,88,96,104	0
2	SO4	D	302	5/5	0.82	0.30	96,98,101,104	0
3	CA	C	303	1/1	0.89	0.18	70,70,70,70	0
2	SO4	C	302	5/5	0.91	0.25	74,75,78,81	0
2	SO4	B	301	5/5	0.92	0.16	48,55,57,60	0
2	SO4	D	301	5/5	0.92	0.26	76,79,81,85	0
2	SO4	C	301	5/5	0.94	0.18	47,54,61,67	0
3	CA	A	303	1/1	0.96	0.22	61,61,61,61	0
2	SO4	A	301	5/5	0.97	0.18	32,38,45,52	0
2	SO4	B	302	5/5	0.97	0.16	39,50,53,56	0

## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.