



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 10, 2024 – 12:13 AM JST

PDB ID : 8WCG
Title : Crystal structure of SARS-CoV-1 RBD in complex with nanobody aSR29 and aSR347
Authors : Zeng, W.H.; Ma, H.; Jin, T.C.
Deposited on : 2023-09-12
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.002 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.38.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	3775 (2.60-2.60)
Clashscore	180529	4181 (2.60-2.60)
Ramachandran outliers	177936	4129 (2.60-2.60)
Sidechain outliers	177891	4129 (2.60-2.60)
RSRZ outliers	164620	3775 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	197	
1	B	197	
2	C	127	
2	D	127	
3	E	131	
3	b	131	

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 7111 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Spike protein S1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	195	1541	993	253	286	9	0	0	0
1	B	194	1534	989	252	284	9	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	436	HIS	TYR	conflict	UNP P59594
B	436	HIS	TYR	conflict	UNP P59594

- Molecule 2 is a protein called aSR29 nanobody.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	C	126	970	609	170	188	3	0	0	0
2	D	126	970	609	170	188	3	0	0	0

- Molecule 3 is a protein called aSR347 nanobody.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	b	131	992	617	168	200	7	0	0	0
3	E	129	976	609	166	194	7	0	0	0

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
4	A	1	15	8	1	6	0	0
4	B	1	15	8	1	6	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
5	A	19	19	19	0	0
5	C	9	9	9	0	0
5	B	30	30	30	0	0
5	D	8	8	8	0	0
5	b	17	17	17	0	0
5	E	15	15	15	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Spike protein S1




- Molecule 1: Spike protein S1




- Molecule 2: aSR29 nanobody




- Molecule 2: aSR29 nanobody



- Molecule 3: aSR347 nanobody



- Molecule 3: aSR347 nanobody



4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	76.85Å 134.68Å 136.69Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.97 – 2.60 47.97 – 2.60	Depositor EDS
% Data completeness (in resolution range)	98.8 (47.97-2.60) 91.1 (47.97-2.60)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.73 (at 2.62Å)	Xtrriage
Refinement program	PHENIX (dev_2481: ???)	Depositor
R, R_{free}	0.232 , 0.279 0.232 , 0.279	Depositor DCC
R_{free} test set	2209 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	47.8	Xtrriage
Anisotropy	0.489	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 32.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.012 for -h,l,k	Xtrriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	7111	wwPDB-VP
Average B, all atoms (Å ²)	57.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.23% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.26	0/1590	0.44	0/2173
1	B	0.26	0/1583	0.44	0/2163
2	C	0.28	0/991	0.48	0/1344
2	D	0.28	0/991	0.46	0/1344
3	E	0.26	0/997	0.44	0/1354
3	b	0.28	0/1013	0.45	0/1375
All	All	0.27	0/7165	0.45	0/9753

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1541	0	1448	20	0
1	B	1534	0	1439	21	0
2	C	970	0	914	5	0
2	D	970	0	914	14	0
3	E	976	0	926	7	0
3	b	992	0	940	0	0
4	A	15	0	15	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	15	0	13	0	0
5	A	19	0	0	0	0
5	B	30	0	0	0	0
5	C	9	0	0	0	0
5	D	8	0	0	1	0
5	E	15	0	0	0	0
5	b	17	0	0	0	0
All	All	7111	0	6609	62	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:11:LEU:HD23	2:D:124:THR:HB	1.41	0.99
2:D:91:THR:HG22	2:D:125:VAL:H	1.35	0.90
1:B:439:LYS:CD	1:B:480:ASP:HB3	2.06	0.85
1:B:439:LYS:HD3	1:B:480:ASP:HB3	1.70	0.72
2:D:11:LEU:CD2	2:D:124:THR:HB	2.22	0.69
1:B:468:THR:HG22	1:B:470:PRO:HD2	1.76	0.66
1:B:439:LYS:NZ	1:B:480:ASP:HB3	2.11	0.65
1:A:414:ASP:OD2	2:C:27:ARG:NH1	2.34	0.60
1:A:487:THR:HB	1:A:491:TYR:HD2	1.68	0.58
1:A:337:VAL:HG11	1:A:389:VAL:HG12	1.86	0.58
1:B:361:PHE:O	2:D:107:ARG:NH2	2.37	0.57
1:A:351:ASP:HA	1:A:513:PRO:HG3	1.87	0.56
1:A:425:THR:O	1:A:493:PRO:HG2	2.06	0.55
1:B:469:PRO:HG3	1:B:476:TRP:CZ2	2.42	0.55
1:B:337:VAL:HA	1:B:387:PHE:HB2	1.89	0.54
3:E:83:MET:C	3:E:84:ASN:HD22	2.10	0.54
1:B:439:LYS:HZ3	1:B:480:ASP:HB3	1.72	0.54
1:A:396:GLN:HA	1:A:401:GLN:HG2	1.88	0.54
1:A:431:THR:HG22	1:A:433:THR:H	1.72	0.53
1:A:443:LEU:HD21	2:D:25:PRO:HG3	1.90	0.53
3:E:30:ALA:O	3:E:53:SER:OG	2.28	0.52
2:C:11:LEU:HG	2:C:124:THR:OG1	2.10	0.51
1:A:425:THR:HG21	1:A:495:ARG:HG3	1.94	0.50
1:A:337:VAL:HG12	1:A:387:PHE:CD1	2.47	0.49
3:E:91:THR:HG23	3:E:128:THR:HA	1.95	0.49
1:B:388:VAL:HG22	1:B:495:ARG:HG2	1.95	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:335:PRO:HG3	1:B:341:GLU:HG3	1.95	0.48
2:D:110:SER:H	2:D:114:TRP:HD1	1.60	0.48
1:A:351:ASP:O	1:A:354:VAL:HG12	2.14	0.48
3:E:67:ARG:HD2	3:E:85:ILE:HG23	1.95	0.48
1:B:321:ASN:OD1	1:B:321:ASN:N	2.46	0.48
2:C:6:GLU:OE1	2:C:6:GLU:N	2.46	0.48
2:D:8:GLY:O	2:D:18:LEU:HD11	2.13	0.48
2:D:67:ARG:NH2	2:D:90:ASP:OD2	2.48	0.47
1:A:484:TYR:HE2	2:D:11:LEU:HD21	1.80	0.47
3:E:82:GLN:HE21	3:E:84:ASN:ND2	2.13	0.47
1:A:433:THR:O	2:D:11:LEU:HD11	2.14	0.47
1:B:439:LYS:NZ	1:B:480:ASP:OD2	2.47	0.47
1:B:439:LYS:HD2	1:B:480:ASP:HB3	1.94	0.47
1:A:330:ASN:OD1	4:A:601:NAG:O1	2.32	0.46
1:B:426:ARG:NH1	1:B:485:THR:O	2.33	0.46
1:B:370:SER:HB3	1:B:373:LYS:HB2	1.98	0.46
1:B:439:LYS:HA	1:B:480:ASP:HA	1.97	0.45
2:D:3:GLN:NE2	5:D:203:HOH:O	2.49	0.45
1:B:407:ASP:OD1	1:B:411:LYS:NZ	2.49	0.45
1:B:363:THR:HG23	1:B:365:LYS:HD3	1.99	0.44
1:B:366:CYS:HA	1:B:419:CYS:HA	2.00	0.44
1:A:347:ASN:H	1:A:509:THR:HB	1.83	0.44
2:C:91:THR:HG23	2:C:124:THR:HG22	1.99	0.43
1:A:426:ARG:O	1:A:430:ALA:HB3	2.19	0.43
2:D:122:GLN:NE2	2:D:124:THR:OG1	2.52	0.43
1:A:353:SER:O	1:A:357:ASN:HB2	2.19	0.43
2:C:12:VAL:O	2:C:125:VAL:HA	2.19	0.42
1:B:439:LYS:CE	1:B:480:ASP:HB3	2.49	0.42
1:A:366:CYS:HA	1:A:419:CYS:HA	2.01	0.42
2:D:30:LYS:O	2:D:53:PHE:HB2	2.20	0.42
3:E:40:ALA:HB3	3:E:43:LYS:HB2	2.00	0.42
1:A:461:SER:HB2	1:A:467:CYS:SG	2.60	0.42
1:B:352:TYR:CD2	1:B:374:LEU:HB3	2.55	0.41
3:E:82:GLN:HE21	3:E:84:ASN:HD21	1.66	0.41
2:D:23:THR:HG23	2:D:77:ASN:O	2.20	0.41
1:A:388:VAL:HG22	1:A:495:ARG:HG2	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	193/197 (98%)	184 (95%)	9 (5%)	0	100	100
1	B	192/197 (98%)	181 (94%)	11 (6%)	0	100	100
2	C	124/127 (98%)	118 (95%)	6 (5%)	0	100	100
2	D	124/127 (98%)	118 (95%)	6 (5%)	0	100	100
3	E	127/131 (97%)	122 (96%)	4 (3%)	1 (1%)	16	34
3	b	129/131 (98%)	121 (94%)	8 (6%)	0	100	100
All	All	889/910 (98%)	844 (95%)	44 (5%)	1 (0%)	48	71

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	E	118	TYR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	167/172 (97%)	163 (98%)	4 (2%)	44	70
1	B	166/172 (96%)	161 (97%)	5 (3%)	36	63
2	C	99/104 (95%)	96 (97%)	3 (3%)	36	63
2	D	99/104 (95%)	94 (95%)	5 (5%)	20	42
3	E	104/107 (97%)	100 (96%)	4 (4%)	28	54
3	b	107/107 (100%)	102 (95%)	5 (5%)	22	45

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	742/766 (97%)	716 (96%)	26 (4%)	31 57

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	362	SER
1	A	364	PHE
1	A	395	ARG
1	A	414	ASP
2	C	57	SER
2	C	62	ASP
2	C	107	ARG
1	B	364	PHE
1	B	411	LYS
1	B	426	ARG
1	B	472	LEU
1	B	491	TYR
2	D	21	SER
2	D	56	ASP
2	D	67	ARG
2	D	96	CYS
2	D	110	SER
3	b	18	LEU
3	b	25	SER
3	b	69	SER
3	b	100	ASP
3	b	113	MET
3	E	44	GLU
3	E	77	ASN
3	E	108	GLN
3	E	111	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	321	ASN
1	B	321	ASN
2	D	82	GLN
3	b	82	GLN
3	E	84	ASN
3	E	123	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	A	601	-	15,15,15	0.54	0	21,21,21	0.83	2 (9%)
4	NAG	B	600	1	15,15,15	1.27	1 (6%)	21,21,21	1.64	5 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	601	-	-	0/6/26/26	0/1/1/1
4	NAG	B	600	1	-	2/6/26/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	600	NAG	C1-C2	-4.32	1.47	1.52

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	600	NAG	O5-C5-C4	3.87	116.73	109.69
4	B	600	NAG	C1-O5-C5	3.74	120.72	113.66
4	B	600	NAG	O1-C1-O5	-2.85	101.84	110.38
4	B	600	NAG	C1-C2-C3	-2.66	106.92	110.54
4	B	600	NAG	C3-C4-C5	2.49	114.69	110.24
4	A	601	NAG	C3-C4-C5	2.48	114.67	110.24
4	A	601	NAG	O5-C5-C4	2.19	113.66	109.69

There are no chirality outliers.

All (2) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
4	B	600	NAG	C4-C5-C6-O6
4	B	600	NAG	O5-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	601	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	195/197 (98%)	0.63	18 (9%) 16 13	40, 51, 71, 92	0
1	B	194/197 (98%)	0.51	14 (7%) 23 18	39, 52, 69, 79	0
2	C	126/127 (99%)	1.04	21 (16%) 5 4	42, 57, 76, 87	0
2	D	126/127 (99%)	1.18	24 (19%) 4 3	46, 63, 94, 98	0
3	E	129/131 (98%)	0.88	19 (14%) 7 5	46, 58, 71, 83	0
3	b	131/131 (100%)	0.72	14 (10%) 12 10	44, 53, 71, 93	0
All	All	901/910 (99%)	0.79	110 (12%) 10 7	39, 55, 79, 98	0

All (110) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	C	108	TYR	9.5
2	C	109	PHE	8.1
2	D	109	PHE	6.2
2	C	25	PRO	5.2
2	D	112	ARG	5.1
2	C	105	SER	5.1
1	B	331	ALA	5.1
1	B	469	PRO	5.0
2	C	106	ALA	4.8
1	B	504	LEU	4.6
2	D	25	PRO	4.5
3	E	118	TYR	4.4
1	B	470	PRO	4.4
1	A	470	PRO	4.2
2	D	126	SER	4.2
2	C	107	ARG	4.2
2	C	26	GLY	3.9
3	E	112	SER	3.9
2	C	112	ARG	3.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	b	118	TYR	3.9
1	A	469	PRO	3.8
1	A	514	LYS	3.7
2	C	111	GLU	3.7
1	A	513	PRO	3.6
3	b	89	GLU	3.6
2	D	108	TYR	3.6
3	b	119	HIS	3.6
3	E	111	CYS	3.6
2	C	1	GLN	3.5
3	b	27	GLY	3.5
2	D	113	GLU	3.3
3	E	26	GLU	3.3
2	C	8	GLY	3.3
1	B	330	ASN	3.2
2	C	110	SER	3.2
2	D	13	GLN	3.1
3	E	25	SER	3.1
3	b	120	ASP	3.1
3	E	41	PRO	3.0
2	D	94	TYR	3.0
2	D	11	LEU	3.0
2	D	124	THR	3.0
2	C	24	ALA	2.9
3	E	85	ILE	2.9
3	E	119	HIS	2.9
2	D	12	VAL	2.8
2	D	86	LEU	2.8
1	A	360	PHE	2.8
1	A	320	THR	2.7
2	C	28	THR	2.7
3	b	24	ALA	2.7
2	D	91	THR	2.7
3	b	131	SER	2.7
2	D	39	GLN	2.7
3	E	13	GLN	2.7
3	E	40	ALA	2.6
2	D	65	LYS	2.6
2	C	126	SER	2.6
2	D	18	LEU	2.6
3	E	18	LEU	2.6
1	A	414	ASP	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	E	17	SER	2.5
1	A	321	ASN	2.5
3	E	129	VAL	2.5
3	b	130	SER	2.5
3	E	110	PRO	2.5
1	B	332	THR	2.5
3	b	1	GLN	2.5
3	E	27	GLY	2.5
1	A	428	ILE	2.4
2	D	3	GLN	2.4
1	B	468	THR	2.4
1	B	325	PHE	2.4
3	b	28	PHE	2.4
1	A	491	TYR	2.4
2	D	110	SER	2.4
3	E	116	ASN	2.4
1	B	329	PHE	2.3
2	C	113	GLU	2.3
2	D	123	VAL	2.3
3	E	87	LYS	2.3
2	C	10	GLY	2.3
2	D	23	THR	2.3
2	D	15	GLY	2.2
2	C	104	ASP	2.2
2	C	114	TRP	2.2
1	B	491	TYR	2.2
3	E	28	PHE	2.2
2	D	14	ALA	2.2
1	A	343	LYS	2.2
1	B	514	LYS	2.2
1	B	506	ALA	2.2
1	A	322	LEU	2.2
1	B	322	LEU	2.2
1	A	431	THR	2.2
2	C	63	THR	2.2
1	A	468	THR	2.1
1	A	345	ILE	2.1
1	A	356	TYR	2.1
3	E	114	THR	2.1
1	A	328	VAL	2.1
2	D	114	TRP	2.1
3	b	102	MET	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	C	11	LEU	2.1
3	b	129	VAL	2.1
2	D	107	ARG	2.1
3	b	88	PRO	2.0
1	B	328	VAL	2.0
1	A	359	THR	2.0
3	b	51	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NAG	B	600	15/15	0.52	0.20	63,87,91,91	0
4	NAG	A	601	15/15	0.69	0.15	70,85,93,94	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around NAG B 600:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.