

Full wwPDB X-ray Structure Validation Report (i)

Nov 22, 2023 – 01:24 PM JST

PDB ID : 7WDI

Title: Crystal structure of the P450 BM3 heme domain mutant F87K in complex

with N-imidazolyl-hexanoyl-L-phenylalanine and hydroxylamine

Authors : Jiang, Y.; Dong, S.; Feng, Y.; Cong, Z.

Deposited on : 2021-12-21

Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.36

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY\ DIFFRACTION$

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},{\rm resolution\ range}({\rm \AA})) \end{array}$
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain					
1	A	465	86%	9%	5%			
1	В	465	84%	10%	6%			



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 7912 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Bifunctional cytochrome P450/NADPH-P450 reductase.


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	Λ	444	Total	С	N	О	S	0	0	0
1	Λ	444	3489	2234	597	641	17	U		
1	D	439	Total	С	N	О	S	0	0	0
1	Б	409	3450	2213	588	632	17	0		

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	expression tag	UNP A0A1Q8UP87
A	87	LYS	PHE	engineered mutation	UNP A0A1Q8UP87
A	456	LEU	-	expression tag	UNP A0A1Q8UP87
A	457	GLU	-	expression tag	UNP A0A1Q8UP87
A	458	HIS	-	expression tag	UNP A0A1Q8UP87
A	459	HIS	-	expression tag	UNP A0A1Q8UP87
A	460	HIS	-	expression tag	UNP A0A1Q8UP87
A	461	HIS	-	expression tag	UNP A0A1Q8UP87
A	462	HIS	-	expression tag	UNP A0A1Q8UP87
A	463	HIS	-	expression tag	UNP A0A1Q8UP87
В	-1	GLY	-	expression tag	UNP A0A1Q8UP87
В	87	LYS	PHE	engineered mutation	UNP A0A1Q8UP87
В	456	LEU	-	expression tag	UNP A0A1Q8UP87
В	457	GLU	-	expression tag	UNP A0A1Q8UP87
В	458	HIS	-	expression tag	UNP A0A1Q8UP87
В	459	HIS	-	expression tag	UNP A0A1Q8UP87
В	460	HIS	-	expression tag	UNP A0A1Q8UP87
В	461	HIS	-	expression tag	UNP A0A1Q8UP87
В	462	HIS	-	expression tag	UNP A0A1Q8UP87
В	463	HIS	-	expression tag	UNP A0A1Q8UP87


• Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: C₃₄H₃₂FeN₄O₄) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf		
2	Λ	1	Total	С	Fe	N	О	0	0	
$\begin{array}{ c c c c c } \hline Z & A \\ \hline \end{array}$	Λ	1	43	34	1	4	4			
2	B	1	Total	С	Fe	N	О	0	0	
2	D	1	43	34	1	4	4			

• Molecule 3 is (2S)-2-(6-imidazol-1-ylhexanoylamino)-3-phenyl-propanoic acid (three-letter code: IC6) (formula: $C_{18}H_{23}N_3O_3$) (labeled as "Ligand of Interest" by depositor).




\mathbf{M}	ol	Chain	Residues	Atoms				ZeroOcc	AltConf
3		A	1	Total 24	C 18	N 3	O 3	0	1



Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
3	Δ	1	Total				0	1	
	3 A	1	24	18	3	3	O	1	
3	В	1	Total	С	Ν	Ο	0	1	
9	Ъ	1	24	18	3	3	0		
2	D	1	Total	С	N	О	0	1	
3	Ъ	1	24	18	3	3	U	1	

 \bullet Molecule 4 is HYDROXYAMINE (three-letter code: HOA) (formula: H₃NO) (labeled as "Ligand of Interest" by depositor).



\mathbf{Mol}	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total N O 2 1 1	0	0
4	В	1	Total N O 2 1 1	0	0

• Molecule 5 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	376	Total O 376 376	0	0
5	В	411	Total O 411 411	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Bifunctional cytochrome P450/NADPH-P450 reductase



• Molecule 1: Bifunctional cytochrome P450/NADPH-P450 reductase





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	58.41Å 149.02Å 65.28Å	Donositon
a, b, c, α , β , γ	90.00° 100.35° 90.00°	Depositor
Resolution (Å)	29.25 - 2.10	Depositor
Resolution (A)	29.25 - 2.10	EDS
% Data completeness	96.2 (29.25-2.10)	Depositor
(in resolution range)	96.2 (29.25-2.10)	EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.33 (at 2.10Å)	Xtriage
Refinement program	PHENIX 1.18.2_3874	Depositor
D D	0.183 , 0.219	Depositor
R, R_{free}	0.184 , 0.218	DCC
R_{free} test set	1999 reflections (3.24%)	wwPDB-VP
Wilson B-factor (Å ²)	25.3	Xtriage
Anisotropy	0.047	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.35, 55.1	EDS
L-test for twinning ²	$ < L >=0.48, < L^2>=0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7912	wwPDB-VP
Average B, all atoms $(Å^2)$	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.04% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: HOA, HEM, IC6

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
IVIOI		RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.34	0/3568	0.51	0/4830	
1	В	0.32	0/3529	0.49	0/4775	
All	All	0.33	0/7097	0.50	0/9605	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3489	0	3401	27	0
1	В	3450	0	3371	38	0
2	A	43	0	30	1	0
2	В	43	0	30	2	0
3	A	48	0	0	0	0
3	В	48	0	0	1	0
4	A	2	0	0	0	0
4	В	2	0	0	0	0
5	A	376	0	0	5	0
5	В	411	0	0	7	0
All	All	7912	0	6832	64	0



The all-atom clash score is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clash score for this structure is 5.

All (64) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic	Clash
4 4 40 1 511 115 04	4 4 440 1141 11604	distance (Å)	overlap (Å)
1:A:148:LEU:HD21	1:A:413:VAL:HG21	1.60	0.80
1:B:134:ASN:HB2	1:B:137:GLU:OE2	1.81	0.80
1:B:150:LEU:HD22	1:B:174:ILE:HD11	1.65	0.77
1:A:237:MET:HE1	1:A:258:ILE:HG13	1.73	0.70
1:A:231:ASP:CA	5:A:812:HOH:O	2.39	0.69
1:B:134:ASN:ND2	1:B:137:GLU:OE2	2.23	0.67
1:B:306:LYS:HG2	5:B:847:HOH:O	1.97	0.65
1:B:230:SER:N	5:B:607:HOH:O	2.29	0.65
1:A:296:ARG:HD2	5:A:602:HOH:O	2.01	0.60
1:B:103:LEU:HD21	1:B:237:MET:HG2	1.86	0.58
1:B:134:ASN:HB2	1:B:137:GLU:CD	2.23	0.58
1:B:120:VAL:HG11	1:B:302:VAL:HG13	1.85	0.58
1:B:216:VAL:HG11	1:B:255:ARG:HG3	1.87	0.57
1:B:312:LYS:HB2	5:B:864:HOH:O	2.04	0.57
1:A:367:TRP:HB2	1:A:371:VAL:HG12	1.85	0.57
1:B:170:PRO:HB3	1:B:174:ILE:HD12	1.88	0.56
1:A:87:LYS:NZ	5:A:612:HOH:O	2.36	0.54
1:B:47:ARG:NH1	3:B:503[B]:IC6:O15	2.35	0.53
1:B:223:ARG:HG2	5:B:656:HOH:O	2.08	0.53
1:B:43:GLU:HG2	1:B:48:VAL:HG22	1.91	0.52
1:A:120:VAL:HG11	1:A:302:VAL:HG13	1.91	0.52
1:A:175:THR:HG22	1:A:179:ARG:HH12	1.74	0.51
1:A:216:VAL:HG21	1:A:259:ILE:HG13	1.91	0.51
1:B:158:PHE:HE2	1:B:258:ILE:HG12	1.74	0.51
1:A:171:HIS:CG	1:A:172:PRO:HD2	2.46	0.50
1:A:213:ASN:OD1	1:A:255:ARG:HD2	2.12	0.50
1:A:388:HIS:HA	1:A:391:LYS:HD3	1.94	0.50
1:A:179:ARG:NH2	1:A:208:ASP:OD1	2.45	0.49
1:A:21:ASN:HA	1:A:189:GLN:HB2	1.95	0.49
1:B:158:PHE:CE2	1:B:258:ILE:HG12	2.47	0.49
1:A:368:GLY:O	1:A:371:VAL:HG13	2.13	0.49
1:A:59:LYS:HE3	1:B:296:ARG:HD3	1.95	0.48
1:B:73:GLN:HB3	1:B:188:LEU:HD21	1.96	0.48
1:A:253:ASN:O	1:A:257:GLN:HG2	2.14	0.48
1:A:298:LEU:HD22	1:A:303:PRO:HB3	1.96	0.48
1:B:253:ASN:O	1:B:257:GLN:HG2	2.16	0.45
1:B:298:LEU:HD22	1:B:303:PRO:HB3	1.98	0.45



Continued from previous page...

Continued from prec		Interatomic	Clash
Atom-1	Atom-2	${\rm distance}\ ({\rm \AA})$	overlap (Å)
1:B:393:PHE:HB3	1:B:400:CYS:HB3	1.98	0.45
1:B:382:PRO:O	5:B:601:HOH:O	2.21	0.45
1:B:185:MET:O	1:B:188:LEU:HB2	2.17	0.45
1:B:134:ASN:CB	1:B:137:GLU:OE2	2.61	0.45
1:B:141:VAL:HB	1:B:142:PRO:HD3	2.00	0.44
1:A:351:ASP:OD1	5:A:601:HOH:O	2.21	0.44
1:B:150:LEU:HD22	1:B:174:ILE:CD1	2.41	0.44
1:B:388:HIS:HA	1:B:391:LYS:HD3	1.98	0.44
1:A:81:PHE:HB3	1:A:209:ILE:HG12	2.00	0.43
1:B:223:ARG:NH2	1:B:232:ASP:OD2	2.28	0.43
1:A:173:PHE:CE1	1:A:177:MET:SD	3.12	0.43
1:B:130:TRP:CD2	1:B:139:ILE:HD13	2.54	0.43
1:B:134:ASN:H	1:B:137:GLU:CD	2.23	0.42
1:A:26:VAL:O	1:A:30:MET:HG3	2.20	0.42
1:A:366:ILE:HG21	1:A:389:ALA:HB1	2.01	0.42
1:A:111:ALA:HA	5:A:672:HOH:O	2.19	0.42
1:A:400:CYS:HB2	2:A:501:HEM:NA	2.34	0.42
1:B:434:LYS:HB2	1:B:442:GLU:HB2	2.01	0.41
1:B:177:MET:SD	1:B:263:ILE:HG12	2.61	0.41
1:B:265:GLY:HA2	2:B:501:HEM:C2C	2.56	0.41
1:B:282:LYS:NZ	1:B:425:ASP:OD2	2.43	0.41
1:B:312:LYS:NZ	5:B:622:HOH:O	2.46	0.41
1:B:204:GLN:OE1	5:B:603:HOH:O	2.22	0.41
1:A:266:HIS:CG	1:A:267:GLU:N	2.88	0.40
1:B:118:MET:HE3	1:B:118:MET:HB2	1.92	0.40
1:B:261:PHE:HD1	2:B:501:HEM:HAC	1.85	0.40
1:A:440:LYS:HB3	1:A:440:LYS:HE3	1.87	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	438/465~(94%)	423 (97%)	15 (3%)	0	100	100
1	В	433/465 (93%)	418 (96%)	15 (4%)	0	100	100
All	All	871/930 (94%)	841 (97%)	30 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percent	iles
1	A	366/408 (90%)	366 (100%)	0	100 1	00
1	В	362/408~(89%)	362 (100%)	0	100 1	00
All	All	728/816~(89%)	728 (100%)	0	100 1	00

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.



5.6 Ligand geometry (i)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Tuno	Chain	Res	Link	Во	ond leng	ths	В	ond ang	les
MIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	IC6	В	502[A]	-	24,25,25	1.47	3 (12%)	29,31,31	0.89	2 (6%)
3	IC6	A	502[A]	-	24,25,25	1.51	3 (12%)	29,31,31	0.98	2 (6%)
2	HEM	A	501	1,4	41,50,50	1.46	4 (9%)	45,82,82	1.55	7 (15%)
3	IC6	В	503[B]	-	24,25,25	1.49	3 (12%)	29,31,31	1.11	2 (6%)
3	IC6	A	503[B]	-	24,25,25	1.57	4 (16%)	29,31,31	1.42	5 (17%)
4	HOA	A	504	2	0,1,1	-	-	-		
4	HOA	В	504	2	0,1,1	-	-	-		
2	HEM	В	501	1,4	41,50,50	1.48	5 (12%)	45,82,82	1.47	9 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	IC6	В	502[A]	-	-	3/20/20/20	0/2/2/2
3	IC6	A	502[A]	-	-	1/20/20/20	0/2/2/2
2	HEM	A	501	1,4	-	2/12/54/54	-
3	IC6	В	503[B]	-	-	5/20/20/20	0/2/2/2
3	IC6	A	503[B]	-	-	6/20/20/20	0/2/2/2
2	HEM	В	501	1,4	-	2/12/54/54	-

All (22) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\text{\AA})$	Ideal(Å)
3	A	503[B]	IC6	C11-N12	5.79	1.46	1.34
3	A	502[A]	IC6	C11-N12	5.62	1.46	1.34
3	В	503[B]	IC6	C11-N12	5.54	1.45	1.34
3	В	502[A]	IC6	C11-N12	5.48	1.45	1.34



Continued from previous page...

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(\mathbf{\mathring{A}})$	$Ideal(\AA)$
2	A	501	HEM	C3C-C2C	-3.91	1.34	1.40
2	В	501	HEM	C3C-C2C	-3.78	1.35	1.40
2	A	501	HEM	C3C-CAC	3.76	1.55	1.47
2	В	501	HEM	C3C-CAC	3.72	1.55	1.47
2	В	501	HEM	CAB-C3B	2.99	1.55	1.47
2	A	501	HEM	CAB-C3B	2.96	1.55	1.47
3	В	502[A]	IC6	C4-N5	-2.67	1.32	1.37
3	A	502[A]	IC6	C4-N5	-2.66	1.32	1.37
3	В	503[B]	IC6	C4-N5	-2.57	1.32	1.37
3	A	503[B]	IC6	C4-N5	-2.49	1.33	1.37
3	A	502[A]	IC6	O24-C11	-2.19	1.18	1.23
2	В	501	HEM	CAA-C2A	2.16	1.55	1.52
3	В	502[A]	IC6	O24-C11	-2.14	1.18	1.23
3	A	503[B]	IC6	C10-C11	2.10	1.55	1.51
3	A	503[B]	IC6	O24-C11	-2.04	1.19	1.23
2	В	501	HEM	CMB-C2B	2.03	1.55	1.50
3	В	503[B]	IC6	O24-C11	-2.01	1.19	1.23
2	A	501	HEM	CMD-C2D	2.00	1.55	1.50

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$Ideal(^{o})$
2	A	501	HEM	CBA-CAA-C2A	-4.14	105.55	112.62
3	A	503[B]	IC6	C10-C11-N12	3.92	122.63	115.83
3	В	503[B]	IC6	C14-C13-N12	-3.75	101.66	110.55
2	В	501	HEM	C1B-NB-C4B	3.25	108.43	105.07
3	A	503[B]	IC6	C14-C13-N12	-3.20	102.97	110.55
2	A	501	HEM	C4D-ND-C1D	2.97	108.14	105.07
2	A	501	HEM	C1B-NB-C4B	2.94	108.11	105.07
2	В	501	HEM	C4D-ND-C1D	2.94	108.11	105.07
2	A	501	HEM	C4B-CHC-C1C	2.87	126.34	122.56
2	В	501	HEM	CBA-CAA-C2A	-2.71	107.99	112.62
3	A	503[B]	IC6	O24-C11-N12	-2.68	118.42	122.95
2	В	501	HEM	C4B-CHC-C1C	2.65	126.06	122.56
2	В	501	HEM	C4C-CHD-C1D	2.59	125.97	122.56
3	В	503[B]	IC6	O15-C14-C13	2.56	121.89	113.40
3	A	502[A]	IC6	O15-C14-C13	2.33	121.15	113.40
3	A	503[B]	IC6	C09-C10-C11	2.32	119.78	113.26
2	A	501	HEM	C3D-C4D-ND	-2.26	107.65	110.17
2	A	501	HEM	C4C-CHD-C1D	2.18	125.44	122.56
3	A	503[B]	IC6	C17-C13-N12	2.17	115.36	110.79
2	В	501	HEM	C3B-C2B-C1B	2.17	108.10	106.49



Continued from previous page...

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
2	В	501	HEM	C3D-C4D-ND	-2.16	107.76	110.17
2	A	501	HEM	CMC-C2C-C3C	2.15	128.70	124.68
3	В	502[A]	IC6	O15-C14-C13	2.15	120.53	113.40
2	В	501	HEM	CMC-C2C-C3C	2.11	128.63	124.68
3	A	502[A]	IC6	C10-C11-N12	2.10	119.47	115.83
2	В	501	HEM	C4A-C3A-C2A	2.10	108.45	107.00
3	В	502[A]	IC6	C10-C11-N12	2.02	119.33	115.83

There are no chirality outliers.

All (19) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	503[B]	IC6	C8-C09-C10-C11
3	A	503[B]	IC6	N5-C6-C7-C8
3	В	502[A]	IC6	C10-C09-C8-C7
3	A	503[B]	IC6	C09-C10-C11-N12
3	A	503[B]	IC6	C09-C10-C11-O24
3	A	503[B]	IC6	C10-C09-C8-C7
3	В	502[A]	IC6	C8-C09-C10-C11
3	A	503[B]	IC6	C6-C7-C8-C09
3	В	503[B]	IC6	C8-C09-C10-C11
3	В	503[B]	IC6	C6-C7-C8-C09
3	В	502[A]	IC6	N5-C6-C7-C8
3	В	503[B]	IC6	C10-C09-C8-C7
3	A	502[A]	IC6	C7-C6-N5-C4
3	В	503[B]	IC6	C09-C10-C11-O24
2	A	501	HEM	CAD-CBD-CGD-O2D
2	В	501	HEM	CAD-CBD-CGD-O2D
2	A	501	HEM	CAD-CBD-CGD-O1D
2	В	501	HEM	CAD-CBD-CGD-O1D
3	В	503[B]	IC6	C09-C10-C11-N12

There are no ring outliers.


3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	HEM	1	0
3	В	503[B]	IC6	1	0
2	В	501	HEM	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,




bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.






























5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle { m RSRZ} \rangle$	# RSRZ > 2		$OWAB(A^2)$	Q<0.9
1	A	444/465 (95%)	0.06	20 (4%) 33	38	16, 27, 53, 74	0
1	В	439/465 (94%)	0.12	27 (6%) 20	25	16, 27, 57, 68	0
All	All	883/930 (94%)	0.09	47 (5%) 26	32	16, 27, 55, 74	0

All (47) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	231	ASP	4.3
1	A	231	ASP	4.2
1	A	225	ALA	4.0
1	В	189	GLN	3.8
1	A	226	SER	3.6
1	В	201	ASN	3.4
1	A	201	ASN	3.4
1	A	2	ILE	3.3
1	A	369	ASP	3.3
1	A	302	VAL	3.2
1	В	186	ASN	3.1
1	В	205	PHE	3.1
1	В	188	LEU	3.1
1	В	221	ALA	3.0
1	В	109	GLN	2.9
1	A	230	SER	2.9
1	В	203	ARG	2.9
1	A	3	LYS	2.8
1	В	214	ASP	2.7
1	A	109	GLN	2.7
1	В	3	LYS	2.7
1	В	181	LEU	2.7
1	A	218	LYS	2.7
1	В	222	ASP	2.7



Continued from previous page...

Mol	Chain	Res	Type	RSRZ	
1	В	2	ILE	2.6	
1	В	262	LEU	2.6	
1	В	169	GLN	2.6	
1	A	190	ARG	2.6	
1	A	108	SER	2.5	
1	В	175	THR	2.5	
1	A	200	GLU	2.5	
1	В	200	GLU	2.5	
1	В	211	VAL	2.4	
1	A	299	VAL	2.4	
1	A	178	VAL	2.4	
1	В	202	LYS	2.3	
1	В	172	PRO	2.3	
1	В	230	SER	2.3	
1	A	179	ARG	2.3	
1	A	229	GLN	2.3	
1	В	185	MET	2.2	
1	В	206	GLN	2.2	
1	A	136	ASP	2.1	
1	В	105	PRO	2.1	
1	В	184	ALA	2.1	
1	A	184	ALA	2.0	
1	В	174	ILE	2.0	

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




 $oxed{Atoms} oxed{ RSCC} oxed{ RSR} oxed{ B-factors(\AA^2)} oxed{ Q<0.9}$

Continued from previous page...

Mol | Type | Chain | Res

							, ,	-
Mol	Type	Chain	Res	Atoms	RSCC	RSR	${f B-factors}({f \AA}^2)$	Q<0.9
3	IC6	A	502[A]	24/24	0.91	0.25	12,19,22,23	24
3	IC6	В	503[B]	24/24	0.91	0.20	17,21,24,27	24
3	IC6	В	502[A]	24/24	0.92	0.24	15,20,23,24	24
3	IC6	A	503[B]	24/24	0.93	0.20	17,20,23,24	24
4	HOA	В	504	2/2	0.95	0.19	23,23,23,25	0
2	HEM	В	501	43/43	0.98	0.14	16,21,24,26	0
2	HEM	A	501	43/43	0.98	0.12	14,19,22,26	0
4	HOA	A	504	2/2	0.99	0.12	21,21,21,23	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around IC6 B 503 (B): 2mF_o-DF_c (at 0.7 rmsd) in gray mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive) Electron density around IC6 B 502 (A):

$2mF_o$ -DF_c (at 0.7 rmsd) in gray mF_o -DF_c (at 3 rmsd) in purple (negative) and green (positive)


























6.5 Other polymers (i)

There are no such residues in this entry.

