



# Full wwPDB X-ray Structure Validation Report ⓘ

Dec 18, 2024 – 04:15 PM JST

PDB ID : 8WEB  
Title : Crystal structure of Arabidopsis thaliana MIK2 ectodomain  
Authors : Wan, L.H.; Hu, Y.X.; Wu, H.M.  
Deposited on : 2023-09-17  
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.21  
EDS : 3.0  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.004 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.40


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	3187 (1.96-1.96)
Clashscore	180529	3412 (1.96-1.96)
Ramachandran outliers	177936	3390 (1.96-1.96)
Sidechain outliers	177891	3390 (1.96-1.96)
RSRZ outliers	164620	3186 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	712	
2	B	2	
2	C	2	
2	D	2	
2	E	2	
2	F	2	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	G	2	 50% 50%

## 2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 5909 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MDIS1-interacting receptor like kinase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	646	5009	3166	843	984	16	0	0	0

There are 15 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	137	GLU	LEU	engineered mutation	UNP Q8VZG8
A	164	LYS	ASP	engineered mutation	UNP Q8VZG8
A	564	PHE	SER	engineered mutation	UNP Q8VZG8
A	701	GLU	-	expression tag	UNP Q8VZG8
A	702	PHE	-	expression tag	UNP Q8VZG8
A	703	HIS	-	expression tag	UNP Q8VZG8
A	704	HIS	-	expression tag	UNP Q8VZG8
A	705	HIS	-	expression tag	UNP Q8VZG8
A	706	HIS	-	expression tag	UNP Q8VZG8
A	707	HIS	-	expression tag	UNP Q8VZG8
A	708	HIS	-	expression tag	UNP Q8VZG8
A	709	HIS	-	expression tag	UNP Q8VZG8
A	710	HIS	-	expression tag	UNP Q8VZG8
A	711	HIS	-	expression tag	UNP Q8VZG8
A	712	HIS	-	expression tag	UNP Q8VZG8

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	2	28	16	2	10	0	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	C	2	Total 28	C 16	N 2	O 10	0	0	0
2	D	2	Total 28	C 16	N 2	O 10	0	0	0
2	E	2	Total 28	C 16	N 2	O 10	0	0	0
2	F	2	Total 28	C 16	N 2	O 10	0	0	0
2	G	2	Total 28	C 16	N 2	O 10	0	0	0

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	Total 14	C 8	N 1	O 5	0	0
3	A	1	Total 14	C 8	N 1	O 5	0	0
3	A	1	Total 14	C 8	N 1	O 5	0	0
3	A	1	Total 14	C 8	N 1	O 5	0	0
3	A	1	Total 14	C 8	N 1	O 5	0	0


- Molecule 4 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
4	A	662	Total 662	O 662	0	0

### 3 Residue-property plots i

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: MDIS1-interacting receptor like kinase 2




- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose




- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucofuranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E:  50% 50%

MAG1  
MAG2

- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucofuranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:  50% 50%

MAG1  
MAG2

- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucofuranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G:  50% 50%

MAG1  
MAG2



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	62.13Å 107.89Å 164.31Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	35.13 – 1.95 35.13 – 1.95	Depositor EDS
% Data completeness (in resolution range)	98.5 (35.13-1.95) 98.4 (35.13-1.95)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.56 (at 1.95Å)	Xtrriage
Refinement program	PHENIX (1.20.1_4487: ???)	Depositor
R, $R_{free}$	0.171 , 0.201 0.171 , 0.201	Depositor DCC
$R_{free}$ test set	4004 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	19.5	Xtrriage
Anisotropy	0.019	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 49.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	5909	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	26.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.94% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.76	0/5111	0.75	0/6952

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5009	0	4964	32	0
2	B	28	0	25	0	0
2	C	28	0	25	0	0
2	D	28	0	25	0	0
2	E	28	0	25	0	0
2	F	28	0	25	1	0
2	G	28	0	25	1	0
3	A	70	0	65	6	0
4	A	662	0	0	16	1
All	All	5909	0	5179	40	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:434:ARG:NH2	4:A:1002:HOH:O	2.09	0.85
1:A:352:ASP:OD2	4:A:1001:HOH:O	2.02	0.78
4:A:1658:HOH:O	2:G:2:NAG:O4	2.02	0.77
1:A:553:GLU:OE2	1:A:575:ARG:NH1	2.20	0.75
1:A:77:ASN:ND2	4:A:1006:HOH:O	2.21	0.72
1:A:633:ASN:ND2	4:A:1004:HOH:O	2.18	0.70
1:A:137:GLU:OE2	1:A:140:ARG:NH2	2.25	0.69
1:A:366:ARG:HG2	1:A:388:GLN:OE1	1.93	0.69
1:A:436:ARG:NE	4:A:1008:HOH:O	2.24	0.68
1:A:469:GLN:NE2	4:A:1010:HOH:O	2.29	0.65
1:A:436:ARG:NH2	4:A:1012:HOH:O	2.30	0.64
1:A:273:GLN:HG3	1:A:297:LYS:HE2	1.81	0.63
3:A:903:NAG:H81	4:A:1600:HOH:O	1.98	0.62
1:A:284:ASN:ND2	4:A:1003:HOH:O	2.16	0.60
1:A:690:GLN:N	4:A:1017:HOH:O	2.38	0.56
3:A:905:NAG:O3	3:A:905:NAG:H83	2.06	0.55
1:A:150:SER:HB3	1:A:174:VAL:HG22	1.93	0.51
3:A:903:NAG:O7	3:A:903:NAG:O3	2.21	0.51
1:A:565:GLU:CD	1:A:565:GLU:H	2.14	0.51
1:A:436:ARG:CZ	4:A:1012:HOH:O	2.58	0.51
1:A:211:GLY:HA3	1:A:233:GLU:HB3	1.96	0.47
1:A:657:ASN:OD1	1:A:680:LYS:HB2	2.16	0.46
1:A:366:ARG:HG2	1:A:388:GLN:CD	2.36	0.45
1:A:113:PRO:HG2	1:A:117:LEU:HD11	1.98	0.45
1:A:473:ASN:ND2	4:A:1025:HOH:O	2.42	0.45
1:A:55:LEU:HG	1:A:85:TRP:HH2	1.82	0.45
1:A:137:GLU:O	1:A:140:ARG:HG2	2.16	0.45
1:A:74:VAL:HG13	1:A:75:ASN:H	1.82	0.44
1:A:530:LYS:HG3	1:A:554:TYR:HB3	2.00	0.44
4:A:1580:HOH:O	2:F:1:NAG:H61	2.18	0.44
1:A:527:ARG:HH11	1:A:527:ARG:HG3	1.83	0.43
1:A:565:GLU:HG3	4:A:1633:HOH:O	2.18	0.43
3:A:902:NAG:O7	3:A:902:NAG:O3	2.27	0.43
1:A:693:LYS:HD2	4:A:1204:HOH:O	2.19	0.43
1:A:487:ASN:HA	1:A:511:SER:O	2.19	0.42
3:A:904:NAG:H3	3:A:904:NAG:H83	2.03	0.41
1:A:215:LYS:HB3	1:A:215:LYS:HE3	1.81	0.41
1:A:470:LEU:HD23	1:A:470:LEU:HA	1.93	0.40
3:A:905:NAG:C8	3:A:905:NAG:C3	2.99	0.40
1:A:279:PRO:HA	1:A:280:PRO:HD3	1.98	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:1406:HOH:O	4:A:1471:HOH:O[4_545]	2.14	0.06

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	642/712 (90%)	612 (95%)	30 (5%)	0	<a href="#">100</a> <a href="#">100</a>

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	583/646 (90%)	580 (100%)	3 (0%)	<a href="#">86</a> <a href="#">86</a>

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	637	GLN
1	A	643	LYS
1	A	646	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	308	ASN
1	A	469	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

12 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	B	1	2,1	14,14,15	0.41	0	17,19,21	0.97	1 (5%)
2	NAG	B	2	2	14,14,15	0.20	0	17,19,21	0.49	0
2	NAG	C	1	2,1	14,14,15	0.32	0	17,19,21	0.53	0
2	NAG	C	2	2	14,14,15	1.47	2 (14%)	17,19,21	0.99	2 (11%)
2	NAG	D	1	2,1	14,14,15	0.76	1 (7%)	17,19,21	0.74	1 (5%)
2	NAG	D	2	2	14,14,15	0.34	0	17,19,21	0.63	1 (5%)
2	NAG	E	1	2,1	14,14,15	0.49	0	17,19,21	0.84	0
2	NAG	E	2	2	14,14,15	0.78	1 (7%)	17,19,21	0.69	0
2	NAG	F	1	2,1	14,14,15	0.91	1 (7%)	17,19,21	1.09	1 (5%)
2	NAG	F	2	2	14,14,15	0.47	0	17,19,21	0.90	1 (5%)
2	NAG	G	1	2,1	14,14,15	0.61	0	17,19,21	0.74	1 (5%)
2	NAG	G	2	2	14,14,15	0.64	0	17,19,21	1.04	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	B	2	2	-	2/6/23/26	0/1/1/1
2	NAG	C	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	C	2	2	-	2/6/23/26	0/1/1/1
2	NAG	D	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	D	2	2	-	2/6/23/26	0/1/1/1
2	NAG	E	1	2,1	-	1/6/23/26	0/1/1/1
2	NAG	E	2	2	-	0/6/23/26	0/1/1/1
2	NAG	F	1	2,1	-	2/6/23/26	0/1/1/1
2	NAG	F	2	2	-	0/6/23/26	0/1/1/1
2	NAG	G	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	G	2	2	-	2/6/23/26	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	2	NAG	C1-C2	3.64	1.57	1.52
2	C	2	NAG	O5-C1	-3.52	1.38	1.43
2	E	2	NAG	C1-C2	2.73	1.56	1.52
2	F	1	NAG	O5-C1	2.53	1.47	1.43
2	D	1	NAG	O5-C1	-2.13	1.40	1.43

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	G	2	NAG	C1-O5-C5	3.94	117.53	112.19
2	F	1	NAG	C1-O5-C5	3.50	116.94	112.19
2	F	2	NAG	C1-O5-C5	3.39	116.78	112.19
2	C	2	NAG	C4-C3-C2	2.62	114.86	111.02
2	B	1	NAG	C1-O5-C5	2.54	115.64	112.19
2	G	1	NAG	O4-C4-C5	-2.35	103.47	109.30
2	D	2	NAG	C1-O5-C5	2.11	115.06	112.19
2	D	1	NAG	C1-O5-C5	2.08	115.01	112.19
2	C	2	NAG	O5-C5-C4	-2.06	105.81	110.83

There are no chirality outliers.

All (11) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	2	NAG	O5-C5-C6-O6
2	G	2	NAG	O5-C5-C6-O6
2	C	2	NAG	C4-C5-C6-O6
2	G	2	NAG	C4-C5-C6-O6
2	D	2	NAG	C4-C5-C6-O6
2	F	1	NAG	C4-C5-C6-O6
2	B	2	NAG	C4-C5-C6-O6
2	C	2	NAG	O5-C5-C6-O6
2	D	2	NAG	O5-C5-C6-O6
2	F	1	NAG	O5-C5-C6-O6
2	E	1	NAG	C4-C5-C6-O6

There are no ring outliers.


2 monomers are involved in 2 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	F	1	NAG	1	0
2	G	2	NAG	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.

















## 5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	NAG	A	903	1	14,14,15	0.29	0	17,19,21	0.69	0
3	NAG	A	904	1	14,14,15	0.41	0	17,19,21	1.95	4 (23%)
3	NAG	A	902	1	14,14,15	0.49	0	17,19,21	0.85	1 (5%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	NAG	A	901	1	14,14,15	1.04	2 (14%)	17,19,21	0.51	0
3	NAG	A	905	1	14,14,15	0.68	0	17,19,21	2.97	5 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	903	1	-	2/6/23/26	0/1/1/1
3	NAG	A	904	1	-	3/6/23/26	0/1/1/1
3	NAG	A	902	1	-	2/6/23/26	0/1/1/1
3	NAG	A	901	1	-	0/6/23/26	0/1/1/1
3	NAG	A	905	1	-	5/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	901	NAG	O5-C1	2.54	1.47	1.43
3	A	901	NAG	C1-C2	2.51	1.56	1.52

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	905	NAG	C2-N2-C7	9.12	135.89	122.90
3	A	905	NAG	C1-O5-C5	6.23	120.63	112.19
3	A	904	NAG	C1-O5-C5	4.50	118.29	112.19
3	A	904	NAG	C2-N2-C7	4.42	129.20	122.90
3	A	904	NAG	C1-C2-N2	3.28	116.10	110.49
3	A	905	NAG	C3-C4-C5	2.71	115.07	110.24
3	A	904	NAG	C4-C3-C2	-2.16	107.85	111.02
3	A	905	NAG	C1-C2-N2	2.15	114.16	110.49
3	A	902	NAG	C1-O5-C5	-2.07	109.39	112.19
3	A	905	NAG	C8-C7-N2	2.04	119.55	116.10

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	902	NAG	C1-C2-N2-C7

*Continued on next page...*

*Continued from previous page...*



Mol	Chain	Res	Type	Atoms
3	A	905	NAG	C3-C2-N2-C7
3	A	905	NAG	O5-C5-C6-O6
3	A	903	NAG	C1-C2-N2-C7
3	A	905	NAG	C4-C5-C6-O6
3	A	904	NAG	C8-C7-N2-C2
3	A	904	NAG	O7-C7-N2-C2
3	A	905	NAG	C8-C7-N2-C2
3	A	905	NAG	O7-C7-N2-C2
3	A	902	NAG	C3-C2-N2-C7
3	A	903	NAG	C3-C2-N2-C7
3	A	904	NAG	C3-C2-N2-C7

There are no ring outliers.



4 monomers are involved in 6 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	903	NAG	2	0
3	A	904	NAG	1	0
3	A	902	NAG	1	0
3	A	905	NAG	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	646/712 (90%)	-0.51	8 (1%) 76 81	11, 21, 41, 89	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	65	THR	4.4
1	A	79	SER	3.7
1	A	696	SER	3.4
1	A	685	SER	3.3
1	A	66	SER	3.0
1	A	78	THR	2.9
1	A	695	CYS	2.6
1	A	690	GLN	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	NAG	B	2	14/15	0.51	0.24	72,84,101,122	0
2	NAG	C	2	14/15	0.57	0.23	61,77,94,97	0
2	NAG	E	2	14/15	0.71	0.14	40,61,73,75	0
2	NAG	F	2	14/15	0.72	0.16	50,65,78,85	0

*Continued on next page...*

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	NAG	B	1	14/15	0.75	0.17	32,50,65,91	0
2	NAG	D	2	14/15	0.80	0.14	37,48,70,76	0
2	NAG	F	1	14/15	0.81	0.15	21,35,58,61	0
2	NAG	G	2	14/15	0.90	0.12	20,35,50,61	0
2	NAG	E	1	14/15	0.93	0.10	23,40,47,55	0
2	NAG	C	1	14/15	0.94	0.08	25,32,41,49	0
2	NAG	D	1	14/15	0.94	0.08	20,28,38,41	0
2	NAG	G	1	14/15	0.97	0.05	12,15,21,22	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





**Electron density around Chain C:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around Chain D:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around Chain E:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around Chain F:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	NAG	A	905	14/15	0.47	0.20	73,81,99,100	0
3	NAG	A	903	14/15	0.58	0.21	65,81,93,97	0
3	NAG	A	904	14/15	0.61	0.17	44,64,75,76	0
3	NAG	A	902	14/15	0.75	0.20	42,55,67,76	0
3	NAG	A	901	14/15	0.78	0.14	25,47,65,65	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around NAG A 905:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around NAG A 903:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







**Electron density around NAG A 904:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around NAG A 902:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.