



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 8, 2020 – 08:44 PM BST

PDB ID : 3WF3
Title : Crystal structure of human beta-galactosidase mutant I51T in complex with Galactose
Authors : Suzuki, H.; Ohto, U.; Shimizu, T.
Deposited on : 2013-07-16
Resolution : 2.15 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1479 (2.16-2.16)
Clashscore	141614	1585 (2.16-2.16)
Ramachandran outliers	138981	1560 (2.16-2.16)
Sidechain outliers	138945	1559 (2.16-2.16)
RSRZ outliers	127900	1456 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	678	 2% 79% 10% 11%
1	B	678	 0% 80% 9% 11%
1	C	678	 2% 79% 9% 11%
1	D	678	 3% 79% 10% 11%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	C	701	-	-	-	X
2	NAG	D	701	-	-	-	X
2	NAG	D	703	-	-	X	-
5	SO4	D	708	-	-	X	-

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 21055 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-galactosidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	605	4841	3140	801	882	18	0	6	0
1	B	605	4825	3131	796	881	17	0	4	0
1	C	603	4808	3120	789	882	17	0	6	0
1	D	603	4805	3115	790	883	17	0	5	0

There are 100 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLU	-	expression tag	UNP P16278
A	1	ALA	-	expression tag	UNP P16278
A	2	GLU	-	expression tag	UNP P16278
A	3	ALA	-	expression tag	UNP P16278
A	4	TYR	-	expression tag	UNP P16278
A	5	VAL	-	expression tag	UNP P16278
A	6	GLU	-	expression tag	UNP P16278
A	7	PHE	-	expression tag	UNP P16278
A	8	HIS	-	expression tag	UNP P16278
A	9	HIS	-	expression tag	UNP P16278
A	10	HIS	-	expression tag	UNP P16278
A	11	HIS	-	expression tag	UNP P16278
A	12	HIS	-	expression tag	UNP P16278
A	13	HIS	-	expression tag	UNP P16278
A	14	ASP	-	expression tag	UNP P16278
A	15	TYR	-	expression tag	UNP P16278
A	16	LYS	-	expression tag	UNP P16278
A	17	ASP	-	expression tag	UNP P16278
A	18	ASP	-	expression tag	UNP P16278
A	19	ASP	-	expression tag	UNP P16278
A	20	ASP	-	expression tag	UNP P16278

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	21	LYS	-	expression tag	UNP P16278
A	22	THR	-	expression tag	UNP P16278
A	23	SER	-	expression tag	UNP P16278
A	51	THR	ILE	engineered mutation	UNP P16278
B	0	GLU	-	expression tag	UNP P16278
B	1	ALA	-	expression tag	UNP P16278
B	2	GLU	-	expression tag	UNP P16278
B	3	ALA	-	expression tag	UNP P16278
B	4	TYR	-	expression tag	UNP P16278
B	5	VAL	-	expression tag	UNP P16278
B	6	GLU	-	expression tag	UNP P16278
B	7	PHE	-	expression tag	UNP P16278
B	8	HIS	-	expression tag	UNP P16278
B	9	HIS	-	expression tag	UNP P16278
B	10	HIS	-	expression tag	UNP P16278
B	11	HIS	-	expression tag	UNP P16278
B	12	HIS	-	expression tag	UNP P16278
B	13	HIS	-	expression tag	UNP P16278
B	14	ASP	-	expression tag	UNP P16278
B	15	TYR	-	expression tag	UNP P16278
B	16	LYS	-	expression tag	UNP P16278
B	17	ASP	-	expression tag	UNP P16278
B	18	ASP	-	expression tag	UNP P16278
B	19	ASP	-	expression tag	UNP P16278
B	20	ASP	-	expression tag	UNP P16278
B	21	LYS	-	expression tag	UNP P16278
B	22	THR	-	expression tag	UNP P16278
B	23	SER	-	expression tag	UNP P16278
B	51	THR	ILE	engineered mutation	UNP P16278
C	0	GLU	-	expression tag	UNP P16278
C	1	ALA	-	expression tag	UNP P16278
C	2	GLU	-	expression tag	UNP P16278
C	3	ALA	-	expression tag	UNP P16278
C	4	TYR	-	expression tag	UNP P16278
C	5	VAL	-	expression tag	UNP P16278
C	6	GLU	-	expression tag	UNP P16278
C	7	PHE	-	expression tag	UNP P16278
C	8	HIS	-	expression tag	UNP P16278
C	9	HIS	-	expression tag	UNP P16278
C	10	HIS	-	expression tag	UNP P16278
C	11	HIS	-	expression tag	UNP P16278
C	12	HIS	-	expression tag	UNP P16278

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	13	HIS	-	expression tag	UNP P16278
C	14	ASP	-	expression tag	UNP P16278
C	15	TYR	-	expression tag	UNP P16278
C	16	LYS	-	expression tag	UNP P16278
C	17	ASP	-	expression tag	UNP P16278
C	18	ASP	-	expression tag	UNP P16278
C	19	ASP	-	expression tag	UNP P16278
C	20	ASP	-	expression tag	UNP P16278
C	21	LYS	-	expression tag	UNP P16278
C	22	THR	-	expression tag	UNP P16278
C	23	SER	-	expression tag	UNP P16278
C	51	THR	ILE	engineered mutation	UNP P16278
D	0	GLU	-	expression tag	UNP P16278
D	1	ALA	-	expression tag	UNP P16278
D	2	GLU	-	expression tag	UNP P16278
D	3	ALA	-	expression tag	UNP P16278
D	4	TYR	-	expression tag	UNP P16278
D	5	VAL	-	expression tag	UNP P16278
D	6	GLU	-	expression tag	UNP P16278
D	7	PHE	-	expression tag	UNP P16278
D	8	HIS	-	expression tag	UNP P16278
D	9	HIS	-	expression tag	UNP P16278
D	10	HIS	-	expression tag	UNP P16278
D	11	HIS	-	expression tag	UNP P16278
D	12	HIS	-	expression tag	UNP P16278
D	13	HIS	-	expression tag	UNP P16278
D	14	ASP	-	expression tag	UNP P16278
D	15	TYR	-	expression tag	UNP P16278
D	16	LYS	-	expression tag	UNP P16278
D	17	ASP	-	expression tag	UNP P16278
D	18	ASP	-	expression tag	UNP P16278
D	19	ASP	-	expression tag	UNP P16278
D	20	ASP	-	expression tag	UNP P16278
D	21	LYS	-	expression tag	UNP P16278
D	22	THR	-	expression tag	UNP P16278
D	23	SER	-	expression tag	UNP P16278
D	51	THR	ILE	engineered mutation	UNP P16278

- Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucofuranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	14	8	1	5	0	0
2	A	1	14	8	1	5	0	0
2	A	1	14	8	1	5	0	0
2	A	1	14	8	1	5	0	0
2	B	1	14	8	1	5	0	0
2	B	1	14	8	1	5	0	0
2	B	1	14	8	1	5	0	0
2	B	1	14	8	1	5	0	0
2	C	1	14	8	1	5	0	0
2	C	1	14	8	1	5	0	0
2	C	1	14	8	1	5	0	0
2	C	1	14	8	1	5	0	0
2	D	1	14	8	1	5	0	0
2	D	1	14	8	1	5	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	D	1	Total	C	N	O	0	0
			14	8	1	5		
2	D	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is beta-D-galactopyranose (three-letter code: GAL) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			12	6	6		
3	B	1	Total	C	O	0	0
			12	6	6		
3	C	1	Total	C	O	0	0
			12	6	6		
3	D	1	Total	C	O	0	0
			12	6	6		

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Cl	0	0
			1	1		
4	A	1	Total	Cl	0	0
			1	1		
4	D	1	Total	Cl	0	0
			1	1		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	C	1	Total Cl 1 1	0	0

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total O S 5 4 1	0	0
5	A	1	Total O S 5 4 1	0	0
5	B	1	Total O S 5 4 1	0	0
5	B	1	Total O S 5 4 1	0	0
5	C	1	Total O S 5 4 1	0	0
5	C	1	Total O S 5 4 1	0	0
5	D	1	Total O S 5 4 1	0	0
5	D	1	Total O S 5 4 1	0	0


- Molecule 6 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	B	1	Total C O 4 2 2	0	0
6	B	1	Total C O 4 2 2	0	0
6	C	1	Total C O 4 2 2	0	0
6	C	1	Total C O 4 2 2	0	0
6	D	1	Total C O 4 2 2	0	0
6	D	1	Total C O 4 2 2	0	0

- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	365	Total O 365 365	0	0
7	B	369	Total O 369 369	0	0
7	C	349	Total O 349 349	0	0
7	D	345	Total O 345 345	0	0



• Molecule 1: Beta-galactosidase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	94.96Å 115.89Å 140.28Å 90.00° 92.23° 90.00°	Depositor
Resolution (Å)	27.30 – 2.15 27.30 – 2.15	Depositor EDS
% Data completeness (in resolution range)	96.4 (27.30-2.15) 96.4 (27.30-2.15)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.73 (at 2.15Å)	Xtrriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.196 , 0.242 0.205 , 0.206	Depositor DCC
R_{free} test set	7959 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	27.0	Xtrriage
Anisotropy	0.088	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 44.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.049 for h,-k,-l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	21055	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.17% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, GAL, EDO, NAG, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.74	0/5020	0.82	6/6846 (0.1%)
1	B	0.77	0/4998	0.83	5/6818 (0.1%)
1	C	0.73	0/4990	0.81	1/6808 (0.0%)
1	D	0.72	0/4979	0.81	3/6795 (0.0%)
All	All	0.74	0/19987	0.82	15/27267 (0.1%)

There are no bond length outliers.

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	206	ARG	NE-CZ-NH2	-6.39	117.10	120.30
1	A	101	ASP	CB-CG-OD1	6.05	123.75	118.30
1	B	35	ASP	CB-CG-OD2	-5.71	113.16	118.30
1	B	256	ARG	NE-CZ-NH2	5.66	123.13	120.30
1	A	609[A]	MET	CA-CB-CG	5.60	122.82	113.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4841	0	4714	44	0
1	B	4825	0	4686	39	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	4808	0	4667	44	0
1	D	4805	0	4669	56	0
2	A	56	0	52	0	0
2	B	56	0	52	0	0
2	C	56	0	52	2	0
2	D	56	0	52	7	0
3	A	12	0	12	0	0
3	B	12	0	12	1	0
3	C	12	0	12	1	0
3	D	12	0	12	2	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0
5	A	10	0	0	0	0
5	B	10	0	0	0	0
5	C	10	0	0	0	0
5	D	10	0	0	2	0
6	A	8	0	12	0	0
6	B	8	0	12	0	0
6	C	8	0	12	1	0
6	D	8	0	12	1	0
7	A	365	0	0	8	0
7	B	369	0	0	9	0
7	C	349	0	0	5	0
7	D	345	0	0	8	0
All	All	21055	0	19040	179	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 179 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:555:ASN:HB2	2:D:703:NAG:O5	1.43	1.12
1:D:555:ASN:CB	2:D:703:NAG:O5	2.09	1.00
2:D:703:NAG:H3	7:D:1117:HOH:O	1.67	0.91
1:A:276:HIS:HE1	1:A:321:ASN:HD22	1.14	0.91
1:B:437:ASN:HD21	1:B:458:ASN:H	1.17	0.88

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	607/678 (90%)	576 (95%)	28 (5%)	3 (0%)	29	22
1	B	605/678 (89%)	579 (96%)	26 (4%)	0	100	100
1	C	605/678 (89%)	578 (96%)	27 (4%)	0	100	100
1	D	604/678 (89%)	576 (95%)	27 (4%)	1 (0%)	47	46
All	All	2421/2712 (89%)	2309 (95%)	108 (4%)	4 (0%)	47	46

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	628	SER
1	A	611	SER
1	A	493	LYS
1	D	423	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	524/585 (90%)	510 (97%)	14 (3%)	44	46
1	B	522/585 (89%)	511 (98%)	11 (2%)	53	57
1	C	522/585 (89%)	509 (98%)	13 (2%)	47	49
1	D	521/585 (89%)	514 (99%)	7 (1%)	69	74
All	All	2089/2340 (89%)	2044 (98%)	45 (2%)	53	55

5 of 45 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	421	THR
1	C	35	ASP
1	D	409	VAL
1	B	611	SER
1	C	49	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 32 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	318	ASN
1	B	424	GLN
1	D	428	ASN
1	B	355	GLN
1	B	437	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 40 ligands modelled in this entry, 4 are monoatomic - leaving 36 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	SO4	D	708	-	4,4,4	0.38	0	6,6,6	0.72	0
6	EDO	A	710	-	3,3,3	0.57	0	2,2,2	0.62	0
6	EDO	D	709	-	3,3,3	0.21	0	2,2,2	0.54	0
5	SO4	D	707	-	4,4,4	0.39	0	6,6,6	0.34	0
6	EDO	B	708	-	3,3,3	0.56	0	2,2,2	0.09	0
2	NAG	D	703	1	14,14,15	0.30	0	17,19,21	0.61	0
3	GAL	D	706	-	12,12,12	0.70	0	17,17,17	0.98	1 (5%)
2	NAG	A	701	1	14,14,15	0.82	0	17,19,21	1.07	1 (5%)
2	NAG	D	701	1	14,14,15	0.84	1 (7%)	17,19,21	1.37	3 (17%)
6	EDO	B	709	-	3,3,3	0.42	0	2,2,2	0.56	0
2	NAG	D	704	1	14,14,15	0.60	0	17,19,21	1.24	2 (11%)
3	GAL	C	706	-	12,12,12	0.85	0	17,17,17	1.02	1 (5%)
3	GAL	B	710	-	12,12,12	0.89	0	17,17,17	1.25	2 (11%)
5	SO4	A	708	-	4,4,4	0.43	0	6,6,6	0.75	0
2	NAG	A	703	1	14,14,15	0.74	0	17,19,21	2.60	6 (35%)
6	EDO	C	710	-	3,3,3	0.50	0	2,2,2	0.09	0
6	EDO	A	709	-	3,3,3	0.45	0	2,2,2	0.42	0
5	SO4	A	707	-	4,4,4	0.41	0	6,6,6	0.41	0
2	NAG	B	701	1	14,14,15	1.04	0	17,19,21	1.81	5 (29%)
2	NAG	C	701	1	14,14,15	0.60	0	17,19,21	1.43	2 (11%)
6	EDO	D	710	-	3,3,3	0.62	0	2,2,2	0.57	0
2	NAG	B	704	1	14,14,15	0.58	0	17,19,21	0.78	0
5	SO4	C	707	-	4,4,4	0.51	0	6,6,6	0.39	0
5	SO4	C	708	-	4,4,4	0.45	0	6,6,6	0.54	0
5	SO4	B	706	-	4,4,4	0.41	0	6,6,6	0.20	0
2	NAG	B	703	1	14,14,15	0.63	0	17,19,21	1.65	2 (11%)
2	NAG	C	704	1	14,14,15	0.67	0	17,19,21	1.40	2 (11%)
2	NAG	B	702	1	14,14,15	0.86	0	17,19,21	2.69	6 (35%)
2	NAG	A	704	1	14,14,15	0.57	0	17,19,21	1.54	3 (17%)
2	NAG	D	702	1	14,14,15	0.62	0	17,19,21	3.29	6 (35%)
5	SO4	B	707	-	4,4,4	0.34	0	6,6,6	0.82	0
6	EDO	C	709	-	3,3,3	0.23	0	2,2,2	0.79	0
3	GAL	A	705	-	12,12,12	0.74	0	17,17,17	1.12	2 (11%)
2	NAG	A	702	1	14,14,15	0.69	0	17,19,21	2.91	6 (35%)
2	NAG	C	702	1	14,14,15	0.61	0	17,19,21	1.35	2 (11%)
2	NAG	C	703	1	14,14,15	0.65	0	17,19,21	1.71	5 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	EDO	A	710	-	-	1/1/1/1	-
6	EDO	D	709	-	-	1/1/1/1	-
6	EDO	B	708	-	-	1/1/1/1	-
2	NAG	D	703	1	-	2/6/23/26	0/1/1/1
3	GAL	D	706	-	-	1/2/22/22	0/1/1/1
2	NAG	A	701	1	-	1/6/23/26	0/1/1/1
2	NAG	D	701	1	-	2/6/23/26	0/1/1/1
6	EDO	B	709	-	-	0/1/1/1	-
2	NAG	D	704	1	-	0/6/23/26	0/1/1/1
3	GAL	C	706	-	-	1/2/22/22	0/1/1/1
3	GAL	B	710	-	-	1/2/22/22	0/1/1/1
2	NAG	A	703	1	-	0/6/23/26	0/1/1/1
6	EDO	C	710	-	-	1/1/1/1	-
6	EDO	A	709	-	-	0/1/1/1	-
2	NAG	B	701	1	-	2/6/23/26	0/1/1/1
2	NAG	C	701	1	-	2/6/23/26	0/1/1/1
6	EDO	D	710	-	-	1/1/1/1	-
2	NAG	B	704	1	-	0/6/23/26	0/1/1/1
2	NAG	B	703	1	-	0/6/23/26	0/1/1/1
2	NAG	C	704	1	-	0/6/23/26	0/1/1/1
2	NAG	B	702	1	-	2/6/23/26	0/1/1/1
2	NAG	A	704	1	-	0/6/23/26	0/1/1/1
2	NAG	D	702	1	-	0/6/23/26	0/1/1/1
6	EDO	C	709	-	-	1/1/1/1	-
3	GAL	A	705	-	-	1/2/22/22	0/1/1/1
2	NAG	A	702	1	-	2/6/23/26	0/1/1/1
2	NAG	C	702	1	-	0/6/23/26	0/1/1/1
2	NAG	C	703	1	-	0/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	701	NAG	C1-C2	2.45	1.56	1.52

The worst 5 of 57 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	702	NAG	C1-O5-C5	10.38	126.26	112.19
2	A	702	NAG	C1-O5-C5	8.20	123.30	112.19

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	702	NAG	C1-O5-C5	6.99	121.67	112.19
2	A	703	NAG	C1-O5-C5	6.78	121.38	112.19
2	B	703	NAG	C1-O5-C5	4.92	118.86	112.19

There are no chirality outliers.

5 of 23 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	701	NAG	C4-C5-C6-O6
2	B	702	NAG	O5-C5-C6-O6
2	B	702	NAG	C4-C5-C6-O6
2	D	703	NAG	O5-C5-C6-O6
2	D	701	NAG	O5-C5-C6-O6

There are no ring outliers.

9 monomers are involved in 17 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	D	708	SO4	2	0
6	D	709	EDO	1	0
2	D	703	NAG	7	0
3	D	706	GAL	2	0
3	C	706	GAL	1	0
3	B	710	GAL	1	0
6	C	709	EDO	1	0
2	C	702	NAG	1	0
2	C	703	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	605/678 (89%)	0.00	15 (2%) 57 65	16, 26, 41, 85	0
1	B	605/678 (89%)	-0.04	9 (1%) 73 79	16, 25, 40, 83	0
1	C	603/678 (88%)	0.05	15 (2%) 57 65	17, 26, 49, 81	0
1	D	603/678 (88%)	0.10	22 (3%) 42 51	16, 28, 50, 83	0
All	All	2416/2712 (89%)	0.03	61 (2%) 57 65	16, 26, 46, 85	0

The worst 5 of 61 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	647	SER	8.4
1	C	527	TRP	6.8
1	D	527	TRP	6.5
1	A	647	SER	6.0
1	A	629	ASP	5.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NAG	D	703	14/15	0.63	0.31	72,93,100,101	0
2	NAG	D	701	14/15	0.69	0.42	81,88,93,93	0
2	NAG	C	701	14/15	0.73	0.43	79,88,98,103	0
2	NAG	D	702	14/15	0.74	0.20	35,46,49,51	0
6	EDO	B	708	4/4	0.75	0.15	45,47,47,49	0
6	EDO	A	709	4/4	0.81	0.20	45,49,52,53	0
2	NAG	B	703	14/15	0.83	0.23	45,53,61,68	0
6	EDO	D	710	4/4	0.84	0.18	28,30,33,33	0
2	NAG	A	704	14/15	0.85	0.39	54,60,64,65	0
2	NAG	A	703	14/15	0.85	0.14	40,45,50,61	0
2	NAG	C	702	14/15	0.85	0.24	43,51,56,56	0
6	EDO	A	710	4/4	0.86	0.22	25,29,30,35	0
5	SO4	D	707	5/5	0.88	0.26	64,69,73,80	0
6	EDO	D	709	4/4	0.88	0.14	35,38,39,45	0
2	NAG	B	701	14/15	0.88	0.14	28,31,34,43	0
2	NAG	C	703	14/15	0.88	0.21	36,41,44,44	0
2	NAG	A	702	14/15	0.89	0.12	28,33,36,42	0
5	SO4	C	707	5/5	0.89	0.24	55,55,59,63	0
2	NAG	B	704	14/15	0.89	0.24	46,51,59,63	0
2	NAG	B	702	14/15	0.91	0.12	25,30,34,35	0
6	EDO	C	710	4/4	0.91	0.14	36,41,41,45	0
2	NAG	D	704	14/15	0.92	0.12	27,30,32,36	0
5	SO4	A	707	5/5	0.92	0.24	69,72,73,78	0
6	EDO	C	709	4/4	0.93	0.18	41,44,46,54	0
2	NAG	C	704	14/15	0.94	0.13	24,26,29,35	0
3	GAL	B	710	12/12	0.94	0.11	19,21,23,30	0
5	SO4	A	708	5/5	0.94	0.17	45,48,54,56	0
5	SO4	D	708	5/5	0.94	0.13	49,50,52,56	0
5	SO4	B	706	5/5	0.95	0.28	56,63,64,67	0
3	GAL	A	705	12/12	0.95	0.11	16,20,24,29	0
2	NAG	A	701	14/15	0.95	0.08	24,30,33,41	0
3	GAL	D	706	12/12	0.95	0.09	21,25,27,31	0
5	SO4	B	707	5/5	0.95	0.15	47,47,54,55	0
6	EDO	B	709	4/4	0.96	0.10	33,33,33,37	0
3	GAL	C	706	12/12	0.97	0.11	18,22,25,32	0
5	SO4	C	708	5/5	0.97	0.15	40,43,47,48	0
4	CL	D	705	1/1	0.99	0.09	20,20,20,20	0
4	CL	A	706	1/1	0.99	0.09	21,21,21,21	0
4	CL	C	705	1/1	0.99	0.09	21,21,21,21	0
4	CL	B	705	1/1	0.99	0.06	20,20,20,20	0

6.5 Other polymers

There are no such residues in this entry.