



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 17, 2024 – 02:08 PM JST

PDB ID : 8WF6  
Title : 14-3-3 zeta complexed with S559 phosphorylated peptide derived from GPIb alpha cytoplasmic domain  
Authors : Shu, Z.M.; Zhou, A.W.  
Deposited on : 2023-09-19  
Resolution : 2.47 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.002 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.38.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.47 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	7106 (2.50-2.46)
Clashscore	180529	7991 (2.50-2.46)
Ramachandran outliers	177936	7888 (2.50-2.46)
Sidechain outliers	177891	7890 (2.50-2.46)
RSRZ outliers	164620	7106 (2.50-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	246	
1	B	246	
1	C	246	
1	D	246	
2	E	12	
2	F	12	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	Q	12	 <p>42% 8% 50% 17%</p>
2	R	12	 <p>17% 50% 8% 42%</p>

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 7160 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 14-3-3 protein zeta/delta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	221	Total 1700	C 1071	N 287	O 333	S 9	0	0	0
1	B	228	Total 1752	C 1098	N 294	O 350	S 10	0	0	0
1	C	226	Total 1741	C 1093	N 297	O 341	S 10	0	0	0
1	D	224	Total 1762	C 1104	N 299	O 350	S 9	0	0	0


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	expression tag	UNP P63104
B	0	SER	-	expression tag	UNP P63104
C	0	SER	-	expression tag	UNP P63104
D	0	SER	-	expression tag	UNP P63104

- Molecule 2 is a protein called S559 phosphorylated peptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	Q	6	Total 47	C 29	N 6	O 11	P 1	0	0	0
2	R	7	Total 52	C 32	N 7	O 12	P 1	0	0	0
2	E	5	Total 36	C 20	N 5	O 10	P 1	0	0	0
2	F	5	Total 36	C 20	N 5	O 10	P 1	0	0	0

- Molecule 3 is SUCCINIC ACID (three-letter code: SIN) (formula: C<sub>4</sub>H<sub>6</sub>O<sub>4</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	C	1	Total C O 8 4 4	0	0
3	D	1	Total C O 8 4 4	0	0



- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	2	Total O 2 2	0	0
4	B	2	Total O 2 2	0	0
4	Q	1	Total O 1 1	0	0
4	R	1	Total O 1 1	0	0
4	C	3	Total O 3 3	0	0
4	D	7	Total O 7 7	0	0
4	E	1	Total O 1 1	0	0
4	F	1	Total O 1 1	0	0



### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 14-3-3 protein zeta/delta





- Molecule 1: 14-3-3 protein zeta/delta



- Molecule 1: 14-3-3 protein zeta/delta




- Molecule 1: 14-3-3 protein zeta/delta





## • Molecule 2: S559 phosphorylated peptide

Chain Q:  42% 8% 50%



## • Molecule 2: S559 phosphorylated peptide

Chain R:  17% 50% 8% 42%

## • Molecule 2: S559 phosphorylated peptide

Chain E:  8% 33% 8% 58%

## • Molecule 2: S559 phosphorylated peptide

Chain F:  8% 33% 8% 58%

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	71.31Å 112.24Å 83.92Å 90.00° 97.52° 90.00°	Depositor
Resolution (Å)	31.40 – 2.47 31.40 – 2.47	Depositor EDS
% Data completeness (in resolution range)	99.7 (31.40-2.47) 99.8 (31.40-2.47)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.87 (at 2.48Å)	Xtrriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, $R_{free}$	0.197 , 0.238 0.195 , 0.237	Depositor DCC
$R_{free}$ test set	2365 reflections (4.86%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	50.0	Xtrriage
Anisotropy	0.478	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 49.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	7160	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	66.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 16.96% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SEP, SIN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.24	0/1722	0.42	0/2326
1	B	0.23	0/1775	0.42	0/2401
1	C	0.23	0/1763	0.41	0/2379
1	D	0.23	0/1784	0.41	0/2402
2	E	0.18	0/25	0.42	0/32
2	F	0.25	0/25	0.51	0/32
2	Q	0.24	0/37	0.40	0/48
2	R	0.24	0/42	0.43	0/55
All	All	0.23	0/7173	0.42	0/9675

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1700	0	1633	23	0
1	B	1752	0	1660	17	0
1	C	1741	0	1681	6	0
1	D	1762	0	1722	9	0
2	E	36	0	31	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	F	36	0	31	1	0
2	Q	47	0	40	0	0
2	R	52	0	42	0	0
3	C	8	0	4	0	0
3	D	8	0	4	0	0
4	A	2	0	0	1	0
4	B	2	0	0	0	0
4	C	3	0	0	0	0
4	D	7	0	0	0	0
4	E	1	0	0	0	0
4	F	1	0	0	0	0
4	Q	1	0	0	0	0
4	R	1	0	0	0	0
All	All	7160	0	6848	54	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:49:LYS:NZ	4:A:301:HOH:O	2.27	0.68
1:B:101:LEU:HA	1:B:105:LEU:HB2	1.78	0.65
1:A:22:MET:HG2	1:A:47:ALA:HB2	1.81	0.62
1:A:115:LYS:HD2	1:A:119:LEU:HD21	1.80	0.62
1:B:161:GLN:NE2	1:B:202:GLU:OE2	2.35	0.59
1:D:83:ARG:NH1	1:D:87:GLU:OE1	2.39	0.55
1:C:49:LYS:NZ	2:F:7:SEP:O3P	2.37	0.54
1:A:59:TRP:CE2	1:A:132:VAL:HG12	2.43	0.54
1:B:80:ARG:O	1:B:84:GLU:HG3	2.09	0.53
1:B:153:PHE:HA	1:B:170:LEU:HD21	1.92	0.52
1:A:36:LEU:HD22	1:A:105:LEU:HD21	1.92	0.52
1:B:38:ASN:O	1:B:42:ASN:ND2	2.42	0.52
1:A:80:ARG:O	1:A:84:GLU:HG3	2.10	0.52
1:C:22:MET:HG2	1:C:47:ALA:HB2	1.93	0.50
1:D:101:LEU:HA	1:D:105:LEU:HB2	1.95	0.49
1:D:77:GLN:HE22	1:D:80:ARG:HH11	1.59	0.48
1:A:51:VAL:HG12	1:A:93:ILE:HD13	1.94	0.48
1:A:119:LEU:HB2	1:A:152:ALA:HB2	1.94	0.48
1:C:58:SER:HB3	1:D:16:ALA:HB1	1.94	0.48
1:D:187:LYS:HB2	1:D:187:LYS:HE3	1.71	0.47

Continued on next page...

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:59:TRP:NE1	1:A:87:GLU:OE2	2.40	0.47
1:A:190:SER:O	1:A:194:THR:HG23	2.15	0.47
1:D:77:GLN:HE21	1:D:77:GLN:HA	1.81	0.46
1:A:63:SER:O	1:A:67:GLN:HG2	2.15	0.46
1:A:210:SER:O	1:A:210:SER:OG	2.26	0.45
1:D:37:SER:OG	1:D:40:GLU:HG3	2.16	0.45
1:B:119:LEU:HB3	1:B:152:ALA:HB2	1.98	0.45
1:B:177:PHE:CE2	1:B:182:LEU:HG	2.51	0.45
1:B:119:LEU:HD13	1:B:151:GLU:HG2	2.00	0.44
1:B:203:LEU:HD23	1:B:203:LEU:HA	1.83	0.44
1:B:193:LYS:HE3	1:B:197:ASP:OD2	2.18	0.43
1:A:47:ALA:O	1:A:51:VAL:HG23	2.18	0.43
1:A:193:LYS:NZ	1:A:197:ASP:OD2	2.49	0.43
1:B:69:THR:O	1:B:76:GLN:NE2	2.43	0.43
1:D:3:LYS:HE2	1:D:40:GLU:OE1	2.19	0.43
1:A:11:LYS:HE2	1:A:11:LYS:HB3	1.77	0.43
1:B:58:SER:O	1:B:62:VAL:HG23	2.18	0.43
1:A:105:LEU:HD22	1:A:118:TYR:CZ	2.54	0.42
1:B:126:TYR:HB3	1:B:145:SER:HB2	2.01	0.42
1:A:12:LEU:HD23	1:A:12:LEU:HA	1.88	0.42
1:A:161:GLN:O	1:A:167:ARG:HD3	2.19	0.42
1:B:87:GLU:OE2	1:B:91:ARG:NH2	2.53	0.42
1:A:117:PHE:HE2	1:A:121:MET:CE	2.33	0.42
1:B:155:ILE:HD12	1:B:155:ILE:H	1.85	0.42
1:A:2:ASP:OD2	1:A:3:LYS:N	2.52	0.42
1:B:41:ARG:HD2	1:B:117:PHE:CD1	2.55	0.42
1:A:164:HIS:CD2	1:A:166:ILE:HD12	2.55	0.41
1:C:13:ALA:HB3	1:C:22:MET:HB2	2.02	0.41
1:A:182:LEU:HD23	1:A:182:LEU:HA	1.90	0.41
1:C:182:LEU:HD12	1:C:182:LEU:HA	1.80	0.40
1:C:126:TYR:HB3	1:C:145:SER:HB2	2.03	0.40
1:D:154:GLU:HA	1:D:157:LYS:HD3	2.03	0.40
1:A:112:ALA:O	1:A:116:VAL:HG23	2.21	0.40
1:B:141:ILE:H	1:B:141:ILE:HG13	1.74	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	215/246 (87%)	209 (97%)	6 (3%)	0	100	100
1	B	222/246 (90%)	217 (98%)	5 (2%)	0	100	100
1	C	220/246 (89%)	214 (97%)	6 (3%)	0	100	100
1	D	218/246 (89%)	213 (98%)	5 (2%)	0	100	100
2	E	2/12 (17%)	1 (50%)	1 (50%)	0	100	100
2	F	2/12 (17%)	2 (100%)	0	0	100	100
2	Q	3/12 (25%)	3 (100%)	0	0	100	100
2	R	4/12 (33%)	4 (100%)	0	0	100	100
All	All	886/1032 (86%)	863 (97%)	23 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	172/210 (82%)	166 (96%)	6 (4%)	31	54
1	B	179/210 (85%)	174 (97%)	5 (3%)	38	63
1	C	178/210 (85%)	175 (98%)	3 (2%)	56	77
1	D	186/210 (89%)	182 (98%)	4 (2%)	47	70
2	E	3/10 (30%)	3 (100%)	0	100	100
2	F	3/10 (30%)	3 (100%)	0	100	100

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	Q	4/10 (40%)	4 (100%)	0	100	100
2	R	4/10 (40%)	4 (100%)	0	100	100
All	All	729/880 (83%)	711 (98%)	18 (2%)	42	67

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	26	MET
1	A	90	LEU
1	A	102	GLU
1	A	121	MET
1	A	174	PHE
1	A	223	ASP
1	B	1	MET
1	B	2	ASP
1	B	77	GLN
1	B	92	ASP
1	B	174	PHE
1	C	121	MET
1	C	144	GLN
1	C	174	PHE
1	D	80	ARG
1	D	96	ASP
1	D	102	GLU
1	D	174	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	146	GLN
1	A	164	HIS
1	B	42	ASN
1	B	76	GLN
1	B	108	ASN
1	C	161	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	SEP	F	7	2	8,9,10	1.55	1 (12%)	8,12,14	1.56	2 (25%)
2	SEP	Q	7	2	8,9,10	1.55	1 (12%)	8,12,14	1.62	2 (25%)
2	SEP	E	7	2	8,9,10	1.56	1 (12%)	8,12,14	1.45	2 (25%)
2	SEP	R	7	2	8,9,10	1.55	1 (12%)	8,12,14	1.69	2 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SEP	F	7	2	-	0/5/8/10	-
2	SEP	Q	7	2	-	0/5/8/10	-
2	SEP	E	7	2	-	0/5/8/10	-
2	SEP	R	7	2	-	0/5/8/10	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	7	SEP	P-O1P	3.40	1.61	1.50
2	Q	7	SEP	P-O1P	3.38	1.61	1.50
2	R	7	SEP	P-O1P	3.38	1.61	1.50
2	F	7	SEP	P-O1P	3.37	1.61	1.50

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	R	7	SEP	OG-CB-CA	3.17	111.23	108.14
2	R	7	SEP	P-OG-CB	-3.01	110.00	118.30
2	Q	7	SEP	P-OG-CB	-2.96	110.15	118.30

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	Q	7	SEP	OG-CB-CA	2.95	111.02	108.14
2	F	7	SEP	P-OG-CB	-2.87	110.38	118.30
2	E	7	SEP	P-OG-CB	-2.78	110.63	118.30
2	F	7	SEP	OG-CB-CA	2.77	110.84	108.14
2	E	7	SEP	OG-CB-CA	2.36	110.45	108.14

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	F	7	SEP	1	0

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	SIN	C	301	-	7,7,7	1.01	0	8,8,8	1.73	3 (37%)
3	SIN	D	301	-	7,7,7	1.01	0	8,8,8	1.72	2 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SIN	C	301	-	-	2/5/5/5	-
3	SIN	D	301	-	-	2/5/5/5	-

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	301	SIN	C3-C2-C1	-2.27	108.71	113.60
3	C	301	SIN	C2-C3-C4	-2.25	108.77	113.60
3	C	301	SIN	C3-C2-C1	-2.04	109.22	113.60
3	C	301	SIN	O4-C4-C3	2.00	120.47	114.03
3	D	301	SIN	O4-C4-C3	2.00	120.46	114.03

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	301	SIN	O1-C1-C2-C3
3	C	301	SIN	O2-C1-C2-C3
3	D	301	SIN	O2-C1-C2-C3
3	D	301	SIN	O1-C1-C2-C3

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	221/246 (89%)	0.57	11 (4%) 35 33	52, 77, 113, 138	0
1	B	228/246 (92%)	0.32	10 (4%) 39 37	46, 67, 106, 122	0
1	C	226/246 (91%)	0.16	8 (3%) 47 45	37, 58, 92, 110	0
1	D	224/246 (91%)	-0.03	5 (2%) 62 60	34, 52, 91, 126	0
2	E	4/12 (33%)	0.78	1 (25%) 2 2	41, 47, 49, 58	0
2	F	4/12 (33%)	0.75	1 (25%) 2 2	47, 52, 59, 69	0
2	Q	5/12 (41%)	0.72	0 100 100	50, 58, 70, 88	0
2	R	6/12 (50%)	1.61	2 (33%) 1 1	65, 78, 114, 114	0
All	All	918/1032 (88%)	0.27	38 (4%) 42 40	34, 64, 105, 138	0

All (38) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	230	SER	4.9
1	D	69	THR	4.1
1	C	211	TYR	4.0
1	A	230	SER	3.4
1	C	206	LEU	3.4
1	D	203	LEU	3.4
2	R	11	PHE	3.3
1	A	1	MET	3.2
1	D	72	ALA	3.1
1	A	69	THR	3.1
2	R	5	ARG	3.0
1	C	203	LEU	3.0
1	B	134	ALA	2.9
1	A	73	GLU	2.8
1	B	136	ASP	2.8
1	C	136	ASP	2.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	D	208	GLU	2.7
1	B	210	SER	2.7
1	A	3	LYS	2.6
1	B	1	MET	2.6
1	B	207	SER	2.5
1	C	72	ALA	2.5
1	A	210	SER	2.5
1	B	232	THR	2.5
1	B	208	GLU	2.4
1	C	210	SER	2.4
1	B	69	THR	2.4
1	B	73	GLU	2.3
1	C	134	ALA	2.3
1	A	163	THR	2.3
2	E	6	GLY	2.3
2	F	10	THR	2.3
1	A	161	GLN	2.2
1	B	211	TYR	2.2
1	A	106	ILE	2.1
1	A	16	ALA	2.1
1	A	78	MET	2.1
1	D	231	ASP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	SEP	Q	7	10/11	0.96	0.08	43,52,58,59	0
2	SEP	R	7	10/11	0.97	0.08	48,60,68,76	0
2	SEP	E	7	10/11	0.98	0.06	32,38,47,57	0
2	SEP	F	7	10/11	0.98	0.06	35,46,51,55	0

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	SIN	C	301	8/8	0.83	0.18	46,57,65,69	0
3	SIN	D	301	8/8	0.88	0.13	48,55,65,73	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.