



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 19, 2024 – 04:11 PM JST

PDB ID : 8WHL  
Title : Crystal structure of CLASP2 in complex with CENP-E  
Authors : Jia, X.; Wei, Z.  
Deposited on : 2023-09-23  
Resolution : 3.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.37.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1133 (3.20-3.20)
Clashscore	141614	1253 (3.20-3.20)
Ramachandran outliers	138981	1234 (3.20-3.20)
Sidechain outliers	138945	1233 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Mol	Chain	Length	Quality of chain
1	A	235	
1	B	235	
1	C	235	
1	D	235	
2	E	66	
2	F	66	
2	G	66	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	H	66	 74% 9% 17%

## 2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 8755 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CLIP-associating protein 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	B	226	Total 1783	C 1137	N 307	O 330	S 9	0	0	0
1	D	226	Total 1780	C 1132	N 306	O 333	S 9	0	0	0
1	C	220	Total 1743	C 1112	N 299	O 323	S 9	0	0	0
1	A	228	Total 1799	C 1144	N 309	O 337	S 9	0	0	0

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	1245	GLY	-	expression tag	UNP O75122
B	1246	PRO	-	expression tag	UNP O75122
B	1247	GLY	-	expression tag	UNP O75122
B	1248	SER	-	expression tag	UNP O75122
B	1249	GLU	-	expression tag	UNP O75122
B	1250	PHE	-	expression tag	UNP O75122
D	1245	GLY	-	expression tag	UNP O75122
D	1246	PRO	-	expression tag	UNP O75122
D	1247	GLY	-	expression tag	UNP O75122
D	1248	SER	-	expression tag	UNP O75122
D	1249	GLU	-	expression tag	UNP O75122
D	1250	PHE	-	expression tag	UNP O75122
C	1245	GLY	-	expression tag	UNP O75122
C	1246	PRO	-	expression tag	UNP O75122
C	1247	GLY	-	expression tag	UNP O75122
C	1248	SER	-	expression tag	UNP O75122
C	1249	GLU	-	expression tag	UNP O75122
C	1250	PHE	-	expression tag	UNP O75122
A	1245	GLY	-	expression tag	UNP O75122
A	1246	PRO	-	expression tag	UNP O75122
A	1247	GLY	-	expression tag	UNP O75122

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
A	1248	SER	-	expression tag	UNP O75122
A	1249	GLU	-	expression tag	UNP O75122
A	1250	PHE	-	expression tag	UNP O75122

- Molecule 2 is a protein called Centromere-associated protein E.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	F	55	431	264	70	96	1	0	0	0
2	H	55	411	257	69	84	1	0	0	0
2	G	52	395	248	62	84	1	0	0	0
2	E	52	398	245	63	89	1	0	0	0

There are 16 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
F	483	GLY	-	expression tag	UNP Q02224
F	484	PRO	-	expression tag	UNP Q02224
F	485	GLY	-	expression tag	UNP Q02224
F	486	SER	-	expression tag	UNP Q02224
H	483	GLY	-	expression tag	UNP Q02224
H	484	PRO	-	expression tag	UNP Q02224
H	485	GLY	-	expression tag	UNP Q02224
H	486	SER	-	expression tag	UNP Q02224
G	483	GLY	-	expression tag	UNP Q02224
G	484	PRO	-	expression tag	UNP Q02224
G	485	GLY	-	expression tag	UNP Q02224
G	486	SER	-	expression tag	UNP Q02224
E	483	GLY	-	expression tag	UNP Q02224
E	484	PRO	-	expression tag	UNP Q02224
E	485	GLY	-	expression tag	UNP Q02224
E	486	SER	-	expression tag	UNP Q02224

- Molecule 3 is ACETATE ION (three-letter code: ACT) (formula: C<sub>2</sub>H<sub>3</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is MALONIC ACID (three-letter code: MLA) (formula: C<sub>3</sub>H<sub>4</sub>O<sub>4</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			7	3	4		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: CLIP-associating protein 2




- Molecule 1: CLIP-associating protein 2



- Molecule 1: CLIP-associating protein 2




- Molecule 1: CLIP-associating protein 2








- Molecule 2: Centromere-associated protein E





- Molecule 2: Centromere-associated protein E



- Molecule 2: Centromere-associated protein E



- Molecule 2: Centromere-associated protein E





## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	106.22Å 106.22Å 327.05Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	46.89 – 3.20 46.89 – 3.20	Depositor EDS
% Data completeness (in resolution range)	99.7 (46.89-3.20) 99.7 (46.89-3.20)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.97 (at 3.19Å)	Xtrriage
Refinement program	PHENIX (1.18.2_3874: ???)	Depositor
R, $R_{free}$	0.246 , 0.278 0.246 , 0.278	Depositor DCC
$R_{free}$ test set	1707 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	101.7	Xtrriage
Anisotropy	0.142	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 79.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.40$ , $\langle L^2 \rangle = 0.22$	Xtrriage
Estimated twinning fraction	0.419 for h,-h-k,-l	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	8755	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	120.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.50% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACT, MLA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.23	0/1825	0.39	0/2467
1	B	0.23	0/1810	0.40	0/2447
1	C	0.23	0/1767	0.39	0/2387
1	D	0.23	0/1806	0.39	0/2442
2	E	0.22	0/400	0.31	0/541
2	F	0.23	0/433	0.31	0/583
2	G	0.23	0/397	0.34	0/535
2	H	0.23	0/413	0.37	0/556
All	All	0.23	0/8851	0.38	0/11958

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1799	0	1881	16	0
1	B	1783	0	1864	18	0
1	C	1743	0	1837	17	0
1	D	1780	0	1854	18	0
2	E	398	0	343	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	F	431	0	383	3	0
2	G	395	0	360	3	0
2	H	411	0	370	4	0
3	A	4	0	3	0	0
3	B	4	0	3	0	0
4	B	7	0	2	0	0
All	All	8755	0	8900	72	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (72) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1275:ARG:NH1	1:B:1311:ASP:OD2	2.16	0.78
1:C:1359:ARG:HD2	2:G:523:GLU:OE2	1.87	0.73
1:B:1343:VAL:HG22	1:B:1368:LEU:HD21	1.79	0.64
1:C:1287:GLN:OE1	1:C:1329:ARG:NH1	2.34	0.61
1:D:1332:PRO:HB3	1:D:1371:SER:HB3	1.83	0.61
1:C:1375:GLU:HG3	1:C:1415:LEU:HD11	1.83	0.59
1:A:1375:GLU:HG3	1:A:1415:LEU:HD11	1.85	0.59
1:D:1336:LYS:O	1:D:1376:GLN:NE2	2.35	0.57
1:B:1348:GLU:OE1	1:B:1351:LYS:NZ	2.38	0.57
2:H:496:ILE:HG23	2:G:500:LEU:HD22	1.87	0.56
1:A:1340:GLU:HG3	1:A:1380:VAL:HG22	1.87	0.56
1:D:1370:THR:HA	1:D:1406:ARG:HD2	1.87	0.56
1:A:1343:VAL:HG22	1:A:1368:LEU:HD21	1.90	0.54
1:B:1265:LEU:HA	1:B:1275:ARG:HD2	1.89	0.54
1:D:1287:GLN:OE1	1:D:1329:ARG:NH1	2.42	0.52
1:D:1291:PHE:HA	1:D:1294:TRP:HD1	1.74	0.52
1:C:1370:THR:HA	1:C:1406:ARG:HD2	1.91	0.52
1:A:1373:SER:HB2	1:A:1376:GLN:HB3	1.90	0.52
1:B:1311:ASP:O	1:B:1317:ARG:NH1	2.33	0.51
1:B:1295:ASP:OD1	1:B:1334:ARG:NH2	2.40	0.51
1:D:1275:ARG:NH1	1:D:1311:ASP:OD2	2.36	0.49
1:B:1409:LYS:HG3	1:B:1449:ILE:HG22	1.94	0.49
1:D:1285:LEU:O	1:D:1290:SER:OG	2.23	0.49
2:H:511:VAL:O	2:H:515:GLU:HG2	2.14	0.48
1:A:1348:GLU:O	1:A:1351:LYS:HG2	2.14	0.48
1:B:1348:GLU:HG3	1:D:1387:THR:HB	1.96	0.48
1:C:1335:PHE:HB2	1:C:1372:ILE:HD12	1.94	0.48

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:1332:PRO:HB3	1:C:1371:SER:HB3	1.97	0.47
1:A:1428:ASP:OD1	1:A:1464:LYS:NZ	2.45	0.47
2:H:515:GLU:OE1	2:H:518:ARG:NH2	2.46	0.47
1:C:1287:GLN:HA	1:C:1330:HIS:CE1	2.50	0.47
1:C:1470:LEU:HD21	2:G:540:LEU:HD22	1.96	0.47
1:C:1264:GLU:HG3	1:C:1275:ARG:HG2	1.97	0.47
1:B:1470:LEU:HD21	2:F:540:LEU:HD22	1.98	0.46
1:B:1393:ASN:OD1	1:B:1394:LEU:N	2.49	0.45
1:C:1331:GLN:HB3	1:C:1334:ARG:HD2	1.98	0.45
1:C:1275:ARG:NH1	1:C:1311:ASP:OD2	2.37	0.45
1:A:1347:LEU:HB3	1:A:1392:ILE:HG23	1.99	0.45
1:B:1264:GLU:HG3	1:B:1275:ARG:HG2	1.98	0.45
1:D:1331:GLN:HB3	1:D:1334:ARG:HD2	1.98	0.45
1:D:1347:LEU:HD21	1:D:1399:MET:HE3	1.98	0.45
1:D:1322:LYS:HG2	1:D:1325:ARG:HH12	1.82	0.44
1:D:1416:LEU:HD11	1:D:1449:ILE:HD13	2.00	0.44
2:F:529:LYS:HE2	2:F:529:LYS:HB3	1.85	0.43
1:A:1321:LEU:HD13	1:A:1361:ALA:HB2	2.01	0.43
1:D:1393:ASN:OD1	1:D:1394:LEU:N	2.52	0.43
2:F:500:LEU:HD22	2:E:496:ILE:HG23	2.01	0.43
1:C:1351:LYS:HD3	1:C:1392:ILE:HD11	2.00	0.42
1:A:1409:LYS:HG3	1:A:1449:ILE:HG22	2.01	0.42
1:A:1416:LEU:HD23	1:A:1416:LEU:HA	1.90	0.42
1:D:1382:CYS:HB2	1:D:1383:PRO:HD3	2.01	0.42
1:B:1331:GLN:HB3	1:B:1334:ARG:HD2	2.01	0.42
1:B:1382:CYS:HB2	1:B:1383:PRO:HD3	2.01	0.42
1:C:1326:GLU:OE1	1:C:1329:ARG:NE	2.46	0.42
1:C:1428:ASP:OD1	1:C:1464:LYS:HE3	2.20	0.42
1:A:1283:MET:O	1:A:1287:GLN:HG2	2.20	0.41
1:C:1390:TYR:CD1	1:C:1431:GLU:HG3	2.55	0.41
1:B:1318:ALA:O	1:B:1322:LYS:HG3	2.20	0.41
1:A:1374:PRO:HG3	1:A:1406:ARG:NH1	2.35	0.41
1:A:1331:GLN:HB3	1:A:1334:ARG:HD2	2.01	0.41
1:C:1382:CYS:HB2	1:C:1383:PRO:HD3	2.02	0.41
1:A:1393:ASN:OD1	1:A:1394:LEU:N	2.53	0.41
1:B:1449:ILE:HD11	1:B:1453:LEU:HB2	2.02	0.41
1:A:1449:ILE:HD11	1:A:1453:LEU:HB2	2.03	0.41
1:B:1340:GLU:OE1	1:D:1380:VAL:HA	2.21	0.40
1:D:1335:PHE:HB2	1:D:1372:ILE:HD12	2.02	0.40
1:C:1394:LEU:HD11	1:C:1398:LYS:HE3	2.03	0.40
1:D:1469:ASN:HA	1:D:1472:ILE:HB	2.02	0.40

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1385:ILE:HG12	1:A:1397:ILE:HG13	2.03	0.40
2:H:514:TYR:CE2	2:H:518:ARG:HD3	2.57	0.40
1:B:1299:LYS:HE2	1:D:1417:PRO:HB2	2.04	0.40
1:B:1379:LYS:NZ	1:B:1418:GLU:OE2	2.55	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	226/235 (96%)	225 (100%)	1 (0%)	0	100	100
1	B	224/235 (95%)	221 (99%)	3 (1%)	0	100	100
1	C	216/235 (92%)	215 (100%)	1 (0%)	0	100	100
1	D	224/235 (95%)	222 (99%)	2 (1%)	0	100	100
2	E	50/66 (76%)	50 (100%)	0	0	100	100
2	F	53/66 (80%)	53 (100%)	0	0	100	100
2	G	50/66 (76%)	50 (100%)	0	0	100	100
2	H	53/66 (80%)	53 (100%)	0	0	100	100
All	All	1096/1204 (91%)	1089 (99%)	7 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	203/209 (97%)	202 (100%)	1 (0%)	88	95
1	B	200/209 (96%)	199 (100%)	1 (0%)	88	95
1	C	197/209 (94%)	196 (100%)	1 (0%)	88	95
1	D	200/209 (96%)	197 (98%)	3 (2%)	65	85
2	E	39/61 (64%)	38 (97%)	1 (3%)	46	76
2	F	44/61 (72%)	43 (98%)	1 (2%)	50	78
2	G	39/61 (64%)	39 (100%)	0	100	100
2	H	38/61 (62%)	37 (97%)	1 (3%)	46	76
All	All	960/1080 (89%)	951 (99%)	9 (1%)	78	91

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	1427	TYR
1	D	1289	GLU
1	D	1375	GLU
1	D	1427	TYR
2	F	518	ARG
2	H	541	GLU
1	C	1427	TYR
1	A	1427	TYR
2	E	504	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	ACT	B	1501	-	3,3,3	0.97	0	3,3,3	0.80	0
3	ACT	A	1501	-	3,3,3	0.96	0	3,3,3	0.80	0
4	MLA	B	1502	-	6,6,6	1.32	0	7,7,7	0.95	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MLA	B	1502	-	-	1/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	1502	MLA	O1B-C1-C2-C3

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.3 Carbohydrates [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.4 Ligands [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.5 Other polymers [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.