



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 7, 2020 – 08:23 AM BST

PDB ID : 3WIR
Title : Crystal structure of kojibiose phosphorylase complexed with glucose
Authors : Okada, S.; Yamamoto, T.; Watanabe, H.; Nishimoto, T.; Chaen, H.; Fukuda, S.; Wakagi, T.; Fushinobu, S.
Deposited on : 2013-09-24
Resolution : 2.05 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1692 (2.04-2.04)
Clashscore	141614	1773 (2.04-2.04)
Ramachandran outliers	138981	1752 (2.04-2.04)
Sidechain outliers	138945	1752 (2.04-2.04)
RSRZ outliers	127900	1672 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	764	 2% 86% 12% ..
1	B	764	 2% 86% 11% ..
1	C	764	 % 85% 12% ..
1	D	764	 2% 84% 13% ..

2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 26306 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Kojibiose phosphorylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	756	6187	3985	1019	1162	21	0	0	0
1	B	756	6187	3985	1019	1162	21	0	0	0
1	C	756	6187	3985	1019	1162	21	0	0	0
1	D	756	6187	3985	1019	1162	21	0	0	0

There are 32 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	757	GLY	-	expression tag	UNP A4XGP2
A	758	SER	-	expression tag	UNP A4XGP2
A	759	HIS	-	expression tag	UNP A4XGP2
A	760	HIS	-	expression tag	UNP A4XGP2
A	761	HIS	-	expression tag	UNP A4XGP2
A	762	HIS	-	expression tag	UNP A4XGP2
A	763	HIS	-	expression tag	UNP A4XGP2
A	764	HIS	-	expression tag	UNP A4XGP2
B	757	GLY	-	expression tag	UNP A4XGP2
B	758	SER	-	expression tag	UNP A4XGP2
B	759	HIS	-	expression tag	UNP A4XGP2
B	760	HIS	-	expression tag	UNP A4XGP2
B	761	HIS	-	expression tag	UNP A4XGP2
B	762	HIS	-	expression tag	UNP A4XGP2
B	763	HIS	-	expression tag	UNP A4XGP2
B	764	HIS	-	expression tag	UNP A4XGP2
C	757	GLY	-	expression tag	UNP A4XGP2
C	758	SER	-	expression tag	UNP A4XGP2
C	759	HIS	-	expression tag	UNP A4XGP2
C	760	HIS	-	expression tag	UNP A4XGP2
C	761	HIS	-	expression tag	UNP A4XGP2

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	762	HIS	-	expression tag	UNP A4XGP2
C	763	HIS	-	expression tag	UNP A4XGP2
C	764	HIS	-	expression tag	UNP A4XGP2
D	757	GLY	-	expression tag	UNP A4XGP2
D	758	SER	-	expression tag	UNP A4XGP2
D	759	HIS	-	expression tag	UNP A4XGP2
D	760	HIS	-	expression tag	UNP A4XGP2
D	761	HIS	-	expression tag	UNP A4XGP2
D	762	HIS	-	expression tag	UNP A4XGP2
D	763	HIS	-	expression tag	UNP A4XGP2
D	764	HIS	-	expression tag	UNP A4XGP2

- Molecule 2 is beta-D-glucopyranose (three-letter code: BGC) (formula: C₆H₁₂O₆).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf		
2	A	1	Total	C	O	0	0
			12	6	6		
2	A	1	Total	C	O	0	0
			12	6	6		
2	B	1	Total	C	O	0	0
			12	6	6		
2	B	1	Total	C	O	0	0
			12	6	6		
2	C	1	Total	C	O	0	0
			12	6	6		
2	C	1	Total	C	O	0	0
			12	6	6		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	D	1	Total	C	O	0	0
			12	6	6		
2	D	1	Total	C	O	0	0
			12	6	6		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O P 5 4 1	0	0
4	B	1	Total O P 5 4 1	0	0
4	B	1	Total O P 5 4 1	0	0
4	C	1	Total O P 5 4 1	0	0
4	C	1	Total O P 5 4 1	0	0
4	D	1	Total O P 5 4 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	371	Total O 371 371	0	0
5	B	387	Total O 387 387	0	0
5	C	299	Total O 299 299	0	0
5	D	321	Total O 321 321	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Kojibiose phosphorylase




- Molecule 1: Kojibiose phosphorylase





- Molecule 1: Kojibiose phosphorylase





● Molecule 1: Kojibiose phosphorylase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	71.58Å 104.46Å 124.16Å 68.83° 86.02° 90.06°	Depositor
Resolution (Å)	46.26 – 2.05 46.26 – 2.05	Depositor EDS
% Data completeness (in resolution range)	97.0 (46.26-2.05) 97.0 (46.26-2.05)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.29 (at 2.05Å)	Xtrriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.191 , 0.251 0.196 , 0.254	Depositor DCC
R_{free} test set	10235 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	26.3	Xtrriage
Anisotropy	0.153	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 48.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	26306	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 20.80 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 8.0405e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, PO4, BGC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.93	0/6328	0.94	12/8550 (0.1%)
1	B	0.91	1/6328 (0.0%)	0.95	12/8550 (0.1%)
1	C	0.81	0/6328	0.88	6/8550 (0.1%)
1	D	0.82	1/6328 (0.0%)	0.93	14/8550 (0.2%)
All	All	0.87	2/25312 (0.0%)	0.93	44/34200 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	C	0	1
All	All	0	2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	658	THR	CB-CG2	-5.97	1.32	1.52
1	B	488	CYS	CB-SG	-5.47	1.72	1.81

The worst 5 of 44 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	657	ARG	NE-CZ-NH1	11.81	126.21	120.30
1	D	95	ARG	NE-CZ-NH1	-9.29	115.65	120.30
1	B	657	ARG	NE-CZ-NH2	-8.81	115.89	120.30
1	D	95	ARG	NE-CZ-NH2	8.66	124.63	120.30
1	D	657	ARG	NE-CZ-NH1	8.03	124.32	120.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	551	ASP	Peptide
1	C	255	ASN	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6187	0	6120	47	0
1	B	6187	0	6119	60	0
1	C	6187	0	6120	57	0
1	D	6187	0	6119	66	0
2	A	24	0	24	0	0
2	B	24	0	24	0	0
2	C	24	0	24	0	0
2	D	24	0	24	0	0
3	A	12	0	16	0	0
3	B	6	0	8	1	0
3	C	12	0	16	0	0
3	D	24	0	32	2	0
4	A	5	0	0	0	0
4	B	10	0	0	1	0
4	C	10	0	0	0	0
4	D	5	0	0	0	0
5	A	371	0	0	6	0
5	B	387	0	0	2	0
5	C	299	0	0	5	0
5	D	321	0	0	9	0
All	All	26306	0	24646	229	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 229 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:636:SER:HB3	1:D:658:THR:HG21	1.54	0.87
1:B:303:ALA:HB2	1:B:656:MET:HE1	1.58	0.86
1:B:303:ALA:HB2	1:B:656:MET:CE	2.07	0.85
1:C:10:ILE:HD12	1:C:99:LEU:HD21	1.65	0.78
1:D:709:GLU:CD	1:D:709:GLU:H	1.88	0.76

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	754/764 (99%)	724 (96%)	27 (4%)	3 (0%)	34 24
1	B	754/764 (99%)	729 (97%)	23 (3%)	2 (0%)	41 31
1	C	754/764 (99%)	726 (96%)	23 (3%)	5 (1%)	22 12
1	D	754/764 (99%)	717 (95%)	33 (4%)	4 (0%)	29 18
All	All	3016/3056 (99%)	2896 (96%)	106 (4%)	14 (0%)	29 18

5 of 14 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	742	LYS
1	D	259	VAL
1	D	741	GLU
1	D	258	ASP
1	B	742	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	669/676 (99%)	645 (96%)	24 (4%)	35	28
1	B	669/676 (99%)	651 (97%)	18 (3%)	44	38
1	C	669/676 (99%)	638 (95%)	31 (5%)	27	19
1	D	669/676 (99%)	631 (94%)	38 (6%)	20	12
All	All	2676/2704 (99%)	2565 (96%)	111 (4%)	30	23

5 of 111 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	277	LYS
1	C	650	ARG
1	D	721	LYS
1	C	288	ARG
1	C	539	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	593	GLN
1	A	752	GLN
1	B	101	GLN
1	B	575	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry

23 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	PO4	C	805	-	4,4,4	0.89	0	6,6,6	1.36	1 (16%)
2	BGC	B	801	-	12,12,12	1.19	1 (8%)	17,17,17	1.20	2 (11%)
3	GOL	D	804	-	5,5,5	0.78	0	5,5,5	1.10	0
2	BGC	C	802	-	12,12,12	1.02	1 (8%)	17,17,17	1.07	2 (11%)
2	BGC	C	801	-	12,12,12	0.53	0	17,17,17	1.34	2 (11%)
2	BGC	D	802	-	12,12,12	1.29	2 (16%)	17,17,17	1.71	6 (35%)
3	GOL	D	806	-	5,5,5	0.62	0	5,5,5	0.48	0
2	BGC	B	802	-	12,12,12	0.59	0	17,17,17	1.22	2 (11%)
3	GOL	A	804	-	5,5,5	0.83	0	5,5,5	1.19	1 (20%)
3	GOL	A	803	-	5,5,5	0.76	0	5,5,5	0.59	0
4	PO4	B	804	-	4,4,4	1.86	1 (25%)	6,6,6	1.54	1 (16%)
4	PO4	B	805	-	4,4,4	1.22	0	6,6,6	1.17	0
4	PO4	C	806	-	4,4,4	0.96	0	6,6,6	0.87	0
2	BGC	A	801	-	12,12,12	0.68	0	17,17,17	1.59	6 (35%)
3	GOL	D	805	-	5,5,5	1.13	1 (20%)	5,5,5	0.97	0
2	BGC	A	802	-	12,12,12	0.60	0	17,17,17	1.86	5 (29%)
3	GOL	B	803	-	5,5,5	0.47	0	5,5,5	0.98	0
2	BGC	D	801	-	12,12,12	0.66	0	17,17,17	1.00	1 (5%)
4	PO4	D	807	-	4,4,4	0.91	0	6,6,6	1.06	0
4	PO4	A	805	-	4,4,4	0.60	0	6,6,6	1.22	1 (16%)
3	GOL	D	803	-	5,5,5	0.73	0	5,5,5	0.42	0
3	GOL	C	803	-	5,5,5	0.69	0	5,5,5	0.47	0
3	GOL	C	804	-	5,5,5	0.42	0	5,5,5	0.27	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	803	-	-	0/4/4/4	-
2	BGC	A	801	-	-	0/2/22/22	0/1/1/1
3	GOL	D	805	-	-	2/4/4/4	-
3	GOL	D	806	-	-	4/4/4/4	-
2	BGC	C	802	-	-	0/2/22/22	0/1/1/1
3	GOL	A	804	-	-	1/4/4/4	-
3	GOL	B	803	-	-	3/4/4/4	-
2	BGC	D	801	-	-	0/2/22/22	0/1/1/1
2	BGC	B	801	-	-	0/2/22/22	0/1/1/1
3	GOL	D	804	-	-	2/4/4/4	-
3	GOL	D	803	-	-	1/4/4/4	-
2	BGC	B	802	-	-	0/2/22/22	0/1/1/1
2	BGC	C	801	-	-	0/2/22/22	0/1/1/1
2	BGC	D	802	-	-	0/2/22/22	0/1/1/1
2	BGC	A	802	-	-	0/2/22/22	0/1/1/1
3	GOL	C	803	-	-	4/4/4/4	-
3	GOL	C	804	-	-	2/4/4/4	-

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	802	BGC	O5-C5	-3.12	1.36	1.44
4	B	804	PO4	P-O1	3.00	1.57	1.50
2	B	801	BGC	O4-C4	2.58	1.49	1.43
2	C	802	BGC	O1-C1	2.40	1.47	1.39
3	D	805	GOL	O2-C2	2.39	1.50	1.43

The worst 5 of 30 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	802	BGC	O2-C2-C1	-3.49	101.07	109.16
4	B	804	PO4	O4-P-O3	3.32	118.62	107.97
2	B	802	BGC	O1-C1-O5	-3.19	100.82	110.38
2	C	801	BGC	C4-C3-C2	-3.15	105.33	110.82
2	B	801	BGC	C1-O5-C5	3.11	119.53	113.66

There are no chirality outliers.

5 of 19 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	D	804	GOL	O1-C1-C2-C3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	D	806	GOL	O1-C1-C2-C3
3	D	806	GOL	C1-C2-C3-O3
3	D	805	GOL	O1-C1-C2-C3
3	C	804	GOL	O1-C1-C2-C3

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	805	PO4	1	0
3	D	805	GOL	2	0
3	B	803	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	756/764 (98%)	-0.11	12 (1%) 72 74	13, 27, 53, 93	0
1	B	756/764 (98%)	-0.13	12 (1%) 72 74	13, 26, 51, 69	0
1	C	756/764 (98%)	-0.18	4 (0%) 91 92	17, 30, 54, 72	0
1	D	756/764 (98%)	-0.11	12 (1%) 72 74	16, 30, 55, 75	0
All	All	3024/3056 (98%)	-0.13	40 (1%) 77 79	13, 29, 53, 93	0

The worst 5 of 40 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	257	GLY	4.4
1	D	743	GLY	4.2
1	A	550	PRO	3.9
1	A	568	VAL	3.8
1	A	549	HIS	3.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	GOL	D	803	6/6	0.72	0.18	44,54,61,61	0
3	GOL	D	804	6/6	0.73	0.29	55,57,60,64	0
3	GOL	D	806	6/6	0.78	0.12	46,47,49,49	0
3	GOL	C	803	6/6	0.79	0.18	46,55,59,62	0
3	GOL	A	804	6/6	0.85	0.24	33,35,40,41	0
3	GOL	D	805	6/6	0.87	0.13	27,36,38,50	0
3	GOL	A	803	6/6	0.88	0.14	29,41,45,47	0
3	GOL	C	804	6/6	0.91	0.10	47,49,50,55	0
3	GOL	B	803	6/6	0.92	0.21	30,40,42,42	0
2	BGC	A	802	12/12	0.93	0.12	21,26,29,32	0
2	BGC	B	802	12/12	0.94	0.13	19,26,29,31	0
2	BGC	C	802	12/12	0.95	0.10	20,25,29,30	0
2	BGC	D	802	12/12	0.95	0.11	21,26,27,27	0
2	BGC	A	801	12/12	0.95	0.11	19,21,23,24	0
2	BGC	D	801	12/12	0.96	0.14	21,24,25,25	0
4	PO4	B	805	5/5	0.96	0.10	42,43,47,47	0
4	PO4	C	806	5/5	0.96	0.09	45,46,46,50	0
2	BGC	B	801	12/12	0.96	0.13	20,22,24,26	0
2	BGC	C	801	12/12	0.97	0.11	19,24,25,25	0
4	PO4	D	807	5/5	0.98	0.07	22,25,27,28	0
4	PO4	C	805	5/5	0.99	0.07	22,24,26,27	0
4	PO4	B	804	5/5	0.99	0.11	18,21,23,26	0
4	PO4	A	805	5/5	0.99	0.09	20,20,23,24	0

6.5 Other polymers

There are no such residues in this entry.