



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 13, 2018 – 03:01 pm GMT

PDB ID : 5WIV
Title : Structure of the sodium-bound human D4 Dopamine receptor in complex with Nemonapride
Authors : Wacker, D.; Wang, S.; Levit, A.; Che, T.; Betz, R.M.; McCorvy, J.D.; Venkatakrisnan, A.J.; Huang, X.-P.; Dror, R.O.; Shoichet, B.K.; Roth, B.L.
Deposited on : 2017-07-20
Resolution : 2.14 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : 1.13
EDS : trunk31020
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk31020


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.14 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	2128 (2.16-2.12)
Clashscore	122126	2253 (2.16-2.12)
Ramachandran outliers	120053	2223 (2.16-2.12)
Sidechain outliers	120020	2222 (2.16-2.12)
RSRZ outliers	108989	2086 (2.16-2.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	422	


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	Cl	N			O
2	A	1	27	21	1	3	2	0	0

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O P		
3	A	1	5	4 1	0	0
3	A	1	5	4 1	0	0
3	A	1	5	4 1	0	0
3	A	1	5	4 1	0	0
3	A	1	5	4 1	0	0

- Molecule 4 is OLEIC ACID (three-letter code: OLA) (formula: C₁₈H₃₄O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			17	15	2		
4	A	1	Total	C	O	0	0
			17	15	2		
4	A	1	Total	C	O	0	0
			17	15	2		
4	A	1	Total	C	O	0	0
			20	18	2		
4	A	1	Total	C	O	0	0
			20	18	2		
4	A	1	Total	C	O	0	0
			16	14	2		
4	A	1	Total	C	O	0	0
			20	18	2		
4	A	1	Total	C	O	0	0
			20	18	2		

- Molecule 5 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 7 4 3	0	0
5	A	1	Total C O 7 4 3	0	0
5	A	1	Total C O 7 4 3	0	0
5	A	1	Total C O 7 4 3	0	0
5	A	1	Total C O 7 4 3	0	0
5	A	1	Total C O 7 4 3	0	0
5	A	1	Total C O 7 4 3	0	0
5	A	1	Total C O 7 4 3	0	0
5	A	1	Total C O 7 4 3	0	0

- Molecule 6 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total Na 1 1	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	67	Total O 67 67	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: D(4) dopamine receptor, soluble cytochrome b562 chimera



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	67.86Å 162.84Å 83.18Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.69 – 2.14 29.69 – 2.14	Depositor EDS
% Data completeness (in resolution range)	96.8 (29.69-2.14) 96.8 (29.69-2.14)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.27 (at 2.14Å)	Xtrriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.195 , 0.227 0.200 , 0.205	Depositor DCC
R_{free} test set	1256 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	47.4	Xtrriage
Anisotropy	0.311	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 68.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3149	wwPDB-VP
Average B, all atoms (Å ²)	59.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.86% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: OLA, PO4, PEG, AQD, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.41	0/2900	0.53	0/3969

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2819	0	2821	30	1
2	A	27	0	0	0	0
3	A	25	0	0	0	0
4	A	147	0	226	1	0
5	A	63	0	81	1	0
6	A	1	0	0	0	0
7	A	67	0	0	4	0
All	All	3149	0	3128	32	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1018:GLU:OE1	7:A:1301:HOH:O	2.09	0.71
1:A:1007:TRP:HH2	1:A:1103:GLN:HE21	1.40	0.69
1:A:123:ILE:HG12	1:A:403:PHE:CZ	2.31	0.65
1:A:1106:LEU:HD23	1:A:385:ILE:HD11	1.79	0.65
1:A:1007:TRP:CH2	1:A:1103:GLN:HG3	2.32	0.65
1:A:93:TYR:O	1:A:97:GLN:HG2	2.03	0.58
5:A:1215:PEG:O4	7:A:1302:HOH:O	2.17	0.56
1:A:127[B]:CYS:HA	1:A:208:MET:HG3	1.88	0.55
1:A:127[A]:CYS:SG	1:A:204:PRO:HB3	2.51	0.51
1:A:127[A]:CYS:HA	1:A:208:MET:HG3	1.92	0.51
1:A:394:ARG:O	1:A:397:PRO:HD2	2.12	0.50
1:A:101:TRP:CH2	1:A:111:LEU:HD12	2.47	0.50
1:A:435:TRP:O	1:A:439:VAL:HG13	2.12	0.50
1:A:436:LEU:O	1:A:439:VAL:HG22	2.13	0.49
1:A:150:ARG:HA	7:A:1362:HOH:O	2.13	0.49
1:A:123:ILE:HG12	1:A:403:PHE:CE1	2.48	0.48
1:A:1003:LEU:HD21	1:A:385:ILE:HG13	1.95	0.48
4:A:1210:OLA:H71	4:A:1210:OLA:H41	1.52	0.47
1:A:223:GLU:OE2	1:A:390:ARG:HD3	2.15	0.47
1:A:1046:PRO:HA	1:A:1049:GLU:CD	2.35	0.46
1:A:222:TRP:CZ2	1:A:1047:LYS:HG2	2.51	0.46
1:A:203:LEU:HB3	1:A:204:PRO:HD3	1.98	0.45
1:A:175:ASN:ND2	7:A:1305:HOH:O	2.41	0.44
1:A:1072:ILE:HG13	1:A:1094:LEU:HD11	2.00	0.44
1:A:124:PHE:HB3	1:A:159:THR:HG23	2.00	0.44
1:A:1003:LEU:CD2	1:A:385:ILE:HG13	2.49	0.43
1:A:211:LEU:HD23	1:A:211:LEU:HA	1.84	0.42
1:A:208:MET:HG2	1:A:212:TYR:CE2	2.55	0.42
1:A:1006:ASN:HD21	1:A:1039:ASP:HB3	1.85	0.42
1:A:1059:LYS:HE3	1:A:1059:LYS:HB3	1.81	0.41
1:A:1071:GLN:HB3	1:A:1094:LEU:HD12	2.01	0.41
1:A:119:CYS:O	1:A:123:ILE:HG13	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:145:ARG:NH1	1:A:171:LEU:O[4_555]	2.14	0.06

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	384/422 (91%)	379 (99%)	5 (1%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	286/330 (87%)	285 (100%)	1 (0%)	93 95

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1096	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1103	GLN
1	A	414	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 24 ligands modelled in this entry, 1 is monoatomic - leaving 23 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	AQD	A	1201	-	29,29,29	2.08	8 (27%)	40,40,40	1.89	10 (25%)
3	PO4	A	1202	-	4,4,4	0.68	0	6,6,6	0.43	0
3	PO4	A	1203	-	4,4,4	0.69	0	6,6,6	0.48	0
3	PO4	A	1204	-	4,4,4	0.76	0	6,6,6	0.41	0
3	PO4	A	1205	-	4,4,4	0.67	0	6,6,6	0.44	0
3	PO4	A	1206	-	4,4,4	0.71	0	6,6,6	0.37	0
4	OLA	A	1207	-	13,16,19	0.24	0	12,16,19	0.64	0
4	OLA	A	1208	-	13,16,19	0.26	0	12,16,19	0.61	0
4	OLA	A	1209	-	13,16,19	0.26	0	12,16,19	0.73	0
4	OLA	A	1210	-	16,19,19	0.22	0	15,19,19	0.63	0
4	OLA	A	1211	-	16,19,19	0.24	0	15,19,19	0.60	0
4	OLA	A	1212	-	12,15,19	0.25	0	11,15,19	0.69	0
4	OLA	A	1213	-	16,19,19	0.29	0	15,19,19	0.43	0
4	OLA	A	1214	-	16,19,19	0.28	0	15,19,19	0.51	0
5	PEG	A	1215	-	6,6,6	1.44	1 (16%)	5,5,5	1.54	0
5	PEG	A	1216	-	6,6,6	1.45	1 (16%)	5,5,5	1.56	1 (20%)
5	PEG	A	1217	-	6,6,6	1.44	1 (16%)	5,5,5	1.55	0
5	PEG	A	1218	-	6,6,6	1.44	1 (16%)	5,5,5	1.53	0
5	PEG	A	1219	-	6,6,6	1.44	1 (16%)	5,5,5	1.45	0
5	PEG	A	1220	-	6,6,6	1.43	1 (16%)	5,5,5	1.49	0
5	PEG	A	1221	-	6,6,6	1.46	1 (16%)	5,5,5	1.43	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	PEG	A	1222	-	6,6,6	1.42	1 (16%)	5,5,5	1.46	0
5	PEG	A	1223	-	6,6,6	1.45	1 (16%)	5,5,5	1.45	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	AQD	A	1201	-	-	0/16/29/29	0/3/3/3
3	PO4	A	1202	-	-	0/0/0/0	0/0/0/0
3	PO4	A	1203	-	-	0/0/0/0	0/0/0/0
3	PO4	A	1204	-	-	0/0/0/0	0/0/0/0
3	PO4	A	1205	-	-	0/0/0/0	0/0/0/0
3	PO4	A	1206	-	-	0/0/0/0	0/0/0/0
4	OLA	A	1207	-	-	0/12/14/17	0/0/0/0
4	OLA	A	1208	-	-	0/12/14/17	0/0/0/0
4	OLA	A	1209	-	-	0/12/14/17	0/0/0/0
4	OLA	A	1210	-	-	0/15/17/17	0/0/0/0
4	OLA	A	1211	-	-	0/15/17/17	0/0/0/0
4	OLA	A	1212	-	-	0/11/13/17	0/0/0/0
4	OLA	A	1213	-	-	0/15/17/17	0/0/0/0
4	OLA	A	1214	-	-	0/15/17/17	0/0/0/0
5	PEG	A	1215	-	-	0/4/4/4	0/0/0/0
5	PEG	A	1216	-	-	0/4/4/4	0/0/0/0
5	PEG	A	1217	-	-	0/4/4/4	0/0/0/0
5	PEG	A	1218	-	-	0/4/4/4	0/0/0/0
5	PEG	A	1219	-	-	0/4/4/4	0/0/0/0
5	PEG	A	1220	-	-	0/4/4/4	0/0/0/0
5	PEG	A	1221	-	-	0/4/4/4	0/0/0/0
5	PEG	A	1222	-	-	0/4/4/4	0/0/0/0
5	PEG	A	1223	-	-	0/4/4/4	0/0/0/0

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1201	AQD	CAY-NBA	-4.24	1.40	1.47
2	A	1201	AQD	CAY-CAZ	-3.20	1.46	1.52
5	A	1221	PEG	O2-C2	-2.78	1.29	1.42
5	A	1216	PEG	O2-C2	-2.75	1.29	1.42
5	A	1223	PEG	O2-C2	-2.73	1.29	1.42
5	A	1217	PEG	O2-C2	-2.72	1.29	1.42

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	1220	PEG	O2-C2	-2.72	1.29	1.42
5	A	1219	PEG	O2-C2	-2.69	1.30	1.42
5	A	1215	PEG	O2-C2	-2.69	1.30	1.42
5	A	1222	PEG	O2-C2	-2.68	1.30	1.42
5	A	1218	PEG	O2-C2	-2.66	1.30	1.42
2	A	1201	AQD	CAO-NBA	-2.48	1.43	1.47
2	A	1201	AQD	CAO-CAT	2.55	1.56	1.51
2	A	1201	AQD	OAR-CAW	3.28	1.42	1.37
2	A	1201	AQD	CAV-NAP	4.20	1.45	1.37
2	A	1201	AQD	CAS-NAQ	4.29	1.43	1.34
2	A	1201	AQD	CAM-CAN	5.17	1.63	1.52

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1201	AQD	CAM-CAZ-NAQ	-4.22	105.84	112.75
2	A	1201	AQD	CAO-NBA-CAY	-3.78	108.27	113.48
2	A	1201	AQD	CAZ-NAQ-CAS	-3.62	116.28	123.02
2	A	1201	AQD	CAW-CAX-CAS	-2.58	121.39	126.14
2	A	1201	AQD	OAR-CAW-CAK	-2.10	120.59	124.15
5	A	1216	PEG	O2-C3-C4	2.02	119.14	110.10
2	A	1201	AQD	CAM-CAZ-CAY	2.30	107.94	104.07
2	A	1201	AQD	CAL-CAX-CAW	2.30	121.51	118.31
2	A	1201	AQD	CAK-CAV-CAU	2.39	120.71	118.12
2	A	1201	AQD	CAX-CAS-NAQ	2.78	122.15	116.86
2	A	1201	AQD	CAZ-CAY-NBA	5.73	106.27	101.46

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1210	OLA	1	0
5	A	1215	PEG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	379/422 (89%)	0.41	33 (8%) 10 13	37, 54, 93, 116	0

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	463	LEU	6.8
1	A	32	GLY	5.4
1	A	453	ALA	4.8
1	A	1096	THR	4.4
1	A	1105	TYR	4.2
1	A	457	ASN	4.0
1	A	459	PHE	3.9
1	A	144	ASN	3.9
1	A	61	THR	3.7
1	A	146	GLN	3.7
1	A	464	ARG	3.6
1	A	1101	TYR	3.6
1	A	1097	THR	3.6
1	A	147	GLY	3.6
1	A	33	GLN	3.5
1	A	462	ALA	3.5
1	A	129	ILE	3.2
1	A	461	LYS	3.2
1	A	1050	ASP	3.2
1	A	1061	PHE	3.0
1	A	145	ARG	2.9
1	A	1098	ARG	2.8
1	A	1102	ILE	2.5
1	A	182	PRO	2.5
1	A	1053	PRO	2.4
1	A	1049	GLU	2.3
1	A	1063	HIS	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	126	LEU	2.3
1	A	1052	SER	2.2
1	A	127[A]	CYS	2.1
1	A	1007	TRP	2.1
1	A	1059	LYS	2.0
1	A	460	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	OLA	A	1213	20/20	0.43	0.39	65,79,101,102	0
5	PEG	A	1220	7/7	0.67	0.16	76,82,93,93	0
5	PEG	A	1219	7/7	0.68	0.21	78,84,87,89	0
4	OLA	A	1210	20/20	0.69	0.29	67,79,89,95	0
4	OLA	A	1209	17/20	0.73	0.27	52,66,72,72	0
5	PEG	A	1221	7/7	0.75	0.17	65,81,95,100	0
3	PO4	A	1204	5/5	0.78	0.40	54,55,57,61	5
4	OLA	A	1214	20/20	0.79	0.21	59,71,77,78	0
5	PEG	A	1222	7/7	0.79	0.28	72,77,83,84	0
5	PEG	A	1218	7/7	0.81	0.15	68,70,75,78	0
4	OLA	A	1211	20/20	0.82	0.16	54,71,93,99	0
4	OLA	A	1212	16/20	0.84	0.18	57,67,83,97	0
5	PEG	A	1217	7/7	0.84	0.19	62,64,78,82	0
5	PEG	A	1215	7/7	0.86	0.16	65,73,78,79	0
4	OLA	A	1207	17/20	0.86	0.13	56,70,80,86	0
5	PEG	A	1216	7/7	0.88	0.14	67,70,74,77	0
3	PO4	A	1203	5/5	0.88	0.39	80,94,101,119	5

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	PO4	A	1206	5/5	0.89	0.21	93,98,106,110	5
4	OLA	A	1208	17/20	0.90	0.14	53,64,98,107	0
5	PEG	A	1223	7/7	0.91	0.10	67,68,72,73	0
3	PO4	A	1205	5/5	0.91	0.22	69,95,97,98	0
6	NA	A	1224	1/1	0.93	0.14	67,67,67,67	0
2	AQD	A	1201	27/27	0.95	0.16	36,45,62,67	0
3	PO4	A	1202	5/5	0.98	0.07	60,74,76,81	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.