



# wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 16, 2024 – 12:09 PM JST

PDB ID : 8WIE  
Title : Unlocking Immunogenic Potential: Innovating a Peptide/Ferritin Fusion Tag Nano-Delivery Platform from de novo design to Significantly Enhance Antigenicity of the Rabies Virus Glycoprotein Domain III  
Authors : Fu, D.; Wang, M.; Guo, Y.  
Deposited on : 2023-09-24  
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.002 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.38.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*




The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	5963 (2.30-2.30)
Clashscore	180529	6698 (2.30-2.30)
Ramachandran outliers	177936	6640 (2.30-2.30)
Sidechain outliers	177891	6640 (2.30-2.30)
RSRZ outliers	164620	5963 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	200	 6% 86% 8% 5%
1	B	200	 5% 86% 7% 6%
1	C	200	 4% 86% 8% 5%
1	D	200	 5% 84% 10% 6%
1	E	200	 4% 82% 11% 6%
1	F	200	 4% 83% 10% 6%

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 9759 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Peptide 10-1,Ferritin heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	C	190	1554	978	271	298	7	0	0	0
1	D	189	1550	976	270	297	7	0	0	0
1	E	188	1546	974	269	296	7	0	0	0
1	F	187	1537	968	267	295	7	0	0	0
1	A	190	1554	978	271	298	7	0	0	0
1	B	188	1546	974	269	296	7	0	0	0

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	15	ARG	GLN	engineered mutation	UNP P02794
C	23	LYS	ARG	engineered mutation	UNP P02794
C	26	THR	ASN	engineered mutation	UNP P02794
C	87	GLN	LYS	engineered mutation	UNP P02794
C	110	GLU	ASN	engineered mutation	UNP P02794
C	114	GLU	SER	engineered mutation	UNP P02794
C	117	ASN	GLU	engineered mutation	UNP P02794
D	15	ARG	GLN	engineered mutation	UNP P02794
D	23	LYS	ARG	engineered mutation	UNP P02794
D	26	THR	ASN	engineered mutation	UNP P02794
D	87	GLN	LYS	engineered mutation	UNP P02794
D	110	GLU	ASN	engineered mutation	UNP P02794
D	114	GLU	SER	engineered mutation	UNP P02794
D	117	ASN	GLU	engineered mutation	UNP P02794
E	15	ARG	GLN	engineered mutation	UNP P02794
E	23	LYS	ARG	engineered mutation	UNP P02794
E	26	THR	ASN	engineered mutation	UNP P02794

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	87	GLN	LYS	engineered mutation	UNP P02794
E	110	GLU	ASN	engineered mutation	UNP P02794
E	114	GLU	SER	engineered mutation	UNP P02794
E	117	ASN	GLU	engineered mutation	UNP P02794
F	15	ARG	GLN	engineered mutation	UNP P02794
F	23	LYS	ARG	engineered mutation	UNP P02794
F	26	THR	ASN	engineered mutation	UNP P02794
F	87	GLN	LYS	engineered mutation	UNP P02794
F	110	GLU	ASN	engineered mutation	UNP P02794
F	114	GLU	SER	engineered mutation	UNP P02794
F	117	ASN	GLU	engineered mutation	UNP P02794
A	15	ARG	GLN	engineered mutation	UNP P02794
A	23	LYS	ARG	engineered mutation	UNP P02794
A	26	THR	ASN	engineered mutation	UNP P02794
A	87	GLN	LYS	engineered mutation	UNP P02794
A	110	GLU	ASN	engineered mutation	UNP P02794
A	114	GLU	SER	engineered mutation	UNP P02794
A	117	ASN	GLU	engineered mutation	UNP P02794
B	15	ARG	GLN	engineered mutation	UNP P02794
B	23	LYS	ARG	engineered mutation	UNP P02794
B	26	THR	ASN	engineered mutation	UNP P02794
B	87	GLN	LYS	engineered mutation	UNP P02794
B	110	GLU	ASN	engineered mutation	UNP P02794
B	114	GLU	SER	engineered mutation	UNP P02794
B	117	ASN	GLU	engineered mutation	UNP P02794

- Molecule 2 is FE (III) ION (three-letter code: FE) (formula: Fe) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	D	1	Total Fe 1 1	0	0

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	D	1	Total Ca 1 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	C	82	Total O 82 82	0	0
4	D	70	Total O 70 70	0	0
4	E	73	Total O 73 73	0	0
4	F	69	Total O 69 69	0	0
4	A	87	Total O 87 87	0	0
4	B	89	Total O 89 89	0	0

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Peptide 10-1,Ferritin heavy chain

Chain C: 





- Molecule 1: Peptide 10-1,Ferritin heavy chain

Chain D: 





- Molecule 1: Peptide 10-1,Ferritin heavy chain

Chain E: 




- Molecule 1: Peptide 10-1,Ferritin heavy chain

Chain F: 





- Molecule 1: Peptide 10-1,Ferritin heavy chain

Chain A:  6% 86% 8% 5%



● Molecule 1: Peptide 10-1, Ferritin heavy chain

Chain B:  5% 86% 7% 6%



## 4 Data and refinement statistics i

Property	Value	Source
Space group	I 4	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	142.83Å 142.83Å 183.92Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	33.67 – 2.30 33.67 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.6 (33.67-2.30) 99.6 (33.67-2.30)	Depositor EDS
$R_{merge}$	0.16	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.13 (at 2.29Å)	Xtrriage
Refinement program	PHENIX (1.19.2_4158: ???)	Depositor
R, $R_{free}$	0.194 , 0.212 0.196 , 0.213	Depositor DCC
$R_{free}$ test set	79525 reflections (2.45%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	31.0	Xtrriage
Anisotropy	0.023	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 47.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.52$ , $\langle L^2 \rangle = 0.36$	Xtrriage
Estimated twinning fraction	0.005 for -h,k,-l	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	9759	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	33.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.93% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FE, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.27	0/1584	0.46	0/2131
1	B	0.25	0/1576	0.43	0/2121
1	C	0.28	0/1584	0.49	0/2131
1	D	0.25	0/1580	0.43	0/2126
1	E	0.25	0/1576	0.45	0/2121
1	F	0.25	0/1567	0.43	0/2110
All	All	0.26	0/9467	0.45	0/12740

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1554	0	1513	9	0
1	B	1546	0	1507	7	0
1	C	1554	0	1513	10	0
1	D	1550	0	1510	14	0
1	E	1546	0	1507	16	0
1	F	1537	0	1494	13	0
2	D	1	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	D	1	0	0	0	0
4	A	87	0	0	1	0
4	B	89	0	0	0	0
4	C	82	0	0	0	0
4	D	70	0	0	2	0
4	E	73	0	0	0	0
4	F	69	0	0	0	0
All	All	9759	0	9044	65	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 65 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:98:LEU:HD23	1:E:162:PRO:HD3	1.69	0.74
1:D:98:LEU:HD23	1:D:162:PRO:HD3	1.71	0.72
1:F:98:LEU:HD23	1:F:162:PRO:HD3	1.71	0.72
1:E:161:ALA:HB1	1:E:162:PRO:HD2	1.73	0.71
1:C:64:ARG:NH1	1:C:68:GLU:OE2	2.20	0.70

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	186/200 (93%)	184 (99%)	2 (1%)	0	100	100
1	B	184/200 (92%)	181 (98%)	3 (2%)	0	100	100
1	C	186/200 (93%)	182 (98%)	3 (2%)	1 (0%)	25	32
1	D	185/200 (92%)	180 (97%)	4 (2%)	1 (0%)	25	32

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	E	184/200 (92%)	180 (98%)	3 (2%)	1 (0%)	25	32
1	F	183/200 (92%)	178 (97%)	4 (2%)	1 (0%)	25	32
All	All	1108/1200 (92%)	1085 (98%)	19 (2%)	4 (0%)	30	39

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	161	ALA
1	D	161	ALA
1	E	161	ALA
1	F	161	ALA

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	169/175 (97%)	163 (96%)	6 (4%)	30	44
1	B	169/175 (97%)	164 (97%)	5 (3%)	36	52
1	C	169/175 (97%)	164 (97%)	5 (3%)	36	52
1	D	169/175 (97%)	167 (99%)	2 (1%)	67	81
1	E	169/175 (97%)	166 (98%)	3 (2%)	54	71
1	F	168/175 (96%)	163 (97%)	5 (3%)	36	52
All	All	1013/1050 (96%)	987 (97%)	26 (3%)	41	58

5 of 26 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	120	LYS
1	A	50	LYS
1	B	84	GLN
1	A	33	TYR
1	A	64	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	74	GLN
1	B	87	GLN
1	B	137	HIS
1	D	140	ASN
1	E	84	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	190/200 (95%)	0.17	11 (5%) 30 32	21, 30, 50, 69	0
1	B	188/200 (94%)	0.07	10 (5%) 33 34	20, 29, 50, 61	0
1	C	190/200 (95%)	0.20	7 (3%) 45 47	24, 31, 51, 63	0
1	D	189/200 (94%)	0.21	10 (5%) 33 34	24, 33, 54, 67	0
1	E	188/200 (94%)	0.26	9 (4%) 36 37	24, 33, 54, 64	0
1	F	187/200 (93%)	0.09	8 (4%) 40 41	24, 32, 52, 65	0
All	All	1132/1200 (94%)	0.17	55 (4%) 36 37	20, 31, 52, 69	0

The worst 5 of 55 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	161	ALA	6.2
1	A	161	ALA	5.5
1	D	161	ALA	5.5
1	F	161	ALA	5.4
1	C	161	ALA	5.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	FE	D	201	1/1	0.73	0.20	110,110,110,110	0
3	CA	D	202	1/1	0.97	0.07	40,40,40,40	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around FE D 201:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







## 6.5 Other polymers ⓘ

There are no such residues in this entry.