



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 14, 2018 – 01:44 pm GMT

PDB ID : 5WL1
Title : Crystal Structure of Human CD1b in Complex with PG
Authors : Shahine, A.; Gras, S.; Rossjohn, J.
Deposited on : 2017-07-25
Resolution : 1.38 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : 1.13
EDS : trunk31020
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk31020


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.38 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	2404 (1.40-1.36)
Clashscore	122126	2520 (1.40-1.36)
Ramachandran outliers	120053	2464 (1.40-1.36)
Sidechain outliers	120020	2463 (1.40-1.36)
RSRZ outliers	108989	2346 (1.40-1.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	300	<p>3% 90% 7%</p>
2	B	99	<p>4% 97%</p>

2 Entry composition

There are 13 unique types of molecules in this entry. The entry contains 7131 atoms, of which 3120 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called T-cell surface glycoprotein CD1b.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	280	4323	1434	2074	385	420	10	0	21	0


There are 23 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	279	GLY	-	expression tag	UNP P29016
A	280	SER	-	expression tag	UNP P29016
A	281	GLY	-	expression tag	UNP P29016
A	282	LEU	-	expression tag	UNP P29016
A	283	ASN	-	expression tag	UNP P29016
A	284	ASP	-	expression tag	UNP P29016
A	285	ILE	-	expression tag	UNP P29016
A	286	PHE	-	expression tag	UNP P29016
A	287	GLU	-	expression tag	UNP P29016
A	288	ALA	-	expression tag	UNP P29016
A	289	GLN	-	expression tag	UNP P29016
A	290	LYS	-	expression tag	UNP P29016
A	291	ILE	-	expression tag	UNP P29016
A	292	GLU	-	expression tag	UNP P29016
A	293	TRP	-	expression tag	UNP P29016
A	294	HIS	-	expression tag	UNP P29016
A	295	GLU	-	expression tag	UNP P29016
A	296	HIS	-	expression tag	UNP P29016
A	297	HIS	-	expression tag	UNP P29016
A	298	HIS	-	expression tag	UNP P29016
A	299	HIS	-	expression tag	UNP P29016
A	300	HIS	-	expression tag	UNP P29016
A	301	HIS	-	expression tag	UNP P29016

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
2	B	98	1545	519	729	139	156	2	0	4	0

- Molecule 3 is tetracosyl octadecanoate (three-letter code: CUY) (formula: $C_{42}H_{84}O_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
3	A	1	87	33	52	2	0	0

- Molecule 4 is IODIDE ION (three-letter code: IOD) (formula: I).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	I		
4	A	1	1	1	0	0

- Molecule 5 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	N	O		
5	A	1	Total	C	H	N	O	0	0
			25	8	11	1	5		
5	A	1	Total	C	H	N	O	0	0
			28	8	14	1	5		
5	A	1	Total	C	H	N	O	0	0
			25	8	11	1	5		
5	A	1	Total	C	H	N	O	0	0
			27	8	13	1	5		
5	A	1	Total	C	H	N	O	0	0
			25	8	11	1	5		
5	A	1	Total	C	H	N	O	0	0
			27	8	13	1	5		

- Molecule 6 is ALPHA-L-FUCOSE (three-letter code: FUC) (formula: C₆H₁₂O₅).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
			Total	C	H			O
6	A	1	21	6	11	4	0	0
6	A	1	21	6	11	4	0	0
6	A	1	21	6	11	4	0	0
6	A	1	21	6	11	4	0	0
6	A	1	21	6	11	4	0	0

- Molecule 7 is ALPHA-D-MANNOSE (three-letter code: MAN) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	A	1	Total	C	H	O	0	0
			21	6	10	5		
7	A	1	Total	C	H	O	0	0
			21	6	10	5		
7	A	1	Total	C	H	O	0	0
			21	6	10	5		

- Molecule 8 is BETA-D-MANNOSE (three-letter code: BMA) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	A	1	Total	C	H	O	0	0
			20	6	9	5		
8	A	1	Total	C	H	O	0	0
			20	6	9	5		

- Molecule 9 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	1	Total	Cl	0	0
			1	1		

- Molecule 10 is SODIUM ION (three-letter code: NA) (formula: Na).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	B	1	Total	Na	0	0
			1	1		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	1	Total	Na	0	0
			1	1		

- Molecule 11 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
11	A	1	Total	C	H	O	0	0
			10	2	6	2		
11	A	1	Total	C	H	O	0	0
			10	2	6	2		
11	B	1	Total	C	H	O	0	0
			10	2	6	2		

- Molecule 12 is (19S,22R,25R)-22,25,26-trihydroxy-16,22-dioxo-17,21,23-trioxa-22lambda 5 -phosphahexacosan-19-yl (9E)-octadec-9-enoate (three-letter code: D3D) (formula: C₄₀H₇₇O₁₀P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	O	P		
12	A	1	120	38	71	10	1	0	0


- Molecule 13 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
13	A	496	Total	O	0	0
			496	496		
13	B	161	Total	O	0	0
			161	161		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: T-cell surface glycoprotein CD1b

Chain A: 



- Molecule 2: Beta-2-microglobulin

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	57.89Å 80.28Å 92.70Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.34 – 1.38 27.63 – 1.38	Depositor EDS
% Data completeness (in resolution range)	98.2 (30.34-1.38) 98.2 (27.63-1.38)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.97 (at 1.38Å)	Xtrriage
Refinement program	BUSTER 2.10.2	Depositor
R, R_{free}	0.188 , 0.206 0.194 , 0.208	Depositor DCC
R_{free} test set	4336 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å ²)	14.1	Xtrriage
Anisotropy	0.168	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 46.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7131	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.48% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, NAG, CL, NA, EDO, CUY, FUC, D3D, IOD, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.49	0/2375	0.66	0/3224
2	B	0.43	0/860	0.60	0/1167
All	All	0.48	0/3235	0.64	0/4391

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2249	2074	2091	10	0
2	B	816	729	743	1	0
3	A	35	52	0	1	0
4	A	1	0	0	0	0
5	A	84	73	68	2	0
6	A	50	55	50	1	0
7	A	33	30	30	0	0
8	A	22	18	17	0	0
9	A	1	0	0	0	0
10	A	1	0	0	0	0
10	B	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
11	A	8	12	12	0	0
11	B	4	6	0	0	0
12	A	49	71	0	4	0
13	A	496	0	0	7	0
13	B	161	0	0	0	0
All	All	4011	3120	3011	14	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (14) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:283:ASN:C	13:A:502:HOH:O	2.18	0.82
1:A:228:GLN:HG3	13:A:804:HOH:O	1.79	0.81
12:A:423:D3D:C16	13:A:501:HOH:O	2.29	0.79
1:A:269:GLU:HG3	13:A:503:HOH:O	1.95	0.67
12:A:423:D3D:O2	13:A:501:HOH:O	2.15	0.63
1:A:208:VAL:HG21	1:A:218[B]:VAL:HG11	1.83	0.60
5:A:403:NAG:H62	6:A:406:FUC:O2	2.02	0.60
3:A:401:CUY:C2	12:A:423:D3D:C37	2.84	0.56
1:A:23:TRP:CZ2	5:A:404:NAG:H82	2.44	0.52
1:A:228:GLN:CG	13:A:804:HOH:O	2.47	0.50
1:A:72:VAL:HG21	12:A:423:D3D:C21	2.45	0.46
1:A:36:GLN:NE2	13:A:504:HOH:O	2.51	0.43
1:A:120:GLY:O	2:B:3:ARG:NH2	2.48	0.42
1:A:262[B]:ARG:CZ	1:A:274:ILE:HD11	2.49	0.42

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	299/300 (100%)	297 (99%)	2 (1%)	0	100	100
2	B	100/99 (101%)	100 (100%)	0	0	100	100
All	All	399/399 (100%)	397 (100%)	2 (0%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	242/247 (98%)	242 (100%)	0	100	100
2	B	93/94 (99%)	92 (99%)	1 (1%)	76	51
All	All	335/341 (98%)	334 (100%)	1 (0%)	93	83

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	70	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	36	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 25 ligands modelled in this entry, 4 are monoatomic - leaving 21 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	CUY	A	401	-	34,34,43	0.69	1 (2%)	34,34,43	0.84	1 (2%)
5	NAG	A	403	1,5,6	14,14,15	0.55	0	17,19,21	2.67	5 (29%)
5	NAG	A	404	5	14,14,15	0.37	0	17,19,21	1.28	1 (5%)
6	FUC	A	405	5	9,10,11	0.19	0	13,14,16	0.46	0
6	FUC	A	406	5	9,10,11	0.59	0	13,14,16	1.03	2 (15%)
5	NAG	A	407	1,5,6	14,14,15	0.29	0	17,19,21	0.56	0
5	NAG	A	408	8,5	14,14,15	0.33	0	17,19,21	0.85	0
6	FUC	A	409	5	9,10,11	0.22	0	13,14,16	0.85	0
7	MAN	A	410	8	11,11,12	0.24	0	15,15,17	0.52	0
8	BMA	A	411	5,7	11,11,12	0.21	0	15,15,17	0.75	0
5	NAG	A	412	1,5,6	14,14,15	0.33	0	17,19,21	1.10	1 (5%)
6	FUC	A	413	5	9,10,11	0.30	0	13,14,16	0.45	0
5	NAG	A	414	8,5	14,14,15	0.26	0	17,19,21	0.71	0
8	BMA	A	415	5,7	11,11,12	0.21	0	15,15,17	0.82	0
7	MAN	A	416	8	11,11,12	0.25	0	15,15,17	0.69	0
6	FUC	A	417	5	9,10,11	0.18	0	13,14,16	0.36	0
7	MAN	A	418	8	11,11,12	0.25	0	15,15,17	0.60	0
11	EDO	A	421	-	3,3,3	0.68	0	2,2,2	0.32	0
11	EDO	A	422	-	3,3,3	0.58	0	2,2,2	0.35	0
12	D3D	A	423	-	48,48,50	0.96	3 (6%)	51,54,56	1.21	3 (5%)
11	EDO	B	101	-	3,3,3	3.52	2 (66%)	2,2,2	0.78	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	CUY	A	401	-	-	0/33/33/42	0/0/0/0
5	NAG	A	403	1,5,6	-	0/6/23/26	0/1/1/1
5	NAG	A	404	5	-	0/6/23/26	0/1/1/1
6	FUC	A	405	5	-	0/0/17/20	0/1/1/1
6	FUC	A	406	5	-	0/0/17/20	0/1/1/1
5	NAG	A	407	1,5,6	-	0/6/23/26	0/1/1/1
5	NAG	A	408	8,5	-	0/6/23/26	0/1/1/1
6	FUC	A	409	5	-	0/0/17/20	0/1/1/1
7	MAN	A	410	8	-	0/2/19/22	0/1/1/1
8	BMA	A	411	5,7	-	0/2/19/22	0/1/1/1
5	NAG	A	412	1,5,6	-	0/6/23/26	0/1/1/1
6	FUC	A	413	5	-	0/0/17/20	0/1/1/1
5	NAG	A	414	8,5	-	0/6/23/26	0/1/1/1
8	BMA	A	415	5,7	-	0/2/19/22	0/1/1/1
7	MAN	A	416	8	-	0/2/19/22	0/1/1/1
6	FUC	A	417	5	-	0/0/17/20	0/1/1/1
7	MAN	A	418	8	-	0/2/19/22	0/1/1/1
11	EDO	A	421	-	-	0/1/1/1	0/0/0/0
11	EDO	A	422	-	-	0/1/1/1	0/0/0/0
12	D3D	A	423	-	-	0/53/53/55	0/0/0/0
11	EDO	B	101	-	-	0/1/1/1	0/0/0/0

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
11	B	101	EDO	C2-C1	-4.26	1.18	1.48
11	B	101	EDO	O1-C1	-4.15	1.20	1.42
12	A	423	D3D	O1-C16	-2.76	1.39	1.45
12	A	423	D3D	O1-C15	2.28	1.39	1.33
12	A	423	D3D	O8-C22	2.65	1.42	1.34
3	A	401	CUY	OAQ-CAP	2.83	1.41	1.33

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
12	A	423	D3D	C18-C17-C16	-5.10	100.35	111.86
5	A	404	NAG	O5-C1-C2	-4.78	104.92	111.52
5	A	403	NAG	O5-C1-C2	-4.62	105.14	111.52
5	A	412	NAG	O5-C1-C2	-2.91	107.50	111.52
5	A	403	NAG	O5-C5-C6	-2.74	102.82	107.15
5	A	403	NAG	C1-C2-N2	-2.64	105.98	110.49

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	406	FUC	O2-C2-C1	2.06	113.32	109.17
5	A	403	NAG	O5-C5-C4	2.17	116.11	110.83
12	A	423	D3D	O8-C22-C23	2.62	117.07	111.55
6	A	406	FUC	C1-C2-C3	2.74	113.12	109.66
3	A	401	CUY	OAQ-CAP-CAO	2.88	120.24	111.92
12	A	423	D3D	O1-C16-C17	3.05	116.25	108.64
5	A	403	NAG	C1-O5-C5	8.50	123.88	112.19

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	401	CUY	1	0
5	A	403	NAG	1	0
5	A	404	NAG	1	0
6	A	406	FUC	1	0
12	A	423	D3D	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	280/300 (93%)	0.04	8 (2%) 51 54	9, 17, 35, 61	0
2	B	98/99 (98%)	0.24	4 (4%) 37 39	10, 21, 45, 48	0
All	All	378/399 (94%)	0.09	12 (3%) 47 50	9, 17, 40, 61	0

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	152	GLN	6.0
1	A	151	TYR	4.4
2	B	1	ILE	3.8
1	A	200	GLY	2.7
1	A	195	PRO	2.6
2	B	75	LYS	2.6
1	A	199	PRO	2.5
2	B	42	ASN	2.5
2	B	73	THR	2.3
1	A	198	GLY	2.2
1	A	202	LEU	2.1
1	A	283	ASN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	FUC	A	405	10/11	0.48	0.26	60,65,66,66	0
11	EDO	A	421	4/4	0.58	0.24	39,40,40,40	0
12	D3D	A	423	49/51	0.59	0.20	29,30,34,38	0
5	NAG	A	403	14/15	0.61	0.19	42,44,48,55	0
5	NAG	A	408	14/15	0.64	0.30	39,42,45,47	0
5	NAG	A	407	14/15	0.69	0.26	39,44,50,50	0
6	FUC	A	409	10/11	0.69	0.33	45,48,50,51	0
3	CUY	A	401	35/44	0.69	0.16	31,35,38,39	0
5	NAG	A	404	14/15	0.70	0.27	36,39,41,43	0
7	MAN	A	410	11/12	0.72	0.23	33,37,40,41	0
6	FUC	A	406	10/11	0.72	0.25	41,42,46,46	0
8	BMA	A	411	11/12	0.79	0.33	37,42,47,48	0
11	EDO	A	422	4/4	0.81	0.15	41,43,43,43	0
6	FUC	A	417	10/11	0.90	0.10	15,20,24,24	0
7	MAN	A	416	11/12	0.93	0.09	12,15,18,19	0
11	EDO	B	101	4/4	0.93	0.25	3,12,25,26	0
7	MAN	A	418	11/12	0.94	0.09	15,18,20,21	0
6	FUC	A	413	10/11	0.95	0.07	15,18,21,22	0
10	NA	A	420	1/1	0.96	0.07	23,23,23,23	0
8	BMA	A	415	11/12	0.97	0.07	11,13,16,18	0
5	NAG	A	414	14/15	0.97	0.06	7,12,17,23	0
9	CL	A	419	1/1	0.98	0.08	17,17,17,17	0
5	NAG	A	412	14/15	0.98	0.05	8,12,15,16	0
4	IOD	A	402	1/1	0.99	0.22	6,6,6,6	1
10	NA	B	102	1/1	0.99	0.07	16,16,16,16	0

6.5 Other polymers [i](#)

There are no such residues in this entry.