

Aug 27, 2024 – 12:37 AM JST

PDB ID	:	8WLH
EMDB ID	:	EMD-37619
Title	:	Cryo-EM structure of the proximal rod-export apparatus and FlgF within the
		motor-hook complex in the CCW state
Authors	:	Tan, J.X.; Zhang, L.; Zhou, Y.; Zhu, Y.Q.
Deposited on	:	2023-09-29
Resolution	:	3.70 Å(reported)
Based on initial model	:	

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/EMValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

EMDB validation analysis	:	0.0.1.dev112
MolProbity	:	4.02b-467
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
MapQ	:	1.9.13
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.38.2

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $ELECTRON\ MICROSCOPY$

The reported resolution of this entry is 3.70 Å.

Sidechain outliers

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.

Metric	Percentile Ran	ks Value								
Ramachandran outliers		0.5%								
Sidechain outliers	3.0%									
Worse	Better									
Percentile relative to all structures										
Percenti	le relative to all EM structures									
Motria	Whole archive	EM structures								
Metric	$(\# { m Entries})$	$(\# {\rm Entries})$								
Ramachandran outliers	207382	16835								

206894

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion < 40%). The numeric value is given above the bar.

16415

Mol	Chain	Length	Quality of chain							
1		20	47%							
1	A	89	92%	7% •						
1	В	89	98%	•						
			15%							
1	С	89	100%							
	-		26%							
1	D	89	100%							
			28%							
2	E	264	93%	• •						
	D	2.15	14%							
3	F'	245	82%	16%						
_			6%							
3	G	245	82% •	15%						
			7%							
3	H	245	82% •	15%						
			7%							
3	I	245	82%	15%						



Chain Length Quality of chain Mol 14% J 3 24580% 15% • 7% Κ 4 10435% 62% 12% \mathbf{L} 1044 67% 31% • 8% 4 Μ 104 70% 29% • 7% Ν 1044 69% • 29% 5% Ο 1044 69% 29% . 8% Р 1044 67% 30% • 9% Q 513883% • 14% 6% R 138522% 77% . \mathbf{S} 513876% • 22% 7% Т 513878% 20% • 5% U 513873% •• 23% 16% V 1346 99% 11% • • W 6 13496% 7% ••• Х 1346 98% 6% 6 Υ 1346% • 93% 5% Ζ • • 6 13494% 7% 6 134 \mathbf{a} 5%• 94% • 7 b 56098% Ļ 7560 \mathbf{c} 97% • 7 \mathbf{d} 56096% 7 560• е 97% f 560796% • • 756097% g 7h 560• 96%

Continued from previous page...



Mol	Chain	Length	Quality of chain	
7	i	560	• 97%	ı
7	j	560	• 96%	
7	k	560	• 97%	1
7	1	560	• 96%	1
8	m	251	97% ···	1
8	n	251	9%	1
8	О	251	8% 98% •	
8	р	251	96% •	
8	q	251	98%	



2 Entry composition (i)

There are 8 unique types of molecules in this entry. The entry contains 36230 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues		At	oms			AltConf	Trace
1	Δ	80	Total	С	Ν	Ο	\mathbf{S}	0	0
1	Π	03	670	449	100	114	7	0	0
1	1 D	80	Total	С	Ν	0	S	0	0
1	D	89	670	449	100	114	$\overline{7}$	0	0
1	С	80	Total	С	Ν	0	S	0	0
	U	89	670	449	100	114	$\overline{7}$	0	
1	Л	80	Total	С	Ν	0	S	0	0
	D	89	670	449	100	114	7	0	

• Molecule 1 is a protein called Flagellar biosynthetic protein FliQ.

• Molecule 2 is a protein called Flagellar biosynthetic protein FliR.

Mol	Chain	Residues		At	AltConf	Trace			
2	Е	253	Total 1945	C 1305	N 307	0 318	S 15	0	0

• Molecule 3 is a protein called Flagellar biosynthetic protein FliP.

Mol	Chain	Residues		At	oms		AltConf	Trace	
3 F	Б	207	Total	С	Ν	0	S	0	0
	207	1605	1072	249	272	12	0	0	
2	3 C	200	Total	С	Ν	0	S	0	0
3 G	G	209	1626	1086	252	276	12	0	0
3	и	208	Total	С	Ν	0	S	0	0
0	11		1614	1077	251	274	12	0	0
2	т	208	Total	С	Ν	0	S	0	0
0	1	208	1614	1077	251	274	12	0	0
3	т	209	Total	С	Ν	0	S	0	0
	J		1623	1084	251	276	12	0	

• Molecule 4 is a protein called Flagellar hook-basal body complex protein FliE.



Mol	Chain	Residues		At	oms			AltConf	Trace
4	Κ	40	Total 300	C 185	N 52	O 57	S 6	0	0
4	L	72	Total 543	C 335	N 99	0 103	S 6	0	0
4	М	74	Total 557	C 344	N 101	0 106	S 6	0	0
4	Ν	74	Total 557	C 344	N 101	0 106	S 6	0	0
4	О	74	Total 557	C 344	N 101	0 106	S 6	0	0
4	Р	73	Total 550	C 340	N 100	0 104	S 6	0	0

• Molecule 5 is a protein called Flagellar basal body rod protein FlgB.

Mol	Chain	Residues		At	oms		AltConf	Trace	
5 Q	0	110	Total	С	Ν	0	S	0	0
	Q	115	922	565	169	183	5	0	
5	5 B	108	Total	С	Ν	Ο	S	0	0
0	п	108	848	523	155	165	5	0	0
5	q	108	Total	С	Ν	Ο	S	0	0
0	G		848	523	155	165	5	0	0
5	Т	110	Total	С	Ν	0	S	0	0
	1	110	863	531	160	167	5	0	0
F	U	106	Total	С	Ν	0	S	0	0
0		100	832	514	150	163	5	U	0

• Molecule 6 is a protein called Flagellar basal-body rod protein FlgC.

Mol	Chain	Residues		At	oms		AltConf	Trace	
6	V	122	Total	С	Ν	0	S	0	0
0	v	100	969	604	167	193	5	0	0
6	W	139	Total	С	Ν	Ο	S	0	0
0	vv	152	964	601	166	192	5	0	0
6	6 X	122	Total	С	Ν	Ο	S	0	0
0		100	969	604	167	193	5		
6	v	122	Total	С	Ν	Ο	S	0	0
0	1	155	969	604	167	193	5		0
6	7	121	Total	С	Ν	Ο	S	0	0
0		101	956	595	165	191	5	0	0
6	0	199	Total	С	Ν	0	S	0	0
0	a	100	969	604	167	193	5		0

R L D W I D E PDB TEIN DATA BANK

• Molecule 7 is a protein called Flagellar M-ring protein.

Mol	Chain	Residues		Ator	\mathbf{ns}		AltConf	Trace
7	h	19	Total	С	Ν	0	0	0
(D	15	81	50	15	16	0	0
7		16	Total	С	Ν	0	0	0
	C	10	103	64	19	20	0	0
7	d	20	Total	С	Ν	0	0	0
	u	20	133	83	23	27	0	0
7	0	16	Total	С	Ν	0	0	0
1		10	103	64	19	20	0	0
7	7 f	91	Total	С	Ν	0	0	0
1		21	140	88	24	28	0	0
7	G	16	Total	С	Ν	0	0	0
1	g	10	103	64	19	20	0	0
7	h	91	Total	С	Ν	0	0	0
1	11	21	140	88	24	28	0	0
7	i	16	Total	С	Ν	0	0	0
1	1	10	103	64	19	20	0	0
7	i	20	Total	С	Ν	Ο	0	0
1	J	20	133	83	23	27	0	0
7	Ŀ	16	Total	С	Ν	0	0	0
I K	10	103	64	19	20		0	
7 1	1	91	Total	С	Ν	0	0	0
'	1		140	88	24	28		

• Molecule 8 is a protein called Flagellar basal-body rod protein FlgF.

Mol	Chain	Residues		At		AltConf	Trace		
8	m	248	Total	С	Ν	Ο	S	0	0
0	111		1804	1106	324	367	7	0	U
8	n	240	Total	С	Ν	Ο	\mathbf{S}	0	0
0	11	249	1812	1111	325	368	8	0	0
8	0	o 250	Total	С	Ν	Ο	\mathbf{S}	0	0
0	0		1820	1116	326	369	9	0	0
8	n	250	Total	С	Ν	Ο	\mathbf{S}	0	0
0	o p	230	1820	1116	326	369	9	0	0
8	8 q	~ <u>240</u>	Total	С	Ν	0	S	0	0
8		249	1812	1111	325	368	8	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Flagellar biosynthetic protein FliQ









0160 8163 A173 A164 A173 A164 A173 A164 A173 A164 A173 A173

• Molecule 3: Flagellar biosynthetic protein FliP



E118 G119 G119 A126 A131 A135 A145 A145 A145 A145 A145 A145 A151 A152 A154 A155 A156 A151 A153 A154 A155 A154 A154 A153 A154 A154 A154 A154 A154 A154 A154 A154 A154 A215 <t

• Molecule 4: Flagellar hook-basal body complex protein FliE

	7%							_
Chain K:		35%		•		62%		
MET ALA ALA ALA GLN GLN ILE ILE	GLY GLY VAL ILE SER	GLN LEU GLN ALA THR	ALA MET ALA ALA ARG GLY GLN ASP	THR HIS SER GLN SER SER	VAL SER PHE ALA GLY GLN LEU HIS	ALA ALA LEU ASP ASP ILE SER SER	ARG GLN ALA ALA ALA ALA ALA	VAL GLN ALA GLU LYS PHE THR



• Molecule 4: Flagellar hook-basal body complex protein FliE

Chain L: 67% · 31%

• Molecule 4: Flagellar hook-basal body complex protein FliE

Molecule 4: Flagellar hook-basal body complex protein FliE
 7%
 Chain N:
 69%
 29%
 Extended to the second s















A ASN VALL VALL PPHEE PP

ILLE ARG GLN TRP MET MET SER ASN ASN ASN HIS GLU

• Molecule 7: Flagellar M-ring protein



• Molecule 7: Flagellar M-ring protein

 Chain e:
 97%

 State
 97%

 State
 84%

 State
 84% 84%

 State
 84% 84% 84%

 State
 84% 84% 84% 84%

 State
 84%</th





Chain g:

97%







ASN ASP HIS GLU

• Molecule 7: Flagellar M-ring protein













• Molecule 8: Flagellar basal-body rod protein FlgF



• Molecule 8: Flagellar basal-body rod protein FlgF







4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	11858	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE	Depositor
	CORRECTION	
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{\AA}^2)$	45	Depositor
Minimum defocus (nm)	1000	Depositor
Maximum defocus (nm)	2500	Depositor
Magnification	105000	Depositor
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	2.735	Depositor
Minimum map value	-1.731	Depositor
Average map value	-0.006	Depositor
Map value standard deviation	0.133	Depositor
Recommended contour level	0.65	Depositor
Map size (Å)	681.984, 681.984, 681.984	wwPDB
Map dimensions	512, 512, 512	wwPDB
Map angles ($^{\circ}$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.332, 1.332, 1.332	Depositor



5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond lengths		Bond angles		
	Unam	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	0.45	0/681	0.67	1/930~(0.1%)	
1	В	0.32	0/681	0.52	0/930	
1	С	0.28	0/681	0.50	0/930	
1	D	0.26	0/681	0.48	0/930	
2	Е	0.31	0/1994	0.52	0/2724	
3	F	0.36	0/1643	0.62	2/2237~(0.1%)	
3	G	0.30	0/1665	0.49	1/2267~(0.0%)	
3	Н	0.29	0/1652	0.49	0/2249	
3	Ι	0.29	0/1652	0.47	0/2249	
3	J	0.33	0/1662	0.52	0/2263	
4	Κ	0.28	0/300	0.52	0/400	
4	L	0.25	0/547	0.44	0/733	
4	М	0.26	0/561	0.44	0/753	
4	Ν	0.25	0/561	0.46	0/753	
4	0	0.27	0/561	0.49	0/753	
4	Р	0.31	0/554	0.48	0/743	
5	Q	0.29	0/930	0.56	0/1251	
5	R	0.26	0/855	0.49	0/1150	
5	S	0.28	0/855	0.53	0/1150	
5	Т	0.26	0/870	0.49	0/1169	
5	U	0.26	0/839	0.47	0/1129	
6	V	0.30	0/981	0.47	0/1334	
6	W	0.29	0/976	0.50	0/1327	
6	Х	0.34	0/981	0.50	0/1334	
6	Y	0.33	0/981	0.62	0/1334	
6	Ζ	0.29	0/968	0.48	0/1316	
6	a	0.36	0/981	0.52	0/1334	
7	b	0.46	0/83	0.86	1/114~(0.9%)	
7	с	0.27	0/107	0.38	0/148	
7	d	0.31	0/137	0.49	0/191	
7	е	0.28	0/107	0.56	0/148	
7	f	0.39	0/145	0.55	0/203	
7	g	0.33	0/107	0.51	0/148	
7	h	0.26	0/145	0.43	0/203	



Mal	Chain	Bond	lengths	Bond angles		
	Ullaill	RMSZ	# Z > 5	RMSZ	# Z > 5	
7	i	0.29	0/107	0.38	0/148	
7	j	0.33	0/137	0.70	0/191	
7	k	0.30	0/107	0.37	0/148	
7	l	0.29	0/145	0.45	0/203	
8	m	0.32	0/1828	0.56	0/2492	
8	n	0.33	0/1836	0.56	2/2502~(0.1%)	
8	0	0.37	0/1844	0.57	0/2512	
8	р	0.33	0/1844	0.57	0/2512	
8	q	0.33	0/1836	0.58	0/2502	
All	All	0.31	0/36808	0.53	7/50037~(0.0%)	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	Н	0	2
3	J	0	2
5	Q	0	1
5	R	0	1
5	Т	0	1
5	U	0	1
6	Y	0	1
6	a	0	1
8	m	0	1
8	n	0	2
8	р	0	1
All	All	0	14

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms		$Observed(^{o})$	$Ideal(^{o})$
3	F	162	PRO	CA-N-CD	-11.98	94.72	111.50
3	F	162	PRO	N-CD-CG	-6.57	93.35	103.20
1	А	85	PRO	CA-N-CD	-5.88	103.27	111.50
8	n	53	THR	CB-CA-C	-5.62	96.43	111.60
3	G	162	PRO	N-CA-CB	-5.30	96.77	102.60
8	n	72	ASP	CB-CG-OD1	5.25	123.03	118.30
7	b	322	PRO	N-CA-CB	-5.19	96.89	102.60



There are no chirality outliers.

All (14) planarity of	outliers	are listed	below:
-----------------------	----------	------------	--------

Mol	Chain	Res	Type	Group
3	Н	135	ARG	Sidechain
3	Н	143	ARG	Sidechain
3	J	140	ARG	Sidechain
3	J	143	ARG	Sidechain
5	Q	106	ARG	Sidechain
5	R	55	ARG	Sidechain
5	Т	4	ARG	Sidechain
5	U	104	ARG	Sidechain
6	Y	110	ARG	Sidechain
6	а	110	ARG	Sidechain
8	m	243	ARG	Sidechain
8	n	225	ARG	Sidechain
8	n	54	ARG	Sidechain
8	р	200	ARG	Sidechain

5.2 Too-close contacts (i)

Due to software issues we are unable to calculate clashes - this section is therefore empty.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentile	es
1	А	87/89~(98%)	81~(93%)	4(5%)	2(2%)	5 31	
1	В	87/89~(98%)	86~(99%)	1 (1%)	0	100 100)
1	С	87/89~(98%)	86~(99%)	1 (1%)	0	100 100)
1	D	87/89~(98%)	85~(98%)	2(2%)	0	100 100)
2	Е	251/264~(95%)	234~(93%)	15 (6%)	2(1%)	16 49	
3	F	205/245~(84%)	197 (96%)	8 (4%)	0	100 100)



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
3	G	207/245~(84%)	199 (96%)	6 (3%)	2 (1%)	13	44
3	Н	206/245~(84%)	201 (98%)	5 (2%)	0	100	100
3	Ι	206/245~(84%)	199 (97%)	6 (3%)	1 (0%)	25	57
3	J	207/245~(84%)	192 (93%)	10 (5%)	5 (2%)	5	30
4	Κ	38/104~(36%)	35~(92%)	3~(8%)	0	100	100
4	L	70/104~(67%)	70 (100%)	0	0	100	100
4	М	72/104~(69%)	70~(97%)	2(3%)	0	100	100
4	Ν	72/104~(69%)	72 (100%)	0	0	100	100
4	Ο	72/104~(69%)	72 (100%)	0	0	100	100
4	Р	71/104~(68%)	71 (100%)	0	0	100	100
5	Q	115/138~(83%)	114 (99%)	1 (1%)	0	100	100
5	R	104/138~(75%)	103 (99%)	1 (1%)	0	100	100
5	S	104/138~(75%)	103 (99%)	1 (1%)	0	100	100
5	Т	106/138~(77%)	105 (99%)	1 (1%)	0	100	100
5	U	102/138~(74%)	101 (99%)	1 (1%)	0	100	100
6	V	131/134~(98%)	122 (93%)	9 (7%)	0	100	100
6	W	130/134~(97%)	123 (95%)	7 (5%)	0	100	100
6	Х	131/134~(98%)	124 (95%)	7 (5%)	0	100	100
6	Y	131/134~(98%)	121 (92%)	7 (5%)	3 (2%)	5	31
6	Ζ	129/134~(96%)	123 (95%)	5 (4%)	1 (1%)	16	49
6	a	131/134~(98%)	124 (95%)	7 (5%)	0	100	100
7	b	11/560~(2%)	9 (82%)	2 (18%)	0	100	100
7	с	14/560~(2%)	12 (86%)	2 (14%)	0	100	100
7	d	18/560~(3%)	18 (100%)	0	0	100	100
7	е	14/560~(2%)	14 (100%)	0	0	100	100
7	f	19/560~(3%)	18 (95%)	1 (5%)	0	100	100
7	g	14/560~(2%)	13 (93%)	1 (7%)	0	100	100
7	h	19/560~(3%)	19 (100%)	0	0	100	100
7	i	14/560~(2%)	14 (100%)	0	0	100	100
7	j	18/560~(3%)	18 (100%)	0	0	100	100
7	k	14/560~(2%)	14 (100%)	0	0	100	100



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
7	1	19/560~(3%)	19 (100%)	0	0	100	100
8	m	246/251~(98%)	238~(97%)	7(3%)	1 (0%)	30	62
8	n	247/251~(98%)	242 (98%)	4 (2%)	1 (0%)	30	62
8	О	248/251~(99%)	233~(94%)	13~(5%)	2(1%)	16	49
8	р	248/251~(99%)	235~(95%)	13~(5%)	0	100	100
8	q	247/251~(98%)	236 (96%)	9 (4%)	2 (1%)	16	49
All	All	4749/11378 (42%)	4565 (96%)	162 (3%)	22 (0%)	27	57

All (22) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	А	84	LEU
3	J	83	ALA
3	J	87	GLN
8	q	155	VAL
2	Е	167	ASN
2	Е	213	ILE
3	G	83	ALA
3	Ι	160	GLN
3	J	86	ASN
6	Y	54	ALA
8	0	50	SER
8	q	53	THR
3	J	140	ARG
6	Y	58	ALA
6	Y	55	PRO
1	А	85	PRO
3	G	162	PRO
8	n	52	ALA
8	0	52	ALA
6	Ζ	58	ALA
8	m	52	ALA
3	J	229	VAL

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	74/74~(100%)	68~(92%)	6 (8%)	9 34
1	В	74/74~(100%)	72~(97%)	2(3%)	40 61
1	\mathbf{C}	74/74~(100%)	74 (100%)	0	100 100
1	D	74/74~(100%)	74 (100%)	0	100 100
2	Е	210/221~(95%)	205~(98%)	5(2%)	44 63
3	F	177/204 (87%)	171 (97%)	6 (3%)	32 56
3	G	179/204~(88%)	172 (96%)	7 (4%)	27 53
3	Н	178/204~(87%)	173 (97%)	5(3%)	38 60
3	Ι	178/204 (87%)	171 (96%)	7 (4%)	27 53
3	J	179/204~(88%)	172 (96%)	7 (4%)	27 53
4	К	33/79~(42%)	29~(88%)	4 (12%)	4 20
4	L	56/79~(71%)	54 (96%)	2(4%)	30 55
4	М	58/79~(73%)	57~(98%)	1 (2%)	56 73
4	Ν	58/79~(73%)	56 (97%)	2(3%)	32 56
4	О	58/79~(73%)	56~(97%)	2(3%)	32 56
4	Р	57/79~(72%)	54 (95%)	3~(5%)	19 46
5	Q	98/113~(87%)	94 (96%)	4 (4%)	26 52
5	R	90/113~(80%)	89~(99%)	1 (1%)	70 80
5	S	90/113~(80%)	87 (97%)	3(3%)	33 57
5	Т	91/113~(80%)	90~(99%)	1 (1%)	70 80
5	U	89/113~(79%)	84 (94%)	5~(6%)	17 45
6	V	104/105~(99%)	104 (100%)	0	100 100
6	W	104/105~(99%)	101 (97%)	3(3%)	37 59
6	Х	104/105~(99%)	102 (98%)	2(2%)	52 70
6	Y	104/105~(99%)	100 (96%)	4 (4%)	28 53
6	Ζ	103/105~(98%)	99~(96%)	4 (4%)	27 53
6	a	104/105~(99%)	98 (94%)	6~(6%)	17 44
7	b	8/467~(2%)	6~(75%)	2(25%)	0 4
7	с	11/467~(2%)	10 (91%)	1 (9%)	7 30
7	d	14/467~(3%)	12 (86%)	2 (14%)	2 16

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Rotameric	Outliers	Perce	entiles
7	е	11/467~(2%)	11 (100%)	0	100	100
7	f	15/467~(3%)	15 (100%)	0	100	100
7	g	11/467~(2%)	11 (100%)	0	100	100
7	h	15/467~(3%)	15 (100%)	0	100	100
7	i	11/467~(2%)	11 (100%)	0	100	100
7	j	14/467~(3%)	13 (93%)	1 (7%)	12	39
7	k	11/467~(2%)	11 (100%)	0	100	100
7	1	15/467~(3%)	14 (93%)	1 (7%)	13	40
8	m	190/193~(98%)	188 (99%)	2 (1%)	70	80
8	n	191/193 (99%)	187 (98%)	4 (2%)	48	67
8	О	192/193~(100%)	189 (98%)	3 (2%)	58	74
8	р	192/193~(100%)	185 (96%)	7 (4%)	30	55
8	q	191/193 (99%)	189 (99%)	2 (1%)	73	82
All	All	3890/9308 (42%)	3773 (97%)	117 (3%)	37	58

All (117) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	7	MET
1	А	69	MET
1	А	73	LEU
1	А	75	ASP
1	А	85	PRO
1	А	87	ILE
1	В	2	THR
1	В	47	MET
2	Е	36	ARG
2	Е	86	PHE
2	Е	214	PHE
2	Е	227	MET
2	Е	229	MET
3	F	38	TRP
3	F	129	LYS
3	F	155	ASN
3	F	160	GLN
3	F	190	PHE
3	F	213	PRO



Mol	Chain	Res	Type
3	G	75	ARG
3	G	102	SER
3	G	116	PHE
3	G	140	ARG
3	G	160	GLN
3	G	162	PRO
3	G	213	PRO
3	Н	135	ARG
3	Н	152	ARG
3	Н	177	SER
3	Н	187	PHE
3	Н	190	PHE
3	Ι	113	TYR
3	Ι	143	ARG
3	Ι	152	ARG
3	Ι	187	PHE
3	Ι	205	MET
3	Ι	238	SER
3	Ι	241	GLN
3	J	65	THR
3	J	117	SER
3	J	140	ARG
3	J	144	GLU
3	J	190	PHE
3	J	224	MET
3	J	230	ASP
4	Κ	74	ASP
4	K	77	LYS
4	Κ	79	SER
4	Κ	101	SER
4	L	75	MET
4	L	84	MET
4	М	70	ASP
4	Ν	33	SER
4	N	81	SER
4	0	101	SER
4	0	102	MET
4	Р	57	GLU
4	Р	91	LYS
4	Р	102	MET
5	Q	34	ASP
5	Q	101	ASP



Mol	Chain	Res	Type
5	Q	113	SER
5	Q	125	SER
5	R	47	SER
5	S	48	GLU
5	S	91	ASP
5	S	128	LYS
5	Т	47	SER
5	U	4	ARG
5	U	18	LEU
5	U	104	ARG
5	U	113	SER
5	U	125	SER
6	W	46	GLN
6	W	123	LYS
6	W	126	MET
6	Х	119	LEU
6	Х	128	LYS
6	Y	3	LEU
6	Y	4	LEU
6	Y	30	ASN
6	Y	120	ASN
6	Ζ	20	ARG
6	Ζ	52	ASP
6	Ζ	84	LEU
6	Ζ	94	PRO
6	a	38	ASP
6	a	86	ASP
6	a	88	ASN
6	a	123	LYS
6	a	125	MET
6	a	127	LEU
7	b	317	SER
7	b	322	PRO
7	с	324	ASN
7	d	318	ASN
7	d	319	GLN
7	j	319	GLN
7	1	318	ASN
8	m	123	GLU
8	m	247	LEU
8	n	53	THR
8	n	72	ASP



Mol	Chain	Res	Type
8	n	153	ASN
8	n	225	ARG
8	0	189	ARG
8	0	200	ARG
8	0	225	ARG
8	р	47	ASP
8	р	49	LEU
8	р	72	ASP
8	р	140	ASP
8	р	160	ARG
8	р	169	ASN
8	р	189	ARG
8	q	51	LEU
8	q	175	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (71) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	39	GLN
2	Е	162	ASN
2	Е	182	ASN
2	Е	205	ASN
3	F	119	GLN
3	F	141	GLN
3	Н	36	GLN
3	Н	132	GLN
3	Н	233	GLN
3	Ι	160	GLN
3	Ι	184	GLN
3	J	114	GLN
3	J	155	ASN
4	Κ	69	ASN
4	Κ	83	GLN
4	Κ	103	GLN
4	L	87	GLN
4	L	90	ASN
4	М	103	GLN
4	N	87	GLN
4	0	55	GLN
5	Q	17	ASN
5	R	23	GLN
5	R	29	ASN



Mol	Chain	Res	Type
5	R	92	GLN
5	R	117	GLN
5	S	32	ASN
5	S	92	GLN
5	Т	17	ASN
5	Т	21	GLN
5	Т	32	ASN
5	Т	117	GLN
5	U	112	ASN
6	V	30	ASN
6	W	5	ASN
6	W	22	ASN
6	W	120	ASN
6	Х	22	ASN
6	Х	50	GLN
6	Х	57	GLN
6	Х	134	GLN
6	Y	22	ASN
6	Ζ	17	GLN
6	Ζ	30	ASN
6	a	57	GLN
6	a	115	ASN
7	b	318	ASN
7	b	319	GLN
7	е	324	ASN
7	1	324	ASN
8	m	102	ASN
8	n	18	GLN
8	n	210	ASN
8	n	223	ASN
8	n	240	ASN
8	0	14	GLN
8	0	18	GLN
8	0	28	ASN
8	0	84	GLN
8	0	116	GLN
8	0	186	GLN
8	0	230	GLN
8	р	28	ASN
8	р	210	ASN
8	р	240	ASN
8	q	18	GLN



Continued from previous page...

Mol	Chain	Res	Type
8	q	112	GLN
8	q	116	GLN
8	q	118	HIS
8	q	153	ASN
8	q	240	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-37619. These allow visual inspection of the internal detail of the map and identification of artifacts.

Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections (i)

6.1.1 Primary map



6.1.2 Raw map



The images above show the map projected in three orthogonal directions.



6.2 Central slices (i)

6.2.1 Primary map



X Index: 256





Z Index: 256

6.2.2 Raw map



X Index: 256

Y Index: 256



The images above show central slices of the map in three orthogonal directions.



6.3 Largest variance slices (i)

6.3.1 Primary map



X Index: 247





Z Index: 249

6.3.2 Raw map



X Index: 0

Y Index: 0



The images above show the largest variance slices of the map in three orthogonal directions.



6.4 Orthogonal standard-deviation projections (False-color) (i)

6.4.1 Primary map



6.4.2 Raw map



The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.



6.5 Orthogonal surface views (i)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.65. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.

6.6 Mask visualisation (i)

This section was not generated. No masks/segmentation were deposited.



7 Map analysis (i)

This section contains the results of statistical analysis of the map.

7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



7.2 Volume estimate (i)



The volume at the recommended contour level is 338 $\rm nm^3;$ this corresponds to an approximate mass of 306 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



7.3 Rotationally averaged power spectrum (i)



*Reported resolution corresponds to spatial frequency of 0.270 \AA^{-1}



8 Fourier-Shell correlation (i)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC (i)



*Reported resolution corresponds to spatial frequency of 0.270 ${\rm \AA^{-1}}$



8.2 Resolution estimates (i)

$\mathbf{Bosolution} \text{ ostimato } (\mathbf{\hat{\lambda}})$	Estimation criterion (FSC cut-off)		
Resolution estimate (A)	0.143	0.5	Half-bit
Reported by author	3.70	-	-
Author-provided FSC curve	3.75	4.40	3.82
Unmasked-calculated*	16.42	28.49	19.57

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 16.42 differs from the reported value 3.7 by more than 10 %



9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-37619 and PDB model 8WLH. Per-residue inclusion information can be found in section 3 on page 8.

9.1 Map-model overlay (i)



The images above show the 3D surface view of the map at the recommended contour level 0.65 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.



9.2 Q-score mapped to coordinate model (i)



The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model (i)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.65).



9.4 Atom inclusion (i)



At the recommended contour level, 80% of all backbone atoms, 60% of all non-hydrogen atoms, are inside the map.



1.0

0.0 <0.0

9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (0.65) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	$\mathbf{Q} extsf{-score}$
All	0.6030	0.3710
А	0.4210	0.2680
В	0.5440	0.3100
С	0.5850	0.3080
D	0.5170	0.2670
Е	0.5120	0.3040
F	0.5710	0.3300
G	0.6190	0.3650
Н	0.6260	0.3760
Ι	0.6250	0.3740
J	0.5800	0.3500
K	0.5620	0.3420
L	0.5960	0.3620
Μ	0.6300	0.3800
Ν	0.6400	0.3850
О	0.6280	0.3710
Р	0.6170	0.3620
Q	0.6130	0.3730
R	0.6780	0.4050
S	0.6650	0.3980
Т	0.6650	0.3980
U	0.6370	0.3950
V	0.5750	0.3790
W	0.6340	0.4040
Х	0.6530	0.4190
Y	0.6540	0.4190
Z	0.6500	0.4090
a	0.6340	0.3820
b	0.5060	0.3730
с	0.5050	0.3240
d	0.4590	0.3070
е	0.6410	0.3840
f	0.6210	0.3700
g	0.5730	0.3360
h	0.6570	0.4110



Chain	Atom inclusion	Q-score
i	0.6020	0.3540
j	0.6470	0.4080
k	0.5730	0.2700
1	0.5500	0.3330
m	0.5930	0.3850
n	0.6400	0.4010
О	0.6340	0.4110
р	0.6100	0.3930
q	0.5360	0.3740

