



wwPDB EM Validation Summary Report ⓘ

May 18, 2024 – 10:01 AM EDT

PDB ID : 6WOV
EMDB ID : EMD-21862
Title : Cryo-EM structure of recombinant mouse Ryanodine Receptor type 2 wild type in complex with FKBP12.6
Authors : Iyer, K.A.; Hu, Y.; Nayak, A.R.; Kurebayashi, N.; Murayama, T.; Samsó, M.
Deposited on : 2020-04-25
Resolution : 5.10 Å (reported)
Based on initial model : 5L1D

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev92
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 5.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	4966	
1	B	4966	
1	C	4966	
1	D	4966	
2	E	107	
2	F	107	
2	G	107	
2	H	107	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 245656 atoms, of which 120412 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ryanodine receptor 2.

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	H	N	O	S		
1	A	3921	59769	19350	29277	5253	5698	191	0	0
1	B	3921	59772	19350	29280	5253	5698	191	0	0
1	C	3921	59773	19350	29281	5253	5698	191	0	0
1	D	3921	59770	19350	29278	5253	5698	191	0	0

- Molecule 2 is a protein called Peptidyl-prolyl cis-trans isomerase FKBP1B.

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	H	N	O	S		
2	E	107	1642	516	824	144	154	4	0	0
2	F	107	1642	516	824	144	154	4	0	0
2	G	107	1642	516	824	144	154	4	0	0
2	H	107	1642	516	824	144	154	4	0	0


- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		AltConf
3	A	1	Total	Zn	0
			1	1	
3	B	1	Total	Zn	0
			1	1	
3	C	1	Total	Zn	0
			1	1	
3	D	1	Total	Zn	0
			1	1	


3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Ryanodine receptor 2












• Molecule 1: Ryanodine receptor 2








I3070	ARG	I3306	A3467	VAL	M3697	S3800	Q3924	I4151	ARG	LYS	GLU
SER	SER	K3340	A3468	GLU	I3628	V3801	Q3924	M4159	ILE	GLY	GLU
ALA	ALA	G3076	L3469	ARG	E3629	L3802	M3940	W4176	VAL	THR	GLU
LEU	LEU	E3341	K3470	VAL	E3633	D3803	V3944	Q4163	THR	SER	LEU
GLY	GLY	E3342	R3471	L3551	H3633	K3812	V3944	Q4176	LEU	LEU	LEU
CYS	CYS	R3943	L3472	V3556	I3634	ALA	S3958	M4177	THR	THR	LEU
LEU	LEU	R3944	L3473	V3566	D3642	GLY	S3978	E4178	VAL	GLY	LEU
ALA	ALA	M3347	L3474	L3560	D3642	LEU	V3978	E4178	VAL	GLY	GLY
ALA	PHE	L3353	P3474	V3566	A3648	GLY	L3981	G4179	VAL	GLY	GLY
PHE	ALA	I3354	I3475	T3360	E3649	GLY	L3981	G4180	VAL	SER	SER
GLY	ALA	I3355	G3476	L3353	L3650	MET	M4018	E4181	VAL	THR	ARG
GLY	ALA	L3355	L3477	T3360	P3651	VAL	PHE	K4182	VAL	SER	VAL
LEU	PHE	L3360	N3478	T3360	E3652	THR	ASP	K4182	GLU	LEU	GLY
SER	ILE	T3360	I3479	T3360	E3652	GLY	L4025	M4184	GLY	LEU	ALA
SER	ILE	K3385	P3482	T3360	D3654	GLY	T4026	M4185	LEU	LEU	LYS
PRO	ALA	P3389	G3483	K3385	E3655	GLY	L4026	V4189	LEU	LEU	LYS
ALA	ALA	GLU	A3489	E3655	A3656	GLY	T4030	I4205	LEU	GLN	VAL
ALA	ALA	ALA	N3493	E3656	M3657	GLY	T4030	I4205	GLY	MET	VAL
GLY	GLN	VAL	I3508	A3657	K3658	VAL	M4051	I4205	GLY	GLY	ALA
GLY	HIS	PHE	ARG	M3658	R3659	LEU	A4069	I4205	LEU	GLY	LEU
LEU	HIS	PHE	ARG	K3659	K3678	GLN	A4069	I4205	LEU	GLY	LEU
LEU	LEU	ARG	SER	R3678	G3679	GLN	A4069	I4205	LEU	GLY	LEU
LEU	GLY	ARG	LYS	C3679	F3685	GLY	A4069	I4205	LEU	GLY	LEU
LEU	VAL	VAL	ASN	F3685	G3688	GLY	A4069	I4205	LEU	GLY	LEU
ASN	VAL	ASP	ALA	HIS	ASP	ASP	A4069	I4205	LEU	GLY	LEU
CYS	ASP	R3407	ALA	ASP	GLY	GLY	A4069	I4205	LEU	GLY	LEU
TYR	ASP	S3408	ALA	GLY	GLY	GLY	A4069	I4205	LEU	GLY	LEU
ARG	ASP	E3423	ALA	GLY	GLY	GLY	A4069	I4205	LEU	GLY	LEU
ILE	ASP	I3424	ALA	GLY	GLY	GLY	A4069	I4205	LEU	GLY	LEU
LEU	ASP	M3427	ALA	GLY	GLY	GLY	A4069	I4205	LEU	GLY	LEU
THR	ASP	S3428	ALA	GLY	GLY	GLY	A4069	I4205	LEU	GLY	LEU
SER	ASP	F3429	ALA	GLY	GLY	GLY	A4069	I4205	LEU	GLY	LEU
LEU	ASP	L3430	ALA	GLY	GLY	GLY	A4069	I4205	LEU	GLY	LEU
LEU	ASP	I3431	ALA	GLY	GLY	GLY	A4069	I4205	LEU	GLY	LEU
LYS	ASP	T3432	ALA	GLY	GLY	GLY	A4069	I4205	LEU	GLY	LEU
LYS	ASP	D3433	ALA	GLY	GLY	GLY	A4069	I4205	LEU	GLY	LEU
LYS	ASP	T3434	ALA	GLY	GLY	GLY	A4069	I4205	LEU	GLY	LEU
SER	ASP	S3436	ALA	GLY	GLY	GLY	A4069	I4205	LEU	GLY	LEU
LEU	ASP	R3437	ALA	GLY	GLY	GLY	A4069	I4205	LEU	GLY	LEU
ALA	ASP	M3438	ALA	GLY	GLY	GLY	A4069	I4205	LEU	GLY	LEU
VAL	ASP	S3439	ALA	GLY	GLY	GLY	A4069	I4205	LEU	GLY	LEU
GLU	ASP	R3453	ALA	GLY	GLY	GLY	A4069	I4205	LEU	GLY	LEU
ARG	ASP	M3460	ALA	GLY	GLY	GLY	A4069	I4205	LEU	GLY	LEU
GLN	ASP		ALA	GLY	GLY	GLY	A4069	I4205	LEU	GLY	LEU




• Molecule 1: Ryanodine receptor 2














• Molecule 1: Ryanodine receptor 2

















- Molecule 2: Peptidyl-prolyl cis-trans isomerase FKBP1B




- Molecule 2: Peptidyl-prolyl cis-trans isomerase FKBP1B



- Molecule 2: Peptidyl-prolyl cis-trans isomerase FKBP1B



- Molecule 2: Peptidyl-prolyl cis-trans isomerase FKBP1B



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C4	Depositor
Number of particles used	103845	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	60	Depositor
Minimum defocus (nm)	2000	Depositor
Maximum defocus (nm)	4500	Depositor
Magnification	105000	Depositor
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	24.070	Depositor
Minimum map value	-7.125	Depositor
Average map value	-0.000	Depositor
Map value standard deviation	1.000	Depositor
Recommended contour level	3.0	Depositor
Map size (\AA)	484.70398, 484.70398, 484.70398	wwPDB
Map dimensions	352, 352, 352	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (\AA)	1.377, 1.377, 1.377	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:
ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.35	0/31108	0.68	15/42056 (0.0%)
1	B	0.35	0/31108	0.68	14/42056 (0.0%)
1	C	0.35	0/31108	0.68	13/42056 (0.0%)
1	D	0.35	0/31108	0.68	14/42056 (0.0%)
2	E	0.35	0/834	0.67	0/1123
2	F	0.35	0/834	0.67	0/1123
2	G	0.35	0/834	0.67	0/1123
2	H	0.35	0/834	0.67	0/1123
All	All	0.35	0/127768	0.68	56/172716 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	15
1	B	0	15
1	C	0	15
1	D	0	15
All	All	0	60

There are no bond length outliers.

The worst 5 of 56 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	2660	LEU	CA-CB-CG	9.27	136.63	115.30
1	D	2660	LEU	CA-CB-CG	9.27	136.63	115.30
1	A	2660	LEU	CA-CB-CG	9.27	136.62	115.30
1	B	2660	LEU	CA-CB-CG	9.25	136.57	115.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	1500	ARG	N-CA-C	7.46	131.14	111.00

There are no chirality outliers.

5 of 60 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	1266	GLU	Peptide
1	A	1500	ARG	Peptide
1	A	544	ASN	Peptide
1	A	608	HIS	Peptide
1	A	618	CYS	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	30492	29277	29270	263	0
1	B	30492	29280	29270	272	0
1	C	30492	29281	29270	269	0
1	D	30492	29278	29270	276	0
2	E	818	824	824	14	0
2	F	818	824	824	14	0
2	G	818	824	824	13	0
2	H	818	824	824	14	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
All	All	125244	120412	120376	1129	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 1129 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:644:LEU:HD11	1:B:1658:THR:HG21	1.70	0.74
1:C:644:LEU:HD11	1:C:1658:THR:HG21	1.69	0.73
1:A:644:LEU:HD11	1:A:1658:THR:HG21	1.69	0.73
1:C:2106:THR:OG1	1:C:3612:ARG:O	2.06	0.73
1:B:288:HIS:O	1:B:290:ARG:NH1	2.22	0.72

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	3849/4966 (78%)	3357 (87%)	482 (12%)	10 (0%)	41	76
1	B	3849/4966 (78%)	3358 (87%)	481 (12%)	10 (0%)	41	76
1	C	3849/4966 (78%)	3358 (87%)	481 (12%)	10 (0%)	41	76
1	D	3849/4966 (78%)	3359 (87%)	480 (12%)	10 (0%)	41	76
2	E	105/107 (98%)	96 (91%)	9 (9%)	0	100	100
2	F	105/107 (98%)	96 (91%)	9 (9%)	0	100	100
2	G	105/107 (98%)	96 (91%)	9 (9%)	0	100	100
2	H	105/107 (98%)	96 (91%)	9 (9%)	0	100	100
All	All	15816/20292 (78%)	13816 (87%)	1960 (12%)	40 (0%)	44	76

5 of 40 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1500	ARG
1	B	1500	ARG
1	C	1500	ARG
1	D	1500	ARG
1	A	607	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	3178/4355 (73%)	3164 (100%)	14 (0%)	91	94
1	B	3178/4355 (73%)	3164 (100%)	14 (0%)	91	94
1	C	3178/4355 (73%)	3164 (100%)	14 (0%)	91	94
1	D	3178/4355 (73%)	3164 (100%)	14 (0%)	91	94
2	E	88/88 (100%)	87 (99%)	1 (1%)	73	85
2	F	88/88 (100%)	87 (99%)	1 (1%)	73	85
2	G	88/88 (100%)	87 (99%)	1 (1%)	73	85
2	H	88/88 (100%)	87 (99%)	1 (1%)	73	85
All	All	13064/17772 (74%)	13004 (100%)	60 (0%)	89	93

5 of 60 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	343	ARG
1	D	3654	ASP
1	C	2203	CYS
1	D	3653	GLU
2	H	49	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 66 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	2210	GLN
1	D	2317	ASN
1	D	4878	GLN
1	B	2176	ASN
1	B	2150	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-21862. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 176




Y Index: 176




Z Index: 176


6.2.2 Raw map



X Index: 352



Y Index: 352




Z Index: 352


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 191



Y Index: 191



Z Index: 159


6.3.2 Raw map



X Index: 330



Y Index: 330




Z Index: 318


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 3.0. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [\(i\)](#)



The volume at the recommended contour level is 2507 nm^3 ; this corresponds to an approximate mass of 2265 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [i](#)




*Reported resolution corresponds to spatial frequency of 0.196 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.196 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	5.10	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	4.99	7.12	5.28

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-21862 and PDB model 6WOV. Per-residue inclusion information can be found in section [3](#) on page [4](#).

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 3.0 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [\(i\)](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [\(i\)](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (3.0).



















9.4 Atom inclusion [i](#)



At the recommended contour level, 79% of all backbone atoms, 78% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (3.0) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.7800	 0.1330
A	 0.7820	 0.1330
B	 0.7810	 0.1330
C	 0.7820	 0.1330
D	 0.7820	 0.1330
E	 0.7670	 0.1520
F	 0.7720	 0.1540
G	 0.7640	 0.1470
H	 0.7640	 0.1430

