



wwPDB EM Validation Summary Report ⓘ

Dec 2, 2024 – 04:18 PM JST

PDB ID : 8WP2
EMDB ID : EMD-37708
Title : MapSPARTA tetramer bound with guide-target
Authors : Huang, P.P.; Li, Z.X.; Guo, L.J.; Xiao, Y.B.; Chen, M.R.
Deposited on : 2023-10-08
Resolution : 3.30 Å (reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev113
MolProbity : 4.02b-467
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.40

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415
RNA backbone	6643	2191

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	507	 57% 31% 10%
1	C	507	 49% 37% 11%
1	I	507	 53% 36% 9%
1	K	507	 61% 29% 9%
2	B	452	 59% 29% 8%
2	D	452	 60% 32% 8%
2	J	452	 63% 29% 7%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	L	452	
3	E	21	
3	G	21	
3	M	21	
3	O	21	
4	F	25	
4	H	25	
4	N	25	
4	P	25	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 31544 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Piwi domain-containing protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	458	Total 3682	C 2386	N 610	O 675	S 11	0	0
1	C	449	Total 3618	C 2347	N 598	O 662	S 11	0	0
1	I	460	Total 3700	C 2398	N 613	O 678	S 11	0	0
1	K	459	Total 3709	C 2406	N 611	O 681	S 11	0	0

- Molecule 2 is a protein called TIR domain-containing protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	B	417	Total 3421	C 2213	N 570	O 626	S 12	0	0
2	D	417	Total 3454	C 2234	N 574	O 634	S 12	0	0
2	J	419	Total 3494	C 2264	N 579	O 640	S 11	0	0
2	L	417	Total 3420	C 2213	N 570	O 625	S 12	0	0

- Molecule 3 is a RNA chain called RNA (5'-R(P*UP*GP*AP*CP*GP*GP*CP*UP*CP*UP*AP*AP*UP*CP*UP*AP*UP*UP*AP*GP*U)-3').

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	P		
3	E	17	Total 341	C 160	N 59	O 105	P 17	0	0
3	G	18	Total 360	C 169	N 61	O 112	P 18	0	0
3	M	18	Total 360	C 169	N 61	O 112	P 18	0	0
3	O	17	Total 341	C 160	N 59	O 105	P 17	0	0


- Molecule 4 is a DNA chain called DNA (25-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	P		
4	F	20	Total 411	196	77	118	20	3	0
4	H	20	Total 411	196	77	118	20	3	0
4	N	20	Total 411	196	77	118	20	3	0
4	P	20	Total 411	196	77	118	20	3	0


3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Piwi domain-containing protein




- Molecule 1: Piwi domain-containing protein







• Molecule 1: Piwi domain-containing protein




• Molecule 1: Piwi domain-containing protein







• Molecule 2: TIR domain-containing protein




• Molecule 2: TIR domain-containing protein







• Molecule 2: TIR domain-containing protein




• Molecule 2: TIR domain-containing protein







• Molecule 3: RNA (5'-R(P*UP*GP*AP*CP*GP*GP*CP*UP*CP*UP*AP*AP*UP*CP*UP*A P*UP*UP*AP*GP*U)-3')





• Molecule 3: RNA (5'-R(P*UP*GP*AP*CP*GP*GP*CP*UP*CP*UP*AP*AP*UP*CP*UP*A P*UP*UP*AP*GP*U)-3')



• Molecule 3: RNA (5'-R(P*UP*GP*AP*CP*GP*GP*CP*UP*CP*UP*AP*AP*UP*CP*UP*A P*UP*UP*AP*GP*U)-3')



• Molecule 3: RNA (5'-R(P*UP*GP*AP*CP*GP*GP*CP*UP*CP*UP*AP*AP*UP*CP*UP*A P*UP*UP*AP*GP*U)-3')





• Molecule 4: DNA (25-MER)





- Molecule 4: DNA (25-MER)



- Molecule 4: DNA (25-MER)



- Molecule 4: DNA (25-MER)



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	183347	Depositor
Resolution determination method	FSC 0.5 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	40	Depositor
Minimum defocus (nm)	1000	Depositor
Maximum defocus (nm)	2400	Depositor
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	2.131	Depositor
Minimum map value	-0.139	Depositor
Average map value	0.016	Depositor
Map value standard deviation	0.032	Depositor
Recommended contour level	0.13	Depositor
Map size (Å)	403.19998, 403.19998, 403.19998	wwPDB
Map dimensions	384, 384, 384	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.05, 1.05, 1.05	Depositor

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.26	0/3775	0.52	3/5111 (0.1%)
1	C	0.58	4/3708 (0.1%)	0.68	8/5016 (0.2%)
1	I	0.27	0/3791	0.50	0/5130
1	K	0.56	3/3804 (0.1%)	0.62	5/5150 (0.1%)
2	B	0.26	0/3504	0.53	1/4728 (0.0%)
2	D	0.25	0/3538	0.50	1/4771 (0.0%)
2	J	0.24	0/3581	0.48	0/4829
2	L	0.25	0/3503	0.50	0/4727
3	E	0.67	1/381 (0.3%)	0.83	0/581
3	G	0.65	1/402 (0.2%)	0.89	2/613 (0.3%)
3	M	0.64	1/402 (0.2%)	0.81	2/613 (0.3%)
3	O	0.66	1/381 (0.3%)	0.87	2/581 (0.3%)
4	F	0.51	0/461	0.87	0/709
4	H	0.52	0/461	0.87	0/709
4	N	0.53	0/461	0.88	0/709
4	P	0.50	0/461	0.88	0/709
All	All	0.39	11/32614 (0.0%)	0.59	24/44686 (0.1%)

The worst 5 of 11 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	144	PRO	CG-CD	-25.71	0.65	1.50
1	K	438	PRO	CG-CD	-25.16	0.67	1.50
1	K	438	PRO	CB-CG	13.33	2.16	1.50
1	C	144	PRO	CB-CG	10.80	2.04	1.50
3	G	1	U	OP3-P	-10.65	1.48	1.61

The worst 5 of 24 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	144	PRO	N-CD-CG	-17.52	76.92	103.20
1	K	438	PRO	N-CD-CG	-15.47	79.99	103.20
1	C	427	PRO	CA-N-CD	-13.04	93.24	111.50

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	K	438	PRO	CA-N-CD	-12.31	94.27	111.50
1	K	438	PRO	CB-CG-CD	-11.54	61.49	106.50

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3682	0	3656	131	0
1	C	3618	0	3625	159	0
1	I	3700	0	3710	137	0
1	K	3709	0	3687	119	0
2	B	3421	0	3366	119	0
2	D	3454	0	3426	110	0
2	J	3494	0	3481	105	0
2	L	3420	0	3363	106	0
3	E	341	0	182	12	0
3	G	360	0	192	14	0
3	M	360	0	192	9	0
3	O	341	0	182	19	0
4	F	411	0	212	4	0
4	H	411	0	212	8	0
4	N	411	0	212	15	0
4	P	411	0	212	12	0
All	All	31544	0	29910	959	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

The worst 5 of 959 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:49:ILE:HD11	2:B:77:GLU:HG2	1.56	0.85
1:I:302:GLN:HG2	1:I:312:VAL:HG13	1.59	0.83

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:K:469:ALA:HB2	3:O:3:A:H4'	1.61	0.83
2:D:178:ILE:HG13	2:D:333:PRO:HB2	1.63	0.81
1:I:255:THR:HG22	1:I:470:CYS:H	1.43	0.81

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	454/507 (90%)	422 (93%)	30 (7%)	2 (0%)	30	61
1	C	443/507 (87%)	402 (91%)	40 (9%)	1 (0%)	44	71
1	I	454/507 (90%)	415 (91%)	38 (8%)	1 (0%)	44	71
1	K	453/507 (89%)	418 (92%)	35 (8%)	0	100	100
2	B	415/452 (92%)	394 (95%)	20 (5%)	1 (0%)	44	71
2	D	415/452 (92%)	375 (90%)	40 (10%)	0	100	100
2	J	417/452 (92%)	385 (92%)	32 (8%)	0	100	100
2	L	415/452 (92%)	384 (92%)	27 (6%)	4 (1%)	13	42
All	All	3466/3836 (90%)	3195 (92%)	262 (8%)	9 (0%)	38	66

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	323	PRO
1	C	412	VAL
2	L	144	VAL
2	L	157	ILE
2	L	158	PHE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	395/446 (89%)	384 (97%)	11 (3%)	38	64
1	C	392/446 (88%)	377 (96%)	15 (4%)	28	56
1	I	402/446 (90%)	381 (95%)	21 (5%)	19	47
1	K	400/446 (90%)	386 (96%)	14 (4%)	31	58
2	B	367/414 (89%)	341 (93%)	26 (7%)	12	37
2	D	377/414 (91%)	359 (95%)	18 (5%)	21	50
2	J	385/414 (93%)	377 (98%)	8 (2%)	48	70
2	L	366/414 (88%)	351 (96%)	15 (4%)	26	54
All	All	3084/3440 (90%)	2956 (96%)	128 (4%)	27	53

5 of 128 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	L	105	TYR
2	L	201	ARG
1	C	430	GLN
1	C	425	TYR
2	L	205	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 13 such sidechains are listed below:

Mol	Chain	Res	Type
2	J	267	GLN
2	J	318	ASN
2	L	138	GLN
1	K	222	GLN
1	K	251	HIS

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
3	E	0/21	-	-
3	G	0/21	-	-
3	M	0/21	-	-
3	O	0/21	-	-
All	All	0/84	-	-

There are no RNA backbone outliers to report.

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-37708. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 192



Y Index: 192




Z Index: 192


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 192



Y Index: 194




Z Index: 239


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.13. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 422 nm³; this corresponds to an approximate mass of 382 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [i](#)



*Reported resolution corresponds to spatial frequency of 0.303\AA^{-1}


8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-37708 and PDB model 8WP2. Per-residue inclusion information can be found in section [3](#) on page [6](#).

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.13 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.13).



































9.4 Atom inclusion [i](#)



At the recommended contour level, 90% of all backbone atoms, 83% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.13) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.8290	 0.2960
A	 0.9280	 0.3340
B	 0.8530	 0.3010
C	 0.9160	 0.3170
D	 0.6250	 0.2190
E	 0.9440	 0.3860
F	 0.8880	 0.3180
G	 0.9110	 0.3020
H	 0.7520	 0.2000
I	 0.9430	 0.3480
J	 0.8480	 0.3230
K	 0.8770	 0.2910
L	 0.5610	 0.2120
M	 0.9530	 0.4190
N	 0.9170	 0.3680
O	 0.9530	 0.3130
P	 0.8520	 0.1910

