

wwPDB EM Validation Summary Report (i)

Dec 11, 2022 – 12:52 AM JST

PDB ID : 7WTV

EMDB ID : EMD-32802

Title : Cryo-EM structure of a human pre-40S ribosomal subunit - State RRP12-A2 Authors : Cheng, J.; Lau, B.; Thoms, M.; Ameismeier, M.; Berninghausen, O.; Hurt, E.;

Beckmann, R.

Deposited on : 2022-02-05

Resolution : 3.50 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp

with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev43

Mogul : 1.8.5 (274361), CSD as541be (2020)

MolProbity : 4.02b-467 buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

MapQ: 1.9.9

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.31.3

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $ELECTRON\ MICROSCOPY$

The reported resolution of this entry is 3.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive $(\# \mathrm{Entries})$	${ m EM\ structures} \ (\#{ m Entries})$
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826
RNA backbone	4643	859

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion <40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain								
1	2	1873	48%	29%	6% •	16%					
2	R	135	23%	9%	40%						
3	b	84		96%							
4	В	264	69%		11%	19%					
5	c	69	8	88%		12%					
6	Е	263	•	90%		9%					
7	e	59	34%	66%							

Continued on next page...







Mol	Chain	Length	Quality of chain							
2.0		225	16%							
33	A	295		60%	13%	27%				
			6%							
34	С	293		63%	10%	26%				
			5%							
35	V	83		84%		16%				



2 Entry composition (i)

There are 37 unique types of molecules in this entry. The entry contains 84177 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a RNA chain called 18S rRNA.

Mol	Chain	Residues		I	AltConf	Trace			
1	2	1574	Total 33614	C 15002	N 6033	O 11005	P 1574	0	0

• Molecule 2 is a protein called 40S ribosomal protein S17.

Mol	Chain	Residues		At	oms	AltConf	Trace		
2	R	81	Total 673	C 420	N 137	O 114	S 2	0	0

• Molecule 3 is a protein called 40S ribosomal protein S27.

Mol	Chain	Residues		At	oms	AltConf	Trace		
3	b	82	Total 640	C 402	N 118	O 113	S 7	0	0

• Molecule 4 is a protein called 40S ribosomal protein S3a.

Mol	Chain	Residues		At	AltConf	Trace			
4	В	213	Total 1729	C 1098	N 309	O 308	S 14	0	0

• Molecule 5 is a protein called 40S ribosomal protein S28.

Mol	Chain	Residues		Ato	ms	AltConf	Trace		
5	С	61	Total 471	C 288	N 95	O 86	S 2	0	0

• Molecule 6 is a protein called 40S ribosomal protein S4, X isoform.

Mol	Chain	Residues		Ato	AltConf	Trace			
6	Е	262	Total	C 1324	N 386	O 358	S	0	0



• Molecule 7 is a protein called 40S ribosomal protein S30.

Mol	Chain	Residues		Ato	$\mathbf{m}\mathbf{s}$	AltConf	Trace		
7	0	20	Total	С	N	О	S	0	0
1	е	20	179	110	43	25	1	0	U

• Molecule 8 is a protein called 40S ribosomal protein S5.

Mol	Chain	Residues		At	oms	AltConf	Trace		
0	E	189	Total	С	N	О	S	0	0
0	Г	109	1494	934	284	269	7	0	U

• Molecule 9 is a protein called 40S ribosomal protein S7.

Mol	Chain	Residues		At	oms	AltConf	Trace		
0	П	186	Total	С	N	О	S	0	0
9	11	100	1501	957	276	267	1	0	U

• Molecule 10 is a protein called 40S ribosomal protein S6.

Mol	Chain	Residues		At	oms			AltConf	Trace
10	G	230	Total 1862	C 1164	N 371	O 320	S 7	0	0

• Molecule 11 is a protein called 40S ribosomal protein S25.

Mol	Chain	Residues		At	oms		AltConf	Trace	
11	Z	72	Total 574	C 368	N 104	O 101	S 1	0	0

• Molecule 12 is a protein called 40S ribosomal protein S24.

Mol	Chain	Residues		At	oms		AltConf	Trace	
12	Y	124	Total 1014	C 641	N 198	O 170	S 5	0	0

• Molecule 13 is a protein called RNA-binding protein PNO1.

Mol	Chain	Residues		Atoms					Trace
13	Х	175	Total 1372	C 881	N 249	O 238	S 4	0	0

• Molecule 14 is a protein called 40S ribosomal protein S23.



Mol	Chain	Residues		At	oms			AltConf	Trace
1.4	v	1./1	Total	С	N	О	S	0	0
14	Λ	141	1098	693	219	183	3	0	U

• Molecule 15 is a protein called Bystin.

Mol	Chain	Residues		At	AltConf	Trace			
15	W	331	Total 2610	C 1671	N 477	O 453	S 9	0	0

• Molecule 16 is a protein called 40S ribosomal protein S15a.

Mol	Chain	Residues		At	oms	AltConf	Trace		
16	117	129	Total	С	N	О	S	0	0
10	VV	129	1033	659	193	175	6	0	U

• Molecule 17 is a protein called Pre-rRNA-processing protein TSR1 homolog.

Mol	Chain	Residues		At	oms			AltConf	Trace
17	u	642	Total 5168	C 3315	N 928	O 901	S 24	0	0

• Molecule 18 is a protein called 40S ribosomal protein S19.

Mol	Chain	Residues		At	oms	AltConf	Trace		
18	Т	144	Total 1122	C 703	N 217	O 199	S 3	0	0

• Molecule 19 is a protein called 40S ribosomal protein S18.

Mol	Chain	Residues		At	oms	AltConf	Trace		
10	C	126	Total	С	N	О	S	0	0
19	S	120	1046	663	204	178	1	U	U

• Molecule 20 is a protein called 40S ribosomal protein S16.

Mol	Chain	Residues		At	oms	AltConf	Trace		
20	Q	125	Total 998	C 637	N 185	O 173	S 3	0	0

• Molecule 21 is a protein called 40S ribosomal protein S15.



Mol	Chain	Residues		Atoms					Trace
21	Р	121	Total 1006	C 643	N 186	O 170	S 7	0	0

• Molecule 22 is a protein called 40S ribosomal protein S14.

Mol	Chain	Residues		At	oms	AltConf	Trace		
22	0	195	Total	С	N	О	S	0	0
22		135	1009	618	198	187	6	0	U

• Molecule 23 is a protein called 40S ribosomal protein S13.

Mol	Chain	Residues		At	oms			AltConf	Trace
99	N	1.40	Total	С	N	О	S	0	0
23	IN .	149	1202	770	228	203	1	0	U

• Molecule 24 is a protein called 40S ribosomal protein S11.

Mol	Chain	Residues		At	oms	AltConf	Trace		
24	т	151	Total	С	N	О	S	0	0
24	Ъ	151	1229	782	230	211	6	U	U

• Molecule 25 is a protein called 40S ribosomal protein S9.

Mol	Chain	Residues		At	oms	AltConf	Trace		
25	J	180	Total 1499	C 955	N 300	O 242	S 2	0	0

• Molecule 26 is a protein called 40S ribosomal protein S8.

Mol	Chain	Residues		Ato	AltConf	Trace			
26	I	205	Total 1682	C 1056	N 331	O 290	S 5	0	0

• Molecule 27 is a protein called Multifunctional methyltransferase subunit TRM112-like protein.

Mol	Chain	Residues		At	oms	AltConf	Trace		
27	r	118	Total 940	C 601	N 166	O 166	S 7	0	0

• Molecule 28 is a protein called Probable 18S rRNA (guanine-N(7))-methyltransferase.



Mol	Chain	Residues		At	oms			AltConf	Trace
28	q	235	Total 1869	C 1182	N 333	O 343	S 11	0	0

• Molecule 29 is a protein called RRP12-like protein.

Mol	Chain	Residues		A	toms			AltConf	Trace
29	K	993	Total	C	N	0	S	0	0
			7707	4938	1337	1387	45		

• Molecule 30 is a protein called 40S ribosomal protein S12.

Mol	Chain	Residues		At	oms			AltConf	Trace
30	M	108	Total	С	N	О	S	0	0
	111	100	837	530	147	153	7		

• Molecule 31 is a protein called Ubiquitin-40S ribosomal protein S27a.

Mol	Chain	Residues		Ato	$\mathbf{m}\mathbf{s}$	AltConf	Trace		
21	f	57	Total	С	N	О	S	0	0
31	1	91	465	295	89	74	7	0	U

• Molecule 32 is a protein called Ribosome biogenesis protein SLX9 homolog.

Mol	Chain	Residues		Ato	$\mathbf{m}\mathbf{s}$	AltConf	Trace		
32	${f z}$	52	Total 416	C 255	N 80	O 79	S 2	0	0

• Molecule 33 is a protein called 40S ribosomal protein SA.

Mol	Chain	Residues		Ato	AltConf	Trace			
33	A	216	Total 1705	C 1083	N 299	O 315	S 8	0	0

• Molecule 34 is a protein called 40S ribosomal protein S2.

Mol	Chain	Residues	Atoms				AltConf	Trace	
34	С	216	Total 1674	C 1085	N 287	O 292	S 10	0	0

• Molecule 35 is a protein called 40S ribosomal protein S21.




Mol	Chain	Residues	Atoms				AltConf	Trace	
35	V	83	Total 636	C 393	N 117	O 121	S 5	0	0

• Molecule 36 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	AltConf
36	2	1	Total Zn 1 1	0

 \bullet Molecule 37 is S-ADENOSYL-L-HOMOCYSTEINE (three-letter code: SAH) (formula: $C_{14}H_{20}N_6O_5S).$




Mol	Chain	Residues	Atoms					AltConf
97	G	1	Total	С	N	О	S	0
31	q	1	26	14	6	5	1	0




Residue-property plots (i) 3

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: 18S rRNA














• Molecule 9: 40S ribosomal protein S7



• Molecule 10: 40S ribosomal protein S6




• Molecule 11: 40S ribosomal protein S25






• Molecule 12: 40S ribosomal protein S24




• Molecule 13: RNA-binding protein PNO1




• Molecule 14: 40S ribosomal protein S23














Chain N: 89% 10% •

• Molecule 24: 40S ribosomal protein S11


Chain L: 78% 16% . .



LYS


• Molecule 25: 40S ribosomal protein S9

Chain J: 85% 8% · 7%



• Molecule 26: 40S ribosomal protein S8

Chain I: 83% 15% •





• Molecule 27: Multifunctional methyltransferase subunit TRM112-like protein

Chain r: 94% 6%



• Molecule 28: Probable 18S rRNA (guanine-N(7))-methyltransferase


Chain q: 83% • 16%









• Molecule 29: RRP12-like protein















 \bullet Molecule 35: 40S ribosomal protein S21

Chain V: 84% 16%





4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	25441	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE	Depositor
	CORRECTION; Relion	
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{Å}^2)$	44	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	2500	Depositor
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.473	Depositor
Minimum map value	-0.222	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.008	Depositor
Recommended contour level	0.02	Depositor
Map size (Å)	381.24, 381.24, 381.24	wwPDB
Map dimensions	360, 360, 360	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.059, 1.059, 1.059	Depositor



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, SAH, G7M

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

N/L-1	Clasia.	Bond	lengths	I	Bond angles
Mol	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	2	0.56	0/37551	1.04	165/58500~(0.3%)
2	R	0.32	0/680	0.66	0/905
3	b	0.31	0/653	0.59	1/876 (0.1%)
4	В	0.28	0/1756	0.56	0/2350
5	С	0.37	0/473	0.72	0/633
6	Е	0.37	0/2118	0.59	0/2849
7	е	0.31	0/180	0.61	0/232
8	F	0.38	0/1515	0.61	0/2037
9	Н	0.29	0/1524	0.60	0/2042
10	G	0.30	0/1885	0.60	1/2510 (0.0%)
11	Z	0.37	0/580	0.68	0/780
12	Y	0.36	0/1031	0.60	0/1370
13	X	0.28	0/1394	0.61	1/1880 (0.1%)
14	X	0.34	0/1116	0.58	0/1490
15	W	0.27	0/2656	0.58	0/3586
16	W	0.37	0/1050	0.59	0/1406
17	u	0.29	0/5296	0.57	0/7154
18	Τ	0.33	0/1142	0.57	0/1530
19	S	0.37	0/1063	0.68	2/1426~(0.1%)
20	Q	0.39	0/1012	0.67	0/1356
21	Р	0.34	0/1025	0.61	0/1369
22	О	0.30	0/1022	0.67	0/1372
23	N	0.31	0/1226	0.56	0/1649
24	L	0.40	0/1250	0.63	0/1673
25	J	0.36	0/1524	0.63	1/2035~(0.0%)
26	I	0.35	0/1711	0.64	0/2282
27	r	0.30	0/961	0.66	0/1301
28	q	0.34	0/1910	0.60	2/2572~(0.1%)
29	K	0.32	0/7851	0.61	2/10624~(0.0%)
30	M	0.24	0/845	0.52	0/1134
31	f	0.26	0/474	0.56	0/626
32	Z	0.27	0/420	0.68	0/564



Mol	Chain	Bond lengths		Bond angles		
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
33	A	0.28	0/1742	0.59	0/2367	
34	С	0.30	0/1710	0.58	1/2310~(0.0%)	
35	V	0.29	0/643	0.57	0/860	
All	All	0.44	0/88989	0.83	$176/127650 \ (0.1\%)$	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a maintain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
17	u	0	1
29	K	0	1
All	All	0	2

There are no bond length outliers.

The worst 5 of 176 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
1	2	501	С	N1-C2-O2	12.14	126.19	118.90
1	2	501	С	C2-N1-C1'	12.11	132.12	118.80
1	2	293	С	N1-C2-O2	11.68	125.91	118.90
1	2	1773	С	N3-C2-O2	-11.65	113.74	121.90
1	2	1838	U	N3-C2-O2	-10.77	114.66	122.20

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
29	K	378	VAL	Peptide
17	u	348	ASP	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	2	33614	0	16977	325	0
2	R	673	0	722	8	0
3	b	640	0	665	0	0
4	В	1729	0	1803	20	0
5	c	471	0	499	0	0
6	Ε	2076	0	2177	14	0
7	е	179	0	200	0	0
8	F	1494	0	1549	9	0
9	Н	1501	0	1593	21	0
10	G	1862	0	2018	23	0
11	Z	574	0	627	10	0
12	Y	1014	0	1082	11	0
13	X	1372	0	1453	0	0
14	X	1098	0	1167	12	0
15	W	2610	0	2644	0	0
16	W	1033	0	1080	7	0
17	u	5168	0	5230	0	0
18	Τ	1122	0	1153	15	0
19	S	1046	0	1094	9	0
20	Q	998	0	1065	12	0
21	Р	1006	0	1056	12	0
22	О	1009	0	1034	18	0
23	N	1202	0	1289	11	0
24	L	1229	0	1302	18	0
25	J	1499	0	1618	11	0
26	I	1682	0	1769	20	0
27	r	940	0	958	0	0
28	q	1869	0	1854	0	0
29	K	7707	0	7986	87	0
30	M	837	0	870	12	0
31	f	465	0	485	0	0
32	Z	416	0	424	0	0
33	A	1705	0	1706	22	0
34	С	1674	0	1764	20	0
35	V	636	0	637	8	0
36	2	1	0	0	0	0
37	q	26	0	19	0	0
All	All	84177	0	69569	647	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 647 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$\begin{array}{c} \text{Clash} \\ \text{overlap } (\text{\AA}) \end{array}$
1:2:536:A:N6	1:2:548:C:H42	1.45	1.14
1:2:748:C:H42	1:2:794:A:N6	1.48	1.12
1:2:536:A:H61	1:2:548:C:N4	1.46	1.11
1:2:748:C:N4	1:2:794:A:H61	1.46	1.11
1:2:1710:C:H42	1:2:1823:A:N6	1.51	1.07

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
2	R	79/135~(58%)	78 (99%)	1 (1%)	0	100	100
3	b	80/84~(95%)	73 (91%)	7 (9%)	0	100	100
4	В	211/264 (80%)	202 (96%)	9 (4%)	0	100	100
5	С	59/69~(86%)	57 (97%)	2 (3%)	0	100	100
6	E	260/263~(99%)	249 (96%)	11 (4%)	0	100	100
7	e	18/59 (30%)	18 (100%)	0	0	100	100
8	F	187/204 (92%)	176 (94%)	10 (5%)	1 (0%)	29	68
9	Н	184/194 (95%)	176 (96%)	8 (4%)	0	100	100
10	G	228/249 (92%)	217 (95%)	11 (5%)	0	100	100
11	Z	70/125 (56%)	63 (90%)	7 (10%)	0	100	100
12	Y	122/133 (92%)	119 (98%)	3 (2%)	0	100	100
13	X	173/252 (69%)	167 (96%)	6 (4%)	0	100	100
14	X	139/143 (97%)	138 (99%)	1 (1%)	0	100	100
15	W	321/437 (74%)	314 (98%)	7 (2%)	0	100	100
16	W	127/130 (98%)	123 (97%)	4 (3%)	0	100	100
17	u	630/804 (78%)	601 (95%)	29 (5%)	0	100	100



Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
18	Т	142/145~(98%)	139 (98%)	3 (2%)	0	100	100
19	S	124/152 (82%)	117 (94%)	7 (6%)	0	100	100
20	Q	123/146 (84%)	116 (94%)	7 (6%)	0	100	100
21	Р	117/145 (81%)	116 (99%)	1 (1%)	0	100	100
22	О	133/151 (88%)	125 (94%)	8 (6%)	0	100	100
23	N	147/151 (97%)	141 (96%)	6 (4%)	0	100	100
24	L	149/158 (94%)	138 (93%)	11 (7%)	0	100	100
25	J	178/194 (92%)	169 (95%)	9 (5%)	0	100	100
26	I	203/208 (98%)	194 (96%)	9 (4%)	0	100	100
27	r	116/125 (93%)	108 (93%)	8 (7%)	0	100	100
28	q	231/281 (82%)	223 (96%)	8 (4%)	0	100	100
29	K	973/1297 (75%)	934 (96%)	39 (4%)	0	100	100
30	M	102/132 (77%)	100 (98%)	2 (2%)	0	100	100
31	f	53/156 (34%)	51 (96%)	2 (4%)	0	100	100
32	Z	50/230 (22%)	49 (98%)	1 (2%)	0	100	100
33	A	214/295 (72%)	209 (98%)	5 (2%)	0	100	100
34	С	214/293 (73%)	206 (96%)	8 (4%)	0	100	100
35	V	81/83 (98%)	78 (96%)	3 (4%)	0	100	100
All	All	6238/7887 (79%)	5984 (96%)	253 (4%)	1 (0%)	100	100

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
8	F	163	PHE

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
2	R	72/122~(59%)	72 (100%)	0	100	100
3	b	74/76~(97%)	74 (100%)	0	100	100
4	В	$194/231\ (84\%)$	194 (100%)	0	100	100
5	c	52/62 (84%)	52 (100%)	0	100	100
6	E	$224/225 \ (100\%)$	224 (100%)	0	100	100
7	e	18/48~(38%)	18 (100%)	0	100	100
8	F	159/170~(94%)	159 (100%)	0	100	100
9	Н	167/174~(96%)	166 (99%)	1 (1%)	86	94
10	G	$200/218 \; (92\%)$	200 (100%)	0	100	100
11	Z	64/103~(62%)	64 (100%)	0	100	100
12	Y	108/115 (94%)	107 (99%)	1 (1%)	78	90
13	X	148/208 (71%)	148 (100%)	0	100	100
14	X	113/115 (98%)	113 (100%)	0	100	100
15	W	262/370~(71%)	260 (99%)	2 (1%)	81	91
16	W	112/113~(99%)	112 (100%)	0	100	100
17	u	561/705~(80%)	560 (100%)	1 (0%)	93	98
18	Т	114/115~(99%)	114 (100%)	0	100	100
19	S	110/132~(83%)	110 (100%)	0	100	100
20	Q	106/121 (88%)	106 (100%)	0	100	100
21	Р	111/130~(85%)	111 (100%)	0	100	100
22	О	105/119~(88%)	105 (100%)	0	100	100
23	N	130/131 (99%)	130 (100%)	0	100	100
24	L	$135/142\ (95\%)$	133 (98%)	2 (2%)	65	84
25	J	160/168 (95%)	159 (99%)	1 (1%)	86	94
26	I	178/180 (99%)	177 (99%)	1 (1%)	86	94
27	r	105/112 (94%)	105 (100%)	0	100	100
28	q	198/240 (82%)	198 (100%)	0	100	100
29	K	846/1094 (77%)	844 (100%)	2 (0%)	93	98
30	M	91/108 (84%)	90 (99%)	1 (1%)	73	88
31	f	51/140 (36%)	51 (100%)	0	100	100
32	Z	44/185 (24%)	44 (100%)	0	100	100
33	A	$180/243 \ (74\%)$	179 (99%)	1 (1%)	86	94



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
34	C	182/225 (81%)	182 (100%)	0	100	100	
35	V	67/67 (100%)	67 (100%)	0	100	100	
All	All	5441/6707 (81%)	5428 (100%)	13 (0%)	93	98	

5 of 13 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
25	J	177	ASN
26	I	141	ARG
33	A	42	LYS
29	K	1245	LYS
30	M	33	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
9	9 H 168		HIS
17	u	298	GLN
17	u	532	ASN
25	J	143	ASN

5.3.3 RNA (i)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers		
1	2	1555/1873 (83%)	353 (22%)	22 (1%)		

5 of 353 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	2	2	A
1	2	3	С
1	2	9	U
1	2	26	U
1	2	33	G

5 of 22 RNA pucker outliers are listed below:

Mol	Chain	Res	Type
1	2	1295	A



Mol	Chain	Res	Type
1	2	1511	U
1	2	1497	G
1	2	1534	С
1	2	332	G

5.4 Non-standard residues in protein, DNA, RNA chains (i)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bo	ond leng	hs	В	ond ang	les
IVIOI	туре	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	G7M	2	1639	1	20,26,27	2.66	7 (35%)	17,39,42	1.25	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	G7M	2	1639	1	=	1/3/25/26	0/3/3/3

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(A)
1	2	1639	G7M	C2-N2	6.43	1.49	1.34
1	2	1639	G7M	C2-N3	5.39	1.46	1.33
1	2	1639	G7M	C4-N3	4.81	1.49	1.37
1	2	1639	G7M	C6-N1	3.95	1.43	1.37
1	2	1639	G7M	C5-C6	3.24	1.53	1.45

All (2) bond angle outliers are listed below:

			V I			$\operatorname{Observed}(^{o})$	100001
1	2	1639	G7M	C2-N1-C6	-3.02	119.54	125.10



\mathbf{Mol}	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
1	2	1639	G7M	N1-C2-N3	-2.06	119.47	123.32

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	2	1639	G7M	C3'-C4'-C5'-O5'

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bo	nd leng	$ ag{ths}$	В	ond ang	les
IVIOI	Туре	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z >2
37	SAH	q	301	-	24,28,28	1.25	3 (12%)	25,40,40	1.85	4 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
37	SAH	q	301	-	-	3/11/31/31	0/3/3/3

All (3) bond length outliers are listed below:



Mol	Chain	Res	Type	Atoms	\mathbf{Z}	Observed(A)	Ideal(Å)
37	q	301	SAH	C2-N3	3.55	1.37	1.32
37	q	301	SAH	OXT-C	-2.22	1.23	1.30
37	q	301	SAH	C2-N1	2.21	1.38	1.33

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\mathrm{Ideal}(^{o})$
37	q	301	SAH	N3-C2-N1	-5.75	119.69	128.68
37	q	301	SAH	C5'-SD-CG	-4.34	89.26	102.27
37	q	301	SAH	OXT-C-O	-2.79	117.75	124.09
37	q	301	SAH	OXT-C-CA	2.25	121.03	113.38

There are no chirality outliers.

All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
37	q	301	SAH	C-CA-CB-CG
37	q	301	SAH	CA-CB-CG-SD
37	q	301	SAH	N-CA-CB-CG

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.




6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-32802. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections (i)


6.1.1 Primary map



The images above show the map projected in three orthogonal directions.

6.2 Central slices (i)

6.2.1 Primary map





The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices (i)

6.3.1 Primary map



The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal surface views (i)

6.4.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.02. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.



6.5 Mask visualisation (i)


This section was not generated. No masks/segmentation were deposited.



7 Map analysis (i)

This section contains the results of statistical analysis of the map.


7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



7.2 Volume estimate (i)




The volume at the recommended contour level is $979~\mathrm{nm}^3$; this corresponds to an approximate mass of $884~\mathrm{kDa}$.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



7.3 Rotationally averaged power spectrum (i)




^{*}Reported resolution corresponds to spatial frequency of 0.286 ${\rm \AA}^{-1}$



8 Fourier-Shell correlation (i)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC (i)



*Reported resolution corresponds to spatial frequency of 0.286 $\rm \AA^{-1}$



8.2 Resolution estimates (i)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)				
resolution estimate (A)	0.143	0.5	Half-bit		
Reported by author	3.50	-	-		
Author-provided FSC curve	3.47	4.48	3.65		
Unmasked-calculated*	-	-	-		


^{*}Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.



9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-32802 and PDB model 7WTV. Per-residue inclusion information can be found in section 3 on page 11.


9.1 Map-model overlay (i)



The images above show the 3D surface view of the map at the recommended contour level 0.02 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.




9.2 Q-score mapped to coordinate model (i)



The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


9.3 Atom inclusion mapped to coordinate model (i)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.02).



9.4 Atom inclusion (i)



At the recommended contour level, 84% of all backbone atoms, 81% of all non-hydrogen atoms, are inside the map.



9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (0.02) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.8082	0.3780
2	0.9354	0.4200
A	0.5763	0.3680
В	0.8511	0.3800
С	0.7485	0.4480
Е	0.9728	0.5700
F	0.8121	0.3130
G	0.9389	0.4770
Н	0.9124	0.4390
I	0.9304	0.4780
J	0.9653	0.5480
K	0.3070	0.1340
L	0.9397	0.5330
M	0.0545	0.0900
N	0.9468	0.4990
О	0.8663	0.3740
Р	0.8094	0.2890
Q	0.7381	0.2690
R	0.4876	0.1420
S	0.7527	0.2760
T	0.7619	0.2810
V	0.7170	0.4530
W	0.9802	0.5620
X	0.9206	0.5240
Y	0.9634	0.5380
Z	0.6542	0.2010
b	0.9713	0.5110
С	0.8455	0.3470
e	0.9821	0.5500
f	0.2860	0.0440
q	0.8115	0.3510
r	0.5261	0.1940
u	0.8810	0.4490
W	0.5150	0.1430
X	0.9159	0.4260
Z	0.0000	0.0370



