



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 4, 2024 – 08:50 PM EST

PDB ID : 1WWH  
Title : Crystal structure of the MPPN domain of mouse Nup35  
Authors : Handa, N.; Murayama, K.; Kukimoto, M.; Hamana, H.; Uchikubo, T.; Takemoto, C.; Terada, T.; Shirouzu, M.; Yokoyama, S.; RIKEN Structural Genomics/Proteomics Initiative (RSGI)  
Deposited on : 2005-01-05  
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtrriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	119	
1	B	119	
1	C	119	
1	D	119	

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2582 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called nucleoporin 35.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	81	Total 649	C 417	N 113	O 114	S 5	0	0	0
1	B	79	Total 631	C 405	N 109	O 112	S 5	0	0	0
1	C	79	Total 631	C 405	N 109	O 112	S 5	0	0	0
1	D	80	Total 639	C 411	N 110	O 113	S 5	0	0	0

There are 52 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	149	GLY	-	cloning artifact	UNP Q9D7J2
A	150	SER	-	cloning artifact	UNP Q9D7J2
A	151	SER	-	cloning artifact	UNP Q9D7J2
A	152	GLY	-	cloning artifact	UNP Q9D7J2
A	153	SER	-	cloning artifact	UNP Q9D7J2
A	154	SER	-	cloning artifact	UNP Q9D7J2
A	155	GLY	-	cloning artifact	UNP Q9D7J2
A	262	SER	-	cloning artifact	UNP Q9D7J2
A	263	GLY	-	cloning artifact	UNP Q9D7J2
A	264	PRO	-	cloning artifact	UNP Q9D7J2
A	265	SER	-	cloning artifact	UNP Q9D7J2
A	266	SER	-	cloning artifact	UNP Q9D7J2
A	267	GLY	-	cloning artifact	UNP Q9D7J2
B	149	GLY	-	cloning artifact	UNP Q9D7J2
B	150	SER	-	cloning artifact	UNP Q9D7J2
B	151	SER	-	cloning artifact	UNP Q9D7J2
B	152	GLY	-	cloning artifact	UNP Q9D7J2
B	153	SER	-	cloning artifact	UNP Q9D7J2
B	154	SER	-	cloning artifact	UNP Q9D7J2
B	155	GLY	-	cloning artifact	UNP Q9D7J2
B	262	SER	-	cloning artifact	UNP Q9D7J2

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	263	GLY	-	cloning artifact	UNP Q9D7J2
B	264	PRO	-	cloning artifact	UNP Q9D7J2
B	265	SER	-	cloning artifact	UNP Q9D7J2
B	266	SER	-	cloning artifact	UNP Q9D7J2
B	267	GLY	-	cloning artifact	UNP Q9D7J2
C	149	GLY	-	cloning artifact	UNP Q9D7J2
C	150	SER	-	cloning artifact	UNP Q9D7J2
C	151	SER	-	cloning artifact	UNP Q9D7J2
C	152	GLY	-	cloning artifact	UNP Q9D7J2
C	153	SER	-	cloning artifact	UNP Q9D7J2
C	154	SER	-	cloning artifact	UNP Q9D7J2
C	155	GLY	-	cloning artifact	UNP Q9D7J2
C	262	SER	-	cloning artifact	UNP Q9D7J2
C	263	GLY	-	cloning artifact	UNP Q9D7J2
C	264	PRO	-	cloning artifact	UNP Q9D7J2
C	265	SER	-	cloning artifact	UNP Q9D7J2
C	266	SER	-	cloning artifact	UNP Q9D7J2
C	267	GLY	-	cloning artifact	UNP Q9D7J2
D	149	GLY	-	cloning artifact	UNP Q9D7J2
D	150	SER	-	cloning artifact	UNP Q9D7J2
D	151	SER	-	cloning artifact	UNP Q9D7J2
D	152	GLY	-	cloning artifact	UNP Q9D7J2
D	153	SER	-	cloning artifact	UNP Q9D7J2
D	154	SER	-	cloning artifact	UNP Q9D7J2
D	155	GLY	-	cloning artifact	UNP Q9D7J2
D	262	SER	-	cloning artifact	UNP Q9D7J2
D	263	GLY	-	cloning artifact	UNP Q9D7J2
D	264	PRO	-	cloning artifact	UNP Q9D7J2
D	265	SER	-	cloning artifact	UNP Q9D7J2
D	266	SER	-	cloning artifact	UNP Q9D7J2
D	267	GLY	-	cloning artifact	UNP Q9D7J2

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	10	Total O 10 10	0	0
2	B	11	Total O 11 11	0	0
2	C	4	Total O 4 4	0	0
2	D	7	Total O 7 7	0	0



## 4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	59.49Å 104.24Å 110.04Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.83 – 2.70	Depositor
% Data completeness (in resolution range)	96.1 (19.83-2.70)	Depositor
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.08	Depositor
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.211 , 0.234	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	2582	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	54.0	wwPDB-VP

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.69	3/664 (0.5%)	0.75	1/894 (0.1%)
1	B	0.66	2/645 (0.3%)	0.72	0/868
1	C	0.69	2/645 (0.3%)	0.70	0/868
1	D	0.69	2/653 (0.3%)	0.68	0/879
All	All	0.68	9/2607 (0.3%)	0.71	1/3509 (0.0%)

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	233	GLU	CD-OE2	6.18	1.32	1.25
1	B	233	GLU	CD-OE1	6.10	1.32	1.25
1	C	233	GLU	CB-CG	5.91	1.63	1.52
1	B	233	GLU	CD-OE2	5.72	1.31	1.25
1	A	233	GLU	CD-OE1	5.58	1.31	1.25

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	218	LEU	CA-CB-CG	5.16	127.16	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	649	0	649	28	0
1	B	631	0	631	16	0
1	C	631	0	631	21	0
1	D	639	0	642	27	0
2	A	10	0	0	2	0
2	B	11	0	0	0	0
2	C	4	0	0	1	0
2	D	7	0	0	0	0
All	All	2582	0	2553	87	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

The worst 5 of 87 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:170:LEU:H	1:A:170:LEU:HD23	1.20	0.99
1:C:174:TRP:HB2	1:C:248:MET:HE3	1.58	0.86
1:A:236:MET:H	1:B:208:ASN:HD22	1.23	0.84
1:D:191:GLN:O	1:D:194:GLN:HG2	1.80	0.81
1:A:170:LEU:H	1:A:170:LEU:CD2	1.96	0.79

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	79/119 (66%)	74 (94%)	3 (4%)	2 (2%)	5	14
1	B	77/119 (65%)	70 (91%)	6 (8%)	1 (1%)	12	30
1	C	77/119 (65%)	70 (91%)	6 (8%)	1 (1%)	12	30
1	D	78/119 (66%)	75 (96%)	2 (3%)	1 (1%)	12	30

*Continued on next page...*



Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	311/476 (65%)	289 (93%)	17 (6%)	5 (2%)	9 24

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	232	GLY
1	A	233	GLU
1	B	233	GLU
1	C	233	GLU
1	D	233	GLU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	70/101 (69%)	64 (91%)	6 (9%)	10 24
1	B	68/101 (67%)	63 (93%)	5 (7%)	13 32
1	C	68/101 (67%)	61 (90%)	7 (10%)	7 16
1	D	69/101 (68%)	61 (88%)	8 (12%)	5 12
All	All	275/404 (68%)	249 (90%)	26 (10%)	8 20

5 of 26 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	205	ASN
1	C	249	GLU
1	D	224	LEU
1	C	246	ASN
1	D	175	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	246	ASN
1	D	182	GLN
1	D	205	ASN
1	C	182	GLN
1	C	191	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

### 6.4 Ligands

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers

EDS was not executed - this section is therefore empty.