

# wwPDB EM Validation Summary Report (i)

Dec 9, 2024 – 08:16 PM JST

PDB ID : 8WWL

EMDB ID : EMD-37892

Title : MCH-MCHR1-Gi complex, T2 state

Authors : Gong, W.; Ye, X.; Liu, G.

Deposited on : 2023-10-25

Resolution : 2.78 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

EMDB validation analysis : FAILED

MolProbity : 4.02b-467

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)

MapQ : FAILED

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.40

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $ELECTRON\ MICROSCOPY$ 

The reported resolution of this entry is 2.78 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	${ m EM\ structures} \ (\#{ m Entries})$		
Clashscore	210492	15764		
Ramachandran outliers	207382	16835		
Sidechain outliers	206894	16415		

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain							
1	A	354	55% 7% •	37%						
2	В	376	72%	19% 9%						
3	С	71	75%	6% 20%						
4	Е	255	78%	13% 9%						
5	R	624	40% 6%	54%						
6	L	19	68%	21% 11%						



# 2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 8964 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Guanine nucleotide-binding protein G(i) subunit alpha-1.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	224	Total 1749	C 1114	N 292	O 330	S 13	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	47	ASN	SER	conflict	UNP P63096
A	203	ALA	GLY	conflict	UNP P63096
A	245	ALA	GLU	conflict	UNP P63096
A	326	SER	ALA	conflict	UNP P63096

• Molecule 2 is a protein called Guanine nucleotide-binding protein G(I)/G(S)/G(T) subunit beta-1.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	В	342	Total 2622	C 1615	N 471	O 515	S 21	0	0

There are 37 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
В	-9	MET	-	initiating methionine	UNP P62873
В	-8	HIS	-	expression tag	UNP P62873
В	-7	HIS	-	expression tag	UNP P62873
В	-6	HIS	-	expression tag	UNP P62873
В	-5	HIS	-	expression tag	UNP P62873
В	-4	HIS	-	expression tag	UNP P62873
В	-3	HIS	-	expression tag	UNP P62873
В	-2	GLY	-	expression tag	UNP P62873
В	-1	SER	-	expression tag	UNP P62873
В	0	SER	-	expression tag	UNP P62873
В	1	GLY	-	expression tag	UNP P62873
В	341	GLY	-	expression tag	UNP P62873

Continued on next page...



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
В	342	SER	-	expression tag	UNP P62873
В	343	SER	-	expression tag	UNP P62873
В	344	GLY	-	expression tag	UNP P62873
В	345	GLY	-	expression tag	UNP P62873
В	346	GLY	-	expression tag	UNP P62873
В	347	GLY	-	expression tag	UNP P62873
В	348	SER	-	expression tag	UNP P62873
В	349	GLY	-	expression tag	UNP P62873
В	350	GLY	-	expression tag	UNP P62873
В	351	GLY	-	expression tag	UNP P62873
В	352	GLY	-	expression tag	UNP P62873
В	353	SER	-	expression tag	UNP P62873
В	354	SER	-	expression tag	UNP P62873
В	355	GLY	-	expression tag	UNP P62873
В	356	VAL	-	expression tag	UNP P62873
В	357	SER	-	expression tag	UNP P62873
В	358	GLY	-	expression tag	UNP P62873
В	359	TRP	-	expression tag	UNP P62873
В	360	ARG	-	expression tag	UNP P62873
В	361	LEU	-	expression tag	UNP P62873
В	362	PHE	-	expression tag	UNP P62873
В	363	LYS	-	expression tag	UNP P62873
В	364	LYS	-	expression tag	UNP P62873
В	365	ILE	-	expression tag	UNP P62873
В	366	SER	-	expression tag	UNP P62873

• Molecule 3 is a protein called Guanine nucleotide-binding protein G(I)/G(S)/G(O) subunit gamma-2.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	С	57	Total 436	C 273	N 77	O 83	S 3	0	0

• Molecule 4 is a protein called Antibody fragment ScFv16.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	E	231	Total 1779	C 1129	N 294	O 346	S 10	0	0

• Molecule 5 is a protein called Fusion protein 1, Melanin-concentrating hormone receptor 1, Fusion protein 2.



Mol	Chain	Residues	Atoms					AltConf	Trace
5	R	289	Total 2239	C 1490	N 358	O 374	S 17	0	0

 $\bullet$  Molecule 6 is a protein called Melanin-concentrating hormone.


Mol	Chain	Residues	Atoms					AltConf	Trace
6	L	17	Total 139	C 87	N 26	O 22	S 4	0	0




## 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Guanine nucleotide-binding protein G(i) subunit alpha-1



• Molecule 2: Guanine nucleotide-binding protein G(I)/G(S)/G(T) subunit beta-1



• Molecule 3: Guanine nucleotide-binding protein G(I)/G(S)/G(O) subunit gamma-2




• Molecule 4: Antibody fragment ScFv16

Chain E: 78% 13% 9%





• Molecule 5: Fusion protein 1, Melanin-concentrating hormone receptor 1, Fusion protein 2



• Molecule 6: Melanin-concentrating hormone

Chain L: 68% 21% 11%





# 4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	588396	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE	Depositor
	CORRECTION	
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{Å}^2)$	55	Depositor
Minimum defocus (nm)	1200	Depositor
Maximum defocus (nm)	2200	Depositor
Magnification	Not provided	
Image detector	GATAN K3 BIOQUANTUM (6k x 4k)	Depositor



## 5 Model quality (i)

### 5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles	
IVIOI	Chain	RMSZ		RMSZ	# Z  > 5
1	A	0.24	0/1779	0.41	0/2396
2	В	0.25	0/2669	0.51	0/3618
3	С	0.23	0/442	0.41	0/597
4	Е	0.26	0/1823	0.48	0/2472
5	R	0.24	0/2301	0.42	0/3152
6	L	0.25	0/141	0.57	0/187
All	All	0.25	0/9155	0.46	0/12422

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1749	0	1692	19	0
2	В	2622	0	2523	42	0
3	С	436	0	448	3	0
4	Е	1779	0	1714	20	0
5	R	2239	0	2281	26	0
6	L	139	0	136	4	0
All	All	8964	0	8794	102	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.



The worst 5 of 102 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$egin{aligned}  ext{Interatomic} \  ext{distance} \ ( ext{Å}) \end{aligned}$	$egin{aligned}  ext{Clash} \  ext{overlap } ( ext{Å}) \end{aligned}$
2:B:152:LEU:HD11	2:B:158:VAL:HG23	1.71	0.73
5:R:194:ASN:HB2	5:R:244:ILE:HD13	1.74	0.68
2:B:96:ARG:NH1	2:B:138:GLU:OE2	2.27	0.67
2:B:25:CYS:SG	2:B:259:GLN:NE2	2.69	0.66
2:B:79:LEU:HB3	2:B:93:ILE:HB	1.76	0.66

There are no symmetry-related clashes.

### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	220/354~(62%)	218 (99%)	2 (1%)	0	100 100
2	В	340/376 (90%)	332 (98%)	8 (2%)	0	100 100
3	С	55/71 (78%)	54 (98%)	1 (2%)	0	100 100
4	E	227/255 (89%)	222 (98%)	5 (2%)	0	100 100
5	R	287/624 (46%)	285 (99%)	2 (1%)	0	100 100
6	L	15/19 (79%)	15 (100%)	0	0	100 100
All	All	1144/1699 (67%)	1126 (98%)	18 (2%)	0	100 100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was



analysed.	and	the	total	number	of	residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	A	184/305~(60%)	180 (98%)	4 (2%)	47	76
2	В	284/306 (93%)	281 (99%)	3 (1%)	70	88
3	C	46/58 (79%)	46 (100%)	0	100	100
4	E	196/208 (94%)	195 (100%)	1 (0%)	86	95
5	R	$243/522 \ (47\%)$	241 (99%)	2 (1%)	79	92
6	L	15/18 (83%)	15 (100%)	0	100	100
All	All	968/1417 (68%)	958 (99%)	10 (1%)	71	89

5 of 10 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	Е	218	ARG
5	R	280	TYR
5	R	308	VAL
1	A	354	PHE
2	В	70	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
4	Е	183	GLN
5	R	194	ASN
5	R	281	GLN
2	В	259	GLN
1	A	188	HIS

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



## 5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry (i)

There are no ligands in this entry.

### 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

