



wwPDB EM Validation Summary Report ⓘ

Oct 20, 2024 – 11:00 AM EDT

PDB ID : 6WXU
EMDB ID : EMD-21963
Title : CryoEM structure of mouse DUOX1-DUOXA1 complex in the dimer-of-dimer state
Authors : Sun, J.
Deposited on : 2020-05-12
Resolution : 2.70 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev113
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4.02b-467
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1536	
1	C	1536	
2	B	341	
2	D	341	
3	E	11	
3	F	11	

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 18156 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Dual oxidase 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	834	6624	4287	1151	1161	25	0	0
1	C	834	6624	4287	1151	1161	25	0	0


There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	16	GLY	-	expression tag	UNP A2AQ92
A	17	PRO	-	expression tag	UNP A2AQ92
A	18	SER	-	expression tag	UNP A2AQ92
A	19	ARG	-	expression tag	UNP A2AQ92
C	16	GLY	-	expression tag	UNP A2AQ92
C	17	PRO	-	expression tag	UNP A2AQ92
C	18	SER	-	expression tag	UNP A2AQ92
C	19	ARG	-	expression tag	UNP A2AQ92

- Molecule 2 is a protein called Dual oxidase maturation factor 1.


Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	B	273	2093	1390	340	351	12	0	0
2	D	273	2093	1390	340	351	12	0	0

- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
3	E	11	127	70	2	55	0	0
3	F	11	127	70	2	55	0	0

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms				AltConf
			Total	C	N	O	
4	A	1	14	8	1	5	0
4	A	1	14	8	1	5	0
4	A	1	14	8	1	5	0
4	B	1	14	8	1	5	0
4	B	1	14	8	1	5	0
4	C	1	14	8	1	5	0
4	C	1	14	8	1	5	0

Continued on next page...



Mol	Chain	Residues	Atoms				AltConf
			Total	C	O	P	
6	A	1	36	27	8	1	0
6	C	1	36	27	8	1	0

- Molecule 7 is DIUNDECYL PHOSPHATIDYL CHOLINE (three-letter code: PLC) (formula: $C_{32}H_{65}NO_8P$).



Mol	Chain	Residues	Atoms				AltConf	
			Total	C	N	O		P
7	B	1	42	32	1	8	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					AltConf
			Total	C	N	O	P	
7	D	1	42	32	1	8	1	0

VAL SER VAL VAL HIS ILE TYR ILE THR VAL VAL ALA ALA GLU LYS PHE ASP ASP ARG THR THR MET LEU LEU TYR VAL ILE ILE CYS GLU ARG HIS PHE PHE LYS VAL LEU LEU ASN ARG THR SER PHE THR THR GLY HIS LEU LEU ASN ARG THR HIS HIS TYR ARG ILE THR HIS PHE ARG PRO PRO PHE PHE PHE PHE PHE ASN SER SER LEU GLN GLU

VAL HIS PRO GLN VAL ARG LYS ILE ILE VAL PHE SER CYS GLY PRO PRO GLY ARG MET THR LYS ASN LEU VAL GLU ILE ILE CYS ARG GLN LEU ILE ASN ARG THR HIS PHE THR SER HIS HIS TYR ARG ILE ASN PHE

- Molecule 2: Dual oxidase maturation factor 1

Chain B:  73% 6% 20%

MET ALA A3 T22 F31 I41 R46 G47 R50 L51 F52 L61 V66 V70 S74 E75 A83 S96 Q102 T115 Q119 T123 Y138 G149 A158 T162 L203 T240 P249 T257 Q275 PRO HIS ARG LEU LYS

ALA PHE PHE ASN GLN SER GLU ASP PRO VAL LEU TRP GLY SER GLU GLY GLY LEU LEU SER LEU PRO HIS TYR ARG ILE ILE ALA GLU SER PRO THR GLN ASP ILE PRO MET SER VAL ALA SER SER GLU THR CYS PHE LYS GLU HIS PRO HIS LYS SER ASP CYS SER

LEU

- Molecule 2: Dual oxidase maturation factor 1

Chain D:  73% 7% 20%

MET ALA A3 T22 F31 I41 R46 G47 R50 L51 F52 L61 V66 V70 S74 E75 A83 F90 S96 Q102 T115 Q119 T123 G149 T162 L198 L203 T240 P249 T257 M268 Q275 PRO HIS ARG

LEU LYS PHE PHE ASN SER SER ASP VAL LEU TRP SER LEU GLU GLY LEU LEU SER LEU PRO HIS TYR SER ARG ILE ILE ALA GLU SER PRO THR GLN ASP ILE PRO MET SER VAL ALA SER SER GLU THR CYS PHE LYS GLU HIS PRO HIS LYS SER ASP CYS

CYS SER LEU

- Molecule 3: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E:  9% 73% 18%

MAG1 MAG2 BMA3 MAN4 MAN5 MAN6 MAN7 MAN8 MAN9 MAN10 MAN11

- Molecule 3: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:  9% 73% 18%

MAG1 MAG2 BMA3 MAN4 MAN5 MAN6 MAN7 MAN8 MAN9 MAN10 MAN11

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C2	Depositor
Number of particles used	302097	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	65.4	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	3.303	Depositor
Minimum map value	-2.021	Depositor
Average map value	0.027	Depositor
Map value standard deviation	0.126	Depositor
Recommended contour level	0.5	Depositor
Map size (\AA)	248.40001, 248.40001, 248.40001	wwPDB
Map dimensions	230, 230, 230	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (\AA)	1.08, 1.08, 1.08	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MAN, BMA, NAG, PLC, PX2, HEC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.37	0/6830	0.46	1/9301 (0.0%)
1	C	0.37	0/6830	0.46	1/9301 (0.0%)
2	B	0.34	0/2154	0.44	0/2950
2	D	0.34	0/2154	0.44	0/2950
All	All	0.37	0/17968	0.45	2/24502 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	1
2	D	0	1
All	All	0	2

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	C	1128	ASP	CB-CG-OD2	5.24	123.01	118.30
1	A	1128	ASP	CB-CG-OD2	5.21	122.98	118.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	115	THR	Peptide
2	D	115	THR	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6624	0	6436	53	0
1	C	6624	0	6436	52	0
2	B	2093	0	2125	12	0
2	D	2093	0	2125	13	0
3	E	127	0	106	2	0
3	F	127	0	106	2	0
4	A	42	0	39	1	0
4	B	28	0	26	1	0
4	C	42	0	39	1	0
4	D	28	0	26	1	0
5	A	86	0	64	7	0
5	C	86	0	64	7	0
6	A	36	0	52	1	0
6	C	36	0	52	1	0
7	B	42	0	64	2	0
7	D	42	0	64	2	0
All	All	18156	0	17824	137	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 137 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:22:THR:OG1	7:D:414:PLC:H62	1.84	0.78
2:B:22:THR:OG1	7:B:414:PLC:H62	1.84	0.77
7:D:414:PLC:H63	3:F:2:NAG:H2	1.68	0.76
1:C:151:TRP:O	1:C:153:ARG:NH1	2.19	0.76
7:B:414:PLC:H63	3:E:2:NAG:H2	1.68	0.75

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	828/1536 (54%)	788 (95%)	40 (5%)	0	100	100
1	C	828/1536 (54%)	788 (95%)	40 (5%)	0	100	100
2	B	271/341 (80%)	258 (95%)	13 (5%)	0	100	100
2	D	271/341 (80%)	258 (95%)	13 (5%)	0	100	100
All	All	2198/3754 (59%)	2092 (95%)	106 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	702/1348 (52%)	694 (99%)	8 (1%)	70	87
1	C	702/1348 (52%)	694 (99%)	8 (1%)	70	87
2	B	219/287 (76%)	215 (98%)	4 (2%)	54	80
2	D	219/287 (76%)	215 (98%)	4 (2%)	54	80
All	All	1842/3270 (56%)	1818 (99%)	24 (1%)	64	85

5 of 24 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	244	PHE
1	C	468	THR
1	C	296	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	480	GLU
1	A	480	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 42 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	289	ASN
1	C	1188	GLN
1	C	302	GLN
1	C	391	GLN
2	D	82	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

22 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	E	1	2,3	14,14,15	0.38	0	17,19,21	0.57	0
3	MAN	E	10	3	11,11,12	0.65	0	15,15,17	1.02	2 (13%)
3	MAN	E	11	3	11,11,12	0.66	0	15,15,17	1.11	2 (13%)
3	NAG	E	2	3	14,14,15	0.68	1 (7%)	17,19,21	0.91	0
3	BMA	E	3	3	11,11,12	0.63	0	15,15,17	0.77	1 (6%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	MAN	E	4	3	11,11,12	0.78	0	15,15,17	1.22	2 (13%)
3	MAN	E	5	3	11,11,12	0.87	0	15,15,17	1.37	2 (13%)
3	MAN	E	6	3	11,11,12	0.86	1 (9%)	15,15,17	0.92	1 (6%)
3	MAN	E	7	3	11,11,12	0.75	0	15,15,17	1.12	2 (13%)
3	MAN	E	8	3	11,11,12	0.67	0	15,15,17	1.21	2 (13%)
3	MAN	E	9	3	11,11,12	0.78	1 (9%)	15,15,17	0.82	1 (6%)
3	NAG	F	1	2,3	14,14,15	0.37	0	17,19,21	0.57	0
3	MAN	F	10	3	11,11,12	0.65	0	15,15,17	1.02	2 (13%)
3	MAN	F	11	3	11,11,12	0.66	0	15,15,17	1.11	2 (13%)
3	NAG	F	2	3	14,14,15	0.68	1 (7%)	17,19,21	0.90	0
3	BMA	F	3	3	11,11,12	0.63	0	15,15,17	0.77	1 (6%)
3	MAN	F	4	3	11,11,12	0.78	0	15,15,17	1.22	2 (13%)
3	MAN	F	5	3	11,11,12	0.87	0	15,15,17	1.37	2 (13%)
3	MAN	F	6	3	11,11,12	0.85	1 (9%)	15,15,17	0.92	1 (6%)
3	MAN	F	7	3	11,11,12	0.75	0	15,15,17	1.12	2 (13%)
3	MAN	F	8	3	11,11,12	0.67	0	15,15,17	1.21	2 (13%)
3	MAN	F	9	3	11,11,12	0.77	1 (9%)	15,15,17	0.82	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	E	1	2,3	-	0/6/23/26	0/1/1/1
3	MAN	E	10	3	-	0/2/19/22	0/1/1/1
3	MAN	E	11	3	-	0/2/19/22	0/1/1/1
3	NAG	E	2	3	-	4/6/23/26	0/1/1/1
3	BMA	E	3	3	-	0/2/19/22	0/1/1/1
3	MAN	E	4	3	-	2/2/19/22	0/1/1/1
3	MAN	E	5	3	-	0/2/19/22	0/1/1/1
3	MAN	E	6	3	-	0/2/19/22	0/1/1/1
3	MAN	E	7	3	-	2/2/19/22	0/1/1/1
3	MAN	E	8	3	-	1/2/19/22	0/1/1/1
3	MAN	E	9	3	-	0/2/19/22	0/1/1/1
3	NAG	F	1	2,3	-	0/6/23/26	0/1/1/1
3	MAN	F	10	3	-	0/2/19/22	0/1/1/1
3	MAN	F	11	3	-	0/2/19/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	F	2	3	-	4/6/23/26	0/1/1/1
3	BMA	F	3	3	-	0/2/19/22	0/1/1/1
3	MAN	F	4	3	-	2/2/19/22	0/1/1/1
3	MAN	F	5	3	-	0/2/19/22	0/1/1/1
3	MAN	F	6	3	-	0/2/19/22	0/1/1/1
3	MAN	F	7	3	-	2/2/19/22	0/1/1/1
3	MAN	F	8	3	-	1/2/19/22	0/1/1/1
3	MAN	F	9	3	-	0/2/19/22	0/1/1/1

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	2	NAG	O5-C1	-2.39	1.39	1.43
3	F	2	NAG	O5-C1	-2.39	1.39	1.43
3	E	9	MAN	O5-C1	-2.38	1.39	1.43
3	F	9	MAN	O5-C1	-2.37	1.39	1.43
3	E	6	MAN	O5-C1	-2.26	1.39	1.43

The worst 5 of 30 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	5	MAN	C1-O5-C5	3.78	117.25	112.19
3	E	5	MAN	C1-O5-C5	3.75	117.21	112.19
3	E	5	MAN	O2-C2-C3	-3.09	103.75	110.15
3	F	5	MAN	O2-C2-C3	-3.08	103.77	110.15
3	E	8	MAN	O2-C2-C3	-3.07	103.79	110.15

There are no chirality outliers.

5 of 18 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	E	7	MAN	C4-C5-C6-O6
3	F	7	MAN	C4-C5-C6-O6
3	E	4	MAN	O5-C5-C6-O6
3	F	4	MAN	O5-C5-C6-O6
3	E	2	NAG	C4-C5-C6-O6


There are no ring outliers.

4 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	F	5	MAN	1	0
3	E	2	NAG	1	0
3	E	5	MAN	1	0
3	F	2	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

18 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	NAG	A	1603	1	14,14,15	0.61	1 (7%)	17,19,21	1.37	2 (11%)
4	NAG	C	1602	1	14,14,15	0.36	0	17,19,21	0.35	0
4	NAG	B	413	2	14,14,15	0.23	0	17,19,21	0.40	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	C	1601	1	14,14,15	0.18	0	17,19,21	0.48	0
4	NAG	D	413	2	14,14,15	0.23	0	17,19,21	0.40	0
7	PLC	D	414	-	41,41,41	1.02	3 (7%)	47,49,49	1.36	4 (8%)
4	NAG	C	1603	1	14,14,15	0.63	1 (7%)	17,19,21	1.36	2 (11%)
4	NAG	D	401	2	14,14,15	0.20	0	17,19,21	0.50	0
5	HEC	A	1605	1	32,50,50	1.72	8 (25%)	30,82,82	1.68	6 (20%)
5	HEC	C	1604	1	32,50,50	1.74	8 (25%)	30,82,82	1.65	8 (26%)
5	HEC	C	1605	1	32,50,50	1.72	8 (25%)	30,82,82	1.67	6 (20%)
7	PLC	B	414	-	41,41,41	1.02	3 (7%)	47,49,49	1.36	4 (8%)
5	HEC	A	1604	1	32,50,50	1.74	8 (25%)	30,82,82	1.65	8 (26%)
4	NAG	A	1602	1	14,14,15	0.36	0	17,19,21	0.35	0
4	NAG	A	1601	1	14,14,15	0.18	0	17,19,21	0.49	0
6	PX2	A	1606	-	35,35,35	1.09	2 (5%)	38,40,40	1.40	5 (13%)
4	NAG	B	401	2	14,14,15	0.20	0	17,19,21	0.50	0
6	PX2	C	1606	-	35,35,35	1.09	2 (5%)	38,40,40	1.40	5 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	1603	1	-	6/6/23/26	0/1/1/1
4	NAG	C	1602	1	-	2/6/23/26	0/1/1/1
4	NAG	B	413	2	-	2/6/23/26	0/1/1/1
4	NAG	C	1601	1	-	2/6/23/26	0/1/1/1
4	NAG	D	413	2	-	2/6/23/26	0/1/1/1
7	PLC	D	414	-	-	18/45/45/45	-
4	NAG	C	1603	1	-	6/6/23/26	0/1/1/1
4	NAG	D	401	2	-	2/6/23/26	0/1/1/1
5	HEC	A	1605	1	-	2/10/54/54	-
5	HEC	C	1604	1	-	4/10/54/54	-
5	HEC	C	1605	1	-	2/10/54/54	-
7	PLC	B	414	-	-	18/45/45/45	-
5	HEC	A	1604	1	-	4/10/54/54	-
4	NAG	A	1602	1	-	2/6/23/26	0/1/1/1
4	NAG	A	1601	1	-	2/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	PX2	A	1606	-	-	18/37/37/37	-
4	NAG	B	401	2	-	2/6/23/26	0/1/1/1
6	PX2	C	1606	-	-	18/37/37/37	-

The worst 5 of 44 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	C	1604	HEC	CBC-CAC	-4.93	1.31	1.49
5	A	1604	HEC	CBC-CAC	-4.92	1.31	1.49
5	C	1605	HEC	CBC-CAC	-4.81	1.31	1.49
5	A	1605	HEC	CBC-CAC	-4.80	1.31	1.49
5	A	1604	HEC	CBB-CAB	-4.79	1.31	1.49

The worst 5 of 50 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	B	414	PLC	O2-C'-C1'	5.66	123.73	111.48
7	D	414	PLC	O2-C'-C1'	5.66	123.72	111.48
4	A	1603	NAG	C2-N2-C7	4.48	128.91	122.90
4	C	1603	NAG	C2-N2-C7	4.48	128.91	122.90
6	A	1606	PX2	C3-C2-C1	-3.80	102.92	111.78

There are no chirality outliers.

5 of 112 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	1606	PX2	C1-O4-P1-O1
6	A	1606	PX2	C1-O4-P1-O2
6	A	1606	PX2	C1-O4-P1-O3
6	C	1606	PX2	C1-O4-P1-O1
6	C	1606	PX2	C1-O4-P1-O2

There are no ring outliers.

12 monomers are involved in 24 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1603	NAG	1	0
7	D	414	PLC	2	0
4	C	1603	NAG	1	0
4	D	401	NAG	1	0


Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	1605	HEC	2	0
5	C	1604	HEC	5	0
5	C	1605	HEC	2	0
7	B	414	PLC	2	0
5	A	1604	HEC	5	0
6	A	1606	PX2	1	0
4	B	401	NAG	1	0
6	C	1606	PX2	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




















5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-21963. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map




The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [\(i\)](#)


6.3.1 Primary map



X Index: 100



Y Index: 118




Z Index: 80


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [\(i\)](#)


6.4.1 Primary map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.5. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 127 nm³; this corresponds to an approximate mass of 115 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [i](#)



*Reported resolution corresponds to spatial frequency of 0.370\AA^{-1}


8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-21963 and PDB model 6WXU. Per-residue inclusion information can be found in section 3 on page 8.

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.5 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.5).















9.4 Atom inclusion [i](#)



At the recommended contour level, 93% of all backbone atoms, 91% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.5) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.9130	 0.5940
A	 0.9220	 0.5980
B	 0.8810	 0.5810
C	 0.9220	 0.5990
D	 0.8810	 0.5820
E	 0.9450	 0.5950
F	 0.9450	 0.5940

