



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 2, 2024 – 10:44 pm GMT

PDB ID : 2WYD
Title : The quorum quenching N-acyl homoserine lactone acylase PvdQ in complex with dodecanoic acid
Authors : Bokhove, M.; Nadal Jimenez, P.; Quax, W.J.; Dijkstra, B.W.
Deposited on : 2009-11-16
Resolution : 1.90 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	7293 (1.90-1.90)
Clashscore	180529	8090 (1.90-1.90)
Ramachandran outliers	177936	8022 (1.90-1.90)
Sidechain outliers	177891	8022 (1.90-1.90)
RSRZ outliers	164620	7292 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	170	 2% 91% 5%
2	B	546	 7% 94% 6%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 6137 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called ACYL-HOMOSERINE LACTONE ACYLASE PVDQ SUB-UNIT ALPHA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	164	1298	818	234	244	2	0	4	0

- Molecule 2 is a protein called ACYL-HOMOSERINE LACTONE ACYLASE PVDQ SUB-UNIT BETA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	546	4331	2714	794	812	11	0	9	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
3	A	1	6	3	3	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is LAURIC ACID (three-letter code: DAO) (formula: $C_{12}H_{24}O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			14	12	2		
4	B	1	Total	C	O	0	0
			14	12	2		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	136	Total 136	O 136	0	0
5	B	296	Total 296	O 296	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: ACYL-HOMOSERINE LACTONE ACYLASE PVDQ SUBUNIT ALPHA

Chain A: 



- Molecule 2: ACYL-HOMOSERINE LACTONE ACYLASE PVDQ SUBUNIT BETA

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	120.78Å 166.37Å 94.31Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.13 – 1.90 39.13 – 1.90	Depositor EDS
% Data completeness (in resolution range)	98.1 (39.13-1.90) 98.0 (39.13-1.90)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.72 (at 1.89Å)	Xtrriage
Refinement program	REFMAC 5.5.0096	Depositor
R, R_{free}	0.167 , 0.194 0.185 , 0.210	Depositor DCC
R_{free} test set	3686 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	21.5	Xtrriage
Anisotropy	0.022	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 44.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6137	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.64% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: DAO, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.55	0/1326	0.56	0/1800
2	B	0.46	0/4436	0.56	1/6032 (0.0%)
All	All	0.48	0/5762	0.56	1/7832 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	497	LEU	CA-CB-CG	5.86	128.78	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1298	0	1250	9	0
2	B	4331	0	4247	21	0
3	A	18	0	24	0	0
3	B	30	0	40	4	0
4	B	28	0	46	5	0
5	A	136	0	0	3	0
5	B	296	0	0	1	0
All	All	6137	0	5607	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:149[B]:GLU:HG3	4:B:1553:DAO:H31	1.66	0.78
1:A:149[A]:GLU:OE1	5:A:2121:HOH:O	2.02	0.73
1:A:149[B]:GLU:CG	4:B:1553:DAO:H31	2.19	0.72
2:B:380:PHE:O	3:B:1550:GOL:H2	2.02	0.59
2:B:1[A]:SER:HB3	2:B:22:PRO:HA	1.83	0.59
2:B:498:LEU:H	2:B:514:GLN:NE2	2.01	0.59
2:B:444[B]:ARG:NH1	5:B:2237:HOH:O	2.37	0.57
1:A:23[B]:ARG:HD3	5:A:2016:HOH:O	2.06	0.54
2:B:202:ASN:HD22	2:B:257:PRO:HG2	1.72	0.53
2:B:185:PRO:HB2	2:B:186:TRP:CD1	2.46	0.49
2:B:25:PRO:HD3	4:B:1553:DAO:H81	1.95	0.49
2:B:24:PHE:CE1	4:B:1553:DAO:H62	2.48	0.49
2:B:350:ARG:CZ	2:B:426:GLU:HG2	2.43	0.48
2:B:387:TRP:CD2	2:B:400:PRO:HB2	2.50	0.47
2:B:477:GLY:HA3	4:B:1553:DAO:H101	1.96	0.46
1:A:68:SER:C	3:B:1548:GOL:H12	2.37	0.45
1:A:90:GLN:HG3	5:A:2069:HOH:O	2.15	0.45
1:A:149[A]:GLU:O	1:A:154[A]:GLN:HG2	2.18	0.44
1:A:28:ARG:NH2	1:A:105:GLU:OE1	2.51	0.44
2:B:241:ARG:HD3	3:B:1551:GOL:C1	2.48	0.44
2:B:188:ASN:HD22	2:B:202:ASN:HA	1.83	0.43
2:B:490:GLU:H	2:B:490:GLU:CD	2.22	0.43
2:B:207:PRO:HB3	2:B:239:TRP:CE2	2.55	0.42
1:A:74:LEU:HB3	1:A:75:PRO:HD3	2.01	0.42
2:B:1[A]:SER:CB	2:B:22:PRO:HA	2.50	0.42
2:B:25:PRO:HG2	2:B:30:MET:HE2	2.02	0.41
2:B:47:GLY:HA3	2:B:58:ILE:O	2.20	0.41
2:B:49:SER:OG	2:B:56:VAL:HA	2.21	0.41
2:B:241:ARG:HD3	3:B:1551:GOL:H12	2.03	0.41
2:B:218:ALA:HB1	2:B:223:VAL:HG21	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	166/170 (98%)	162 (98%)	4 (2%)	0	100	100
2	B	552/546 (101%)	532 (96%)	20 (4%)	0	100	100
All	All	718/716 (100%)	694 (97%)	24 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	123/126 (98%)	123 (100%)	0	100	100
2	B	456/447 (102%)	453 (99%)	3 (1%)	81	83
All	All	579/573 (101%)	576 (100%)	3 (0%)	86	88

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	24	PHE
2	B	162	TRP
2	B	432	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	128	GLN
2	B	34	GLN
2	B	188	ASN
2	B	202	ASN
2	B	282	GLN
2	B	514	GLN
2	B	540	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	B	1549	-	5,5,5	0.37	0	5,5,5	0.29	0
3	GOL	B	1551	-	5,5,5	0.43	0	5,5,5	0.27	0
3	GOL	B	1547	-	5,5,5	0.41	0	5,5,5	0.33	0
4	DAO	B	1553	-	13,13,13	0.52	0	13,13,13	1.06	1 (7%)
3	GOL	A	1171	-	5,5,5	0.35	0	5,5,5	0.23	0
3	GOL	A	1170	-	5,5,5	0.28	0	5,5,5	0.48	0
3	GOL	B	1548	-	5,5,5	0.71	0	5,5,5	1.08	1 (20%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	DAO	B	1552	-	13,13,13	0.61	0	13,13,13	0.62	0
3	GOL	A	1172	-	5,5,5	0.36	0	5,5,5	0.55	0
3	GOL	B	1550	-	5,5,5	0.44	0	5,5,5	0.40	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	B	1549	-	-	0/4/4/4	-
3	GOL	B	1551	-	-	0/4/4/4	-
3	GOL	B	1547	-	-	4/4/4/4	-
4	DAO	B	1553	-	-	7/11/11/11	-
3	GOL	A	1171	-	-	1/4/4/4	-
3	GOL	A	1170	-	-	2/4/4/4	-
3	GOL	B	1548	-	-	4/4/4/4	-
4	DAO	B	1552	-	-	1/11/11/11	-
3	GOL	A	1172	-	-	2/4/4/4	-
3	GOL	B	1550	-	-	3/4/4/4	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	1553	DAO	C3-C2-C1	-2.74	107.57	114.47
3	B	1548	GOL	C3-C2-C1	-2.27	102.87	111.70

There are no chirality outliers.

All (24) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1172	GOL	O1-C1-C2-C3
3	B	1547	GOL	O1-C1-C2-C3
3	B	1547	GOL	O2-C2-C3-O3
3	B	1548	GOL	O1-C1-C2-C3
3	B	1550	GOL	O1-C1-C2-C3
4	B	1553	DAO	C1-C2-C3-C4
4	B	1553	DAO	C5-C6-C7-C8
3	A	1170	GOL	C1-C2-C3-O3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	B	1547	GOL	C1-C2-C3-O3
3	B	1548	GOL	C1-C2-C3-O3
3	A	1172	GOL	O1-C1-C2-O2
3	B	1547	GOL	O1-C1-C2-O2
3	B	1548	GOL	O1-C1-C2-O2
3	B	1550	GOL	O1-C1-C2-O2
4	B	1553	DAO	C11-C10-C9-C8
3	B	1548	GOL	O2-C2-C3-O3
3	A	1171	GOL	O1-C1-C2-C3
4	B	1553	DAO	C9-C10-C11-C12
4	B	1553	DAO	C2-C3-C4-C5
4	B	1553	DAO	C3-C4-C5-C6
3	B	1550	GOL	O2-C2-C3-O3
4	B	1552	DAO	C2-C3-C4-C5
4	B	1553	DAO	C6-C7-C8-C9
3	A	1170	GOL	O2-C2-C3-O3

There are no ring outliers.

4 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	1551	GOL	2	0
4	B	1553	DAO	5	0
3	B	1548	GOL	1	0
3	B	1550	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	164/170 (96%)	-0.41	3 (1%) 67 70	9, 19, 30, 44	4 (2%)
2	B	546/546 (100%)	0.21	37 (6%) 25 26	7, 26, 47, 61	9 (1%)
All	All	710/716 (99%)	0.07	40 (5%) 31 32	7, 24, 46, 61	13 (1%)

All (40) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	430	ILE	4.8
2	B	431	PRO	4.7
2	B	432	ASP	3.9
1	A	6	THR	3.6
2	B	429	GLY	3.4
1	A	167	ALA	3.3
1	A	169	LYS	3.2
2	B	408	PRO	3.1
2	B	344	GLY	3.0
2	B	347	SER	2.8
2	B	433	GLY	2.8
2	B	489	GLU	2.8
2	B	349	ALA	2.7
2	B	470	GLY	2.7
2	B	346	LYS	2.6
2	B	304	ARG	2.5
2	B	348	LEU	2.5
2	B	337	ARG	2.4
2	B	338	LEU	2.3
2	B	112	ALA	2.3
2	B	353	ALA	2.3
2	B	410	VAL	2.3
2	B	343	GLN	2.3
2	B	409	GLN	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	404	ALA	2.3
2	B	419	ALA	2.3
2	B	403	ILE	2.2
2	B	427	LYS	2.2
2	B	447	GLU	2.2
2	B	384	ASP	2.2
2	B	394	GLN	2.2
2	B	356	ALA	2.2
2	B	434	ALA	2.2
2	B	345	GLU	2.2
2	B	241	ARG	2.1
2	B	114	GLY	2.1
2	B	341	ASP	2.0
2	B	309	GLN	2.0
2	B	471	ASP	2.0
2	B	490	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	DAO	B	1553	14/14	0.78	0.19	35,38,41,41	0
3	GOL	A	1171	6/6	0.82	0.15	33,39,40,42	0
3	GOL	B	1547	6/6	0.86	0.20	51,53,54,55	0
3	GOL	B	1550	6/6	0.87	0.12	45,45,46,47	0
3	GOL	A	1172	6/6	0.88	0.12	35,40,42,44	0
3	GOL	B	1551	6/6	0.89	0.12	36,39,40,40	0
3	GOL	B	1549	6/6	0.89	0.16	24,35,37,40	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	GOL	A	1170	6/6	0.93	0.09	26,33,35,38	0
3	GOL	B	1548	6/6	0.93	0.15	19,27,29,33	0
4	DAO	B	1552	14/14	0.96	0.07	20,21,24,24	0

6.5 Other polymers [i](#)

There are no such residues in this entry.