



Full wwPDB EM Validation Report ⓘ

Dec 20, 2020 – 08:12 PM EST

PDB ID : 6X0U
EMDB ID : EMD-21984
Title : Structure of MZT1/GCP3-NHD and MZT1/GCP6-NHD in the gamma-TuRC
lumenal bridge
Authors : Wiczorek, M.; Huang, T.-L.; Urnavicius, L.; Hsia, K.-C.; Kapoor, T.M.
Deposited on : 2020-05-17
Resolution : 3.60 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.0.dev61
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.16


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	907	
2	D	1819	
3	A	82	
3	C	82	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3091 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Gamma-tubulin complex component 3.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	B	116	933	591	171	169	2	0	0

- Molecule 2 is a protein called Gamma-tubulin complex component 6.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	D	158	1220	771	209	232	8	0	0


- Molecule 3 is a protein called Mitotic-spindle organizing protein 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
3	A	65	484	299	85	96	4	0	0
3	C	59	454	281	79	90	4	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Gamma-tubulin complex component 3



THR	HIS	PHE	THR	GLN	GLY	ILE	VAL	GLN	GLN	PHE	THR	THR	THR	SER	SER	SER	SER	ASP	GLY
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

● Molecule 2: Gamma-tubulin complex component 6



PHE	SER	PHE	GLY	ASP	PRO	CYS	GLY	ASP	THR	THR	GLY	VAL	ALA	THR	VAL	GLY	VAL	THR	THR
GLU	ASP	GLY	PHE	GLM	SER	ALA	ASP	ASN	THR	ARG	THR	THR	THR	GLU	ALA	ASP	ASP	VAL	VAL
GLY	ARG	ASP	ALA	PHE	ASP	PHE	LYS	CYS	GLY	GLY	GLY	GLY	VAL	VAL	VAL	VAL	VAL	THR	THR
ALA	PHE	VAL	VAL	ARG	GLY	VAL	GLY	HIS	PRO	PRO	GLY	LEU	GLN	GLY	SER	SER	VAL	VAL	GLY
LEU	ARG	ARG	TYR	GLN	TYR	ARG	TYR	ARG	THR	THR	THR	THR	THR	PRO	GLY	THR	THR	THR	THR
PRO	THR	GLY	VAL	LEU	LEU	SER	TYR	LEU	GLY	THR	THR	THR	GLY	HIS	GLY	ASP	THR	THR	THR
VAL	ASN	HIS	TYR	LEU	VAL	ASP	VAL	TYR	THR	VAL	VAL	VAL	VAL	LEU	LEU	VAL	VAL	VAL	VAL
TRP	SER	ASP	VAL	PRO	VAL	LEU	VAL	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR
ALA	ARG	GLY	ALA	ASP	ARG	VAL	LEU	ASP	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR
ARG	GLY	ARG	ASP	GLY	ARG	ARG	GLY	ARG	GLY	GLY	GLY	GLY	GLY	GLY	GLY	GLY	GLY	GLY	GLY
LEU	GLY	ASP	GLY	HIS	ILE	GLN	VAL	VAL	VAL	VAL	VAL	VAL	VAL	VAL	VAL	VAL	VAL	VAL	VAL
ASP	GLY	TRP	ASN	PRO	THR	LEU	LEU	LEU	LEU	LEU	LEU	LEU	LEU	LEU	LEU	LEU	LEU	LEU	LEU
PRO	LEU	LEU	VAL	ALA	LEU	LEU	LEU	LEU	LEU	LEU	LEU	LEU	LEU	LEU	LEU	LEU	LEU	LEU	LEU
SER	LEU	GLY	SER	SER	GLY	GLY	GLY	GLY	GLY	GLY	GLY	GLY	GLY	GLY	GLY	GLY	GLY	GLY	GLY
THR	GLY	ASP	TYR	ALA	ILE	GLN	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	419138	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING ONLY	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	45	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	0.109	Depositor
Minimum map value	-0.074	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.002	Depositor
Recommended contour level	0.014	Depositor
Map size (\AA)	381.24802, 381.24802, 381.24802	wwPDB
Map dimensions	368, 368, 368	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (\AA)	1.036, 1.036, 1.036	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	B	0.27	0/948	0.42	0/1277
2	D	0.28	0/1235	0.43	0/1664
3	A	0.26	0/484	0.41	0/653
3	C	0.22	0/454	0.40	0/611
All	All	0.27	0/3121	0.42	0/4205

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	933	0	953	14	0
2	D	1220	0	1231	13	0
3	A	484	0	512	11	0
3	C	454	0	482	8	0
All	All	3091	0	3178	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:16:LEU:HD12	2:D:17:PRO:HD2	1.74	0.70
1:B:115:ALA:O	1:B:120:GLN:NE2	2.31	0.63
1:B:39:ILE:HD12	3:A:55:ILE:HD11	1.82	0.62
3:A:55:ILE:HD13	3:A:60:LEU:HD12	1.81	0.62
2:D:48:ASN:HD22	3:C:59:ALA:HB2	1.68	0.59
2:D:131:ASP:HB3	2:D:134:LEU:HD23	1.85	0.59
1:B:44:ALA:O	3:A:56:ASN:ND2	2.38	0.57
2:D:64:LEU:HD12	2:D:65:PRO:HD2	1.87	0.56
2:D:66:ALA:HB1	2:D:111:LEU:HD11	1.89	0.54
1:B:96:LEU:HD11	3:A:60:LEU:HD22	1.90	0.53
2:D:4:ILE:HG13	3:C:53:GLN:HE21	1.73	0.53
3:C:18:LEU:HA	3:C:21:VAL:HG12	1.91	0.51
2:D:88:ARG:NH2	3:C:30:GLU:OE1	2.43	0.51
3:C:59:ALA:O	3:C:62:SER:OG	2.28	0.51
1:B:45:PRO:HA	3:A:56:ASN:OD1	2.11	0.51
1:B:48:GLU:HG2	1:B:124:ARG:HG3	1.93	0.50
3:A:48:VAL:O	3:A:52:GLU:HG2	2.11	0.49
2:D:185:ALA:O	2:D:188:THR:HG23	2.12	0.49
1:B:54:VAL:O	1:B:58:ILE:HG13	2.15	0.47
2:D:69:LYS:O	2:D:73:LEU:HG	2.15	0.46
1:B:10:ASN:OD1	1:B:11:VAL:N	2.49	0.46
2:D:64:LEU:HD21	2:D:69:LYS:HE3	2.00	0.43
3:A:59:ALA:O	3:A:62:SER:OG	2.36	0.43
2:D:9:ASP:OD1	2:D:34:LYS:HE2	2.18	0.43
1:B:58:ILE:HG23	3:A:35:LEU:HD21	2.00	0.42
1:B:23:ARG:HH12	3:A:74:LEU:HD13	1.85	0.42
3:C:60:LEU:HA	3:C:63:VAL:HG22	2.02	0.41
1:B:21:LEU:HD23	3:A:71:THR:HB	2.01	0.41
2:D:83:GLY:N	2:D:84:PRO:HD2	2.35	0.41
2:D:11:LEU:HB2	3:C:46:ILE:HG21	2.03	0.41
1:B:28:VAL:O	1:B:28:VAL:HG13	2.21	0.41
1:B:67:ARG:NH2	3:A:34:ILE:O	2.52	0.40
1:B:54:VAL:HG21	1:B:91:TRP:CD1	2.56	0.40
3:C:32:SER:O	3:C:37:THR:OG1	2.34	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	112/907 (12%)	109 (97%)	3 (3%)	0	100	100
2	D	148/1819 (8%)	138 (93%)	9 (6%)	1 (1%)	22	61
3	A	63/82 (77%)	63 (100%)	0	0	100	100
3	C	57/82 (70%)	55 (96%)	2 (4%)	0	100	100
All	All	380/2890 (13%)	365 (96%)	14 (4%)	1 (0%)	44	75

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	D	134	LEU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	101/798 (13%)	101 (100%)	0	100	100
2	D	135/1546 (9%)	135 (100%)	0	100	100
3	A	53/62 (86%)	53 (100%)	0	100	100
3	C	53/62 (86%)	53 (100%)	0	100	100
All	All	342/2468 (14%)	342 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	30	GLN
1	B	65	GLN
2	D	48	ASN
2	D	172	HIS
2	D	176	GLN
3	C	53	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-21984. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 184



Y Index: 184




Z Index: 184


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 155



Y Index: 184



Z Index: 111

The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal surface views [i](#)


6.4.1 Primary map



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.014. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.5 Mask visualisation

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 241 nm^3 ; this corresponds to an approximate mass of 218 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum i



*Reported resolution corresponds to spatial frequency of 0.278 Å⁻¹


8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-21984 and PDB model 6X0U. Per-residue inclusion information can be found in section 3 on page 4.

9.1 Map-model overlay [i](#)



The images above show the 3D surface view of the map at the recommended contour level 0.014 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Atom inclusion [i](#)



At the recommended contour level, 74% of all backbone atoms, 63% of all non-hydrogen atoms, are inside the map.