



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2018 – 10:05 pm GMT

PDB ID : 4X1D
Title : Ytterbium-bound human serum transferrin
Authors : Wang, M.; Zhang, H.; Sun, H.
Deposited on : 2014-11-24
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtrriage (Phenix) : 1.13
EDS : trunk30967
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk30967


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	2792 (2.80-2.80)
Clashscore	122126	3209 (2.80-2.80)
Ramachandran outliers	120053	3158 (2.80-2.80)
Sidechain outliers	120020	3160 (2.80-2.80)
RSRZ outliers	108989	2726 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	679	 83% 13% ..
1	B	679	 83% 13% ..

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 10389 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serotransferrin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	658	5107	3216	885	961	45	0	0	0
1	B	658	5115	3221	888	961	45	0	1	0


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	429	VAL	ILE	See Sequence details	UNP P02787
B	429	VAL	ILE	See Sequence details	UNP P02787

- Molecule 2 is YTTERBIUM (III) ION (three-letter code: YB) (formula: Yb).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Yb	0	0
			1	1		
2	A	1	Total	Yb	0	0
			1	1		

- Molecule 3 is MALONATE ION (three-letter code: MLI) (formula: C₃H₂O₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			7	3	4		
3	B	1	Total	C	O	0	0
			7	3	4		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	78	Total 78	O 78	0	0
5	B	61	Total 61	O 61	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Serotransferrin



- Molecule 1: Serotransferrin



4 Data and refinement statistics i

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	156.50Å 136.86Å 107.84Å 90.00° 90.41° 90.00°	Depositor
Resolution (Å)	50.00 – 2.80 37.33 – 2.80	Depositor EDS
% Data completeness (in resolution range)	95.4 (50.00-2.80) 93.6 (37.33-2.80)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.36 (at 2.81Å)	Xtrriage
Refinement program	REFMAC 5.8.0073	Depositor
R, R_{free}	0.198 , 0.231 0.191 , 0.225	Depositor DCC
R_{free} test set	2627 reflections (4.92%)	wwPDB-VP
Wilson B-factor (Å ²)	51.1	Xtrriage
Anisotropy	0.071	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 23.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.397 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	10389	wwPDB-VP
Average B, all atoms (Å ²)	53.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.27% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, YB, MLI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.52	0/5222	0.69	1/7051 (0.0%)
1	B	0.52	0/5233	0.68	0/7065
All	All	0.52	0/10455	0.68	1/14116 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	23	ARG	NE-CZ-NH2	5.31	122.95	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	340	LYS	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5107	0	4945	47	0
1	B	5115	0	4958	53	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	7	0	2	1	0
3	B	7	0	2	1	0
4	A	6	0	8	0	0
4	B	6	0	8	0	0
5	A	78	0	0	3	0
5	B	61	0	0	0	0
All	All	10389	0	9923	100	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (100) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:628:ASP:OD2	1:B:632[B]:ARG:NH2	2.24	0.71
1:A:452:THR:HG22	1:A:517:TYR:HA	1.74	0.69
1:B:452:THR:HG22	1:B:517:TYR:HA	1.75	0.66
1:A:69:ASP:OD1	1:A:327:ARG:NH2	2.30	0.65
1:B:69:ASP:OD1	1:B:327:ARG:NH2	2.31	0.62
1:B:265:GLU:O	1:B:269:GLN:HG3	2.01	0.61
1:B:340:LYS:HB3	1:B:341:PRO:CD	2.31	0.60
1:A:265:GLU:O	1:A:269:GLN:HG3	2.01	0.59
1:B:671:LEU:O	1:B:675:THR:HG23	2.03	0.58
1:A:671:LEU:O	1:A:675:THR:HG23	2.03	0.58
1:B:628:ASP:OD2	1:B:632[B]:ARG:CZ	2.52	0.58
1:B:594:GLU:O	1:B:598:HIS:HB2	2.05	0.57
1:A:594:GLU:O	1:A:598:HIS:HB2	2.05	0.57
1:A:634:ASP:N	1:A:634:ASP:OD1	2.38	0.56
1:B:258:GLY:HA2	1:B:260:GLU:OE1	2.05	0.55
1:A:425:GLY:HA2	1:A:584:ASN:OD1	2.07	0.55
1:B:634:ASP:OD1	1:B:634:ASP:N	2.39	0.55
1:B:425:GLY:HA2	1:B:584:ASN:OD1	2.08	0.54
1:B:663:ARG:HD3	1:B:671:LEU:HD21	1.90	0.53
1:A:413:ASN:OD1	1:A:414:LYS:N	2.43	0.52
1:A:381:ILE:HA	1:A:386:ALA:O	2.09	0.52
1:A:538:VAL:HG11	1:A:571:VAL:HG21	1.92	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:23:ARG:HG3	1:B:37:VAL:O	2.09	0.52
1:A:96:TYR:HB2	1:A:207:HIS:HB3	1.92	0.52
1:B:381:ILE:HA	1:B:386:ALA:O	2.10	0.52
1:B:96:TYR:HB2	1:B:207:HIS:HB3	1.92	0.51
1:B:413:ASN:OD1	1:B:414:LYS:N	2.44	0.51
1:B:340:LYS:CG	1:B:341:PRO:HD3	2.40	0.51
1:B:538:VAL:HG11	1:B:571:VAL:HG21	1.92	0.51
1:A:514:TYR:CD2	1:A:523:CYS:HB2	2.45	0.50
1:B:514:TYR:CD2	1:B:523:CYS:HB2	2.45	0.50
1:A:23:ARG:HG3	1:A:37:VAL:O	2.12	0.50
1:B:164:GLY:HA3	1:B:171:CYS:SG	2.52	0.50
1:B:163:ASP:OD2	1:B:166:ASP:OD2	2.29	0.50
1:A:163:ASP:OD2	1:A:166:ASP:OD2	2.30	0.49
1:B:240:ASP:OD1	1:B:678:ARG:NH2	2.44	0.49
1:A:164:GLY:HA3	1:A:171:CYS:SG	2.53	0.49
1:A:79:PRO:HB3	1:A:250:THR:HG21	1.94	0.48
1:B:340:LYS:CB	1:B:341:PRO:CD	2.91	0.48
1:A:350:HIS:HB3	1:A:629:LEU:HD21	1.94	0.48
1:A:240:ASP:OD1	1:A:678:ARG:NH2	2.46	0.48
1:B:79:PRO:HB3	1:B:250:THR:HG21	1.96	0.48
1:B:254:ARG:NH1	1:B:257:GLY:O	2.47	0.47
1:A:254:ARG:NH1	1:A:257:GLY:O	2.47	0.47
1:A:377:CYS:HB3	1:A:389:MET:SD	2.54	0.47
1:B:340:LYS:HB3	1:B:341:PRO:HD3	1.96	0.47
1:A:374:THR:HG21	1:A:395:PHE:CD2	2.50	0.47
1:A:116:LYS:NZ	5:A:803:HOH:O	2.47	0.47
1:B:332:PRO:O	1:B:333:GLU:CG	2.63	0.47
1:B:374:THR:HG21	1:B:395:PHE:CD2	2.50	0.47
1:B:377:CYS:HB3	1:B:389:MET:SD	2.56	0.47
1:B:258:GLY:CA	1:B:260:GLU:OE1	2.63	0.46
1:A:26:MET:CE	1:A:274:PHE:CD2	2.99	0.46
1:A:324:ARG:NH1	5:A:857:HOH:O	2.49	0.46
1:B:350:HIS:HB3	1:B:629:LEU:HD21	1.97	0.46
1:B:26:MET:CE	1:B:274:PHE:CD2	2.99	0.45
1:A:26:MET:CE	1:A:274:PHE:CE2	3.00	0.45
1:B:524:LEU:HB2	1:B:531:ALA:HB2	1.99	0.45
1:A:276:LYS:HD3	5:A:828:HOH:O	2.17	0.45
1:A:524:LEU:HB2	1:A:531:ALA:HB2	1.98	0.45
1:B:135:LEU:O	1:B:136:TYR:C	2.56	0.45
1:A:539:PRO:HD3	1:A:571:VAL:HG11	2.00	0.44
1:B:585:HIS:CE1	3:B:702:MLI:O9	2.70	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:26:MET:CE	1:B:274:PHE:CE2	3.00	0.44
1:B:188:TYR:CE1	1:B:206:LYS:HG3	2.53	0.44
1:A:79:PRO:HB3	1:A:250:THR:CG2	2.47	0.44
1:B:332:PRO:C	1:B:333:GLU:HG3	2.38	0.44
1:B:574:TYR:CD1	1:B:575:ALA:N	2.86	0.44
1:B:538:VAL:HB	1:B:539:PRO:HD3	2.00	0.43
1:B:539:PRO:HD3	1:B:571:VAL:HG11	2.00	0.43
1:A:574:TYR:CD1	1:A:575:ALA:N	2.86	0.43
1:B:272:GLU:O	1:B:278:LYS:HE2	2.18	0.43
1:B:79:PRO:HB3	1:B:250:THR:CG2	2.48	0.43
1:A:427:PHE:HB2	1:A:535:HIS:HB3	2.00	0.43
1:B:382:MET:SD	1:B:402:CYS:HB3	2.58	0.43
1:A:135:LEU:O	1:A:136:TYR:C	2.57	0.43
1:A:382:MET:SD	1:A:402:CYS:HB3	2.58	0.43
1:A:272:GLU:O	1:A:278:LYS:HE2	2.19	0.43
1:B:460:TRP:O	1:B:464:MET:HB2	2.19	0.43
1:B:340:LYS:CB	1:B:341:PRO:HD3	2.48	0.42
1:A:460:TRP:O	1:A:464:MET:HB2	2.19	0.42
1:A:188:TYR:CE1	1:A:206:LYS:HG3	2.53	0.42
1:A:538:VAL:HB	1:A:539:PRO:HD3	2.01	0.42
1:A:272:GLU:O	1:A:278:LYS:CE	2.68	0.41
1:A:464:MET:HB3	1:A:476:PHE:CE2	2.55	0.41
1:B:272:GLU:O	1:B:278:LYS:CE	2.68	0.41
1:B:464:MET:HB3	1:B:476:PHE:CE2	2.55	0.41
1:A:383:ASN:OD1	1:A:383:ASN:C	2.59	0.41
1:A:426:TYR:CD1	1:A:582:ALA:HB3	2.55	0.41
1:B:195:LEU:HB2	1:B:203:ALA:HB2	2.03	0.41
1:B:471:ILE:HD12	1:B:473:HIS:NE2	2.35	0.41
1:A:471:ILE:HD12	1:A:473:HIS:NE2	2.36	0.41
1:B:400:GLY:HA3	1:B:647:TYR:CD2	2.55	0.41
1:A:585:HIS:CE1	3:A:702:MLI:O9	2.74	0.41
1:A:195:LEU:HB2	1:A:203:ALA:HB2	2.02	0.41
1:B:427:PHE:HB2	1:B:535:HIS:HB3	2.01	0.41
1:A:4:LYS:HE3	1:A:262:LEU:HD11	2.03	0.41
1:A:391:LEU:HD13	1:A:395:PHE:HB3	2.02	0.40
1:A:400:GLY:HA3	1:A:647:TYR:CD2	2.56	0.40
1:B:258:GLY:HA3	1:B:259:LYS:HA	1.79	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	650/679 (96%)	608 (94%)	36 (6%)	6 (1%)	19	50
1	B	651/679 (96%)	608 (93%)	38 (6%)	5 (1%)	21	53
All	All	1301/1358 (96%)	1216 (94%)	74 (6%)	11 (1%)	21	53

All (11) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	340	LYS
1	A	413	ASN
1	B	413	ASN
1	A	626	THR
1	B	626	THR
1	A	668	SER
1	B	668	SER
1	A	332	PRO
1	A	363	VAL
1	A	609	GLY
1	B	363	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	551/572 (96%)	534 (97%)	17 (3%)	43	77
1	B	552/572 (96%)	535 (97%)	17 (3%)	43	77

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	1103/1144 (96%)	1069 (97%)	34 (3%)	43 77

All (34) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	89	GLU
1	A	157	SER
1	A	249	HIS
1	A	260	GLU
1	A	291	LYS
1	A	327	ARG
1	A	340	LYS
1	A	365	LYS
1	A	489	LYS
1	A	506	CYS
1	A	555	ASN
1	A	565	ASP
1	A	568	ARG
1	A	571	VAL
1	A	624	SER
1	A	634	ASP
1	A	648	GLU
1	B	89	GLU
1	B	152	ASN
1	B	157	SER
1	B	249	HIS
1	B	291	LYS
1	B	327	ARG
1	B	365	LYS
1	B	437	SER
1	B	501	SER
1	B	506	CYS
1	B	565	ASP
1	B	568	ARG
1	B	571	VAL
1	B	624	SER
1	B	634	ASP
1	B	648	GLU
1	B	663	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	152	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	MLI	A	702	2	0,6,6	0.00	-	0,7,7	0.00	-
4	GOL	A	703	-	5,5,5	0.26	0	5,5,5	0.53	0
3	MLI	B	702	2	0,6,6	0.00	-	0,7,7	0.00	-
4	GOL	B	703	-	5,5,5	0.29	0	5,5,5	0.72	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MLI	A	702	2	-	0/0/4/4	0/0/0/0
4	GOL	A	703	-	-	0/4/4/4	0/0/0/0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MLI	B	702	2	-	0/0/4/4	0/0/0/0
4	GOL	B	703	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	702	MLI	1	0
3	B	702	MLI	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	658/679 (96%)	0.21	2 (0%) 93 92	25, 49, 90, 111	0
1	B	658/679 (96%)	0.22	3 (0%) 90 88	27, 48, 89, 108	0
All	All	1316/1358 (96%)	0.21	5 (0%) 92 90	25, 48, 89, 111	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	89	GLU	2.6
1	A	88	LYS	2.2
1	A	619	PHE	2.2
1	B	619	PHE	2.2
1	B	364	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	GOL	B	703	6/6	0.92	0.31	48,50,55,63	0
4	GOL	A	703	6/6	0.96	0.21	42,48,50,51	0
2	YB	B	701	1/1	0.99	0.22	47,47,47,47	0
3	MLI	A	702	7/7	0.99	0.20	33,36,38,38	0
2	YB	A	701	1/1	0.99	0.22	49,49,49,49	0
3	MLI	B	702	7/7	0.99	0.20	34,35,40,41	0

6.5 Other polymers [i](#)

There are no such residues in this entry.