

Full wwPDB X-ray Structure Validation Report (i)

Oct 28, 2024 - 03:06 PM JST

PDB ID	:	8X18
Title	:	Xanthomonas campestris pv. campestris OpgD mutant-D379N with beta-1,2-
		glucan
Authors	:	Motouchi, S.; Nakajima, M.
Deposited on	:	2023-11-06
Resolution	:	2.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R_{free}	164625	1763 (2.26-2.26)
Clashscore	180529	1919 (2.26-2.26)
Ramachandran outliers	177936	1884 (2.26-2.26)
Sidechain outliers	177891	1885 (2.26-2.26)
RSRZ outliers	164620	1763 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Qu	ality of chain	
1	А	541	3%	37%	6% • 7%
1	В	541	8%	6	8% • 9%
2	С	22	23%	73%	5%
3	D	11	55%	45%	
4	Е	10	50%	50%	



2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 9012 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
1	Λ	504	Total	С	Ν	0	\mathbf{S}	0	0	0
1		504	4020	2577	705	727	11	0	0	0
1	В	400	Total	С	Ν	0	S	0	0	0
ГБ	490	3927	2522	686	708	11	0	0	0	

• Molecule 1 is a protein called Glucans biosynthesis protein D.

Chain	Residue	Modelled	Actual	Comment	Reference
А	379	ASN	ASP	engineered mutation	UNP Q8P3C6
А	534	LEU	-	expression tag	UNP Q8P3C6
А	535	GLU	-	expression tag	UNP Q8P3C6
А	536	HIS	-	expression tag	UNP Q8P3C6
А	537	HIS	-	expression tag	UNP Q8P3C6
А	538	HIS	-	expression tag	UNP Q8P3C6
А	539	HIS	-	expression tag	UNP Q8P3C6
А	540	HIS	-	expression tag	UNP Q8P3C6
А	541	HIS	-	expression tag	UNP Q8P3C6
В	379	ASN	ASP	engineered mutation	UNP Q8P3C6
В	534	LEU	-	expression tag	UNP Q8P3C6
В	535	GLU	-	expression tag	UNP Q8P3C6
В	536	HIS	-	expression tag	UNP Q8P3C6
В	537	HIS	-	expression tag	UNP Q8P3C6
В	538	HIS	-	expression tag	UNP Q8P3C6
В	539	HIS	-	expression tag	UNP Q8P3C6
В	540	HIS	-	expression tag	UNP Q8P3C6
В	541	HIS	-	expression tag	UNP Q8P3C6

There are 18 discrepancies between the modelled and reference sequences:

• Molecule 2 is an oligosaccharide called beta-D-glucopyranose-(1-2)-beta-D



a-D-glucopyranose-(1-2)-beta-D-glucopyrano



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	Trace	
2	С	22	Total 243	C 132	0 111	0	0	0

• Molecule 3 is an oligosaccharide called beta-D-glucopyranose-(1-2)-beta-D



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
3	D	11	Total C O 122 66 56	0	0	0

• Molecule 4 is an oligosaccharide called beta-D-glucopyranose-(1-2)-beta-D



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	Trace	
4	Е	10	Total 111	C 60	O 51	0	0	0

• Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	А	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 6 3 3 \end{array}$	0	0
5	А	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 6 3 3 \end{array}$	0	0
5	А	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 6 3 3 \end{array}$	0	0
5	А	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 6 3 3 \end{array}$	0	0
5	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 6 & 3 & 3 \end{array}$	0	0
5	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 6 & 3 & 3 \end{array}$	0	0
5	В	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 6 3 3 \end{array}$	0	0
5	В	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 6 3 3 \end{array}$	0	0

• Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	А	298	Total O 298 298	0	0
6	В	243	Total O 243 243	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Glucans biosynthesis protein D

 $\label{eq:2} \bullet Molecule 2: beta-D-glucopyranose-(1-2)-beta-D-glucopyranos$





• Molecule 3: beta-D-glucopyranose-(1-2)-beta-D-

Chain D:	55%	45%
BGC1 BGC2 BGC3 BGC4 BGC5 BGC6 BGC7 BGC3 BGC1 BGC11		

 \bullet Molecule 4: beta-D-glucopyranose-(1-2)-beta

Chain E:	50%	50%
0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0		



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 65	Depositor
Cell constants	171.88Å 171.88Å 128.69Å	Deperitor
a, b, c, α , β , γ	90.00° 90.00° 120.00°	Depositor
$\mathbf{P}_{\text{assolution}}(\hat{\mathbf{A}})$	48.72 - 2.25	Depositor
Resolution (A)	48.72 - 2.25	EDS
% Data completeness	100.0 (48.72-2.25)	Depositor
(in resolution range)	$100.0 \ (48.72-2.25)$	EDS
R _{merge}	(Not available)	Depositor
R _{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$2.99 (at 2.24 \text{\AA})$	Xtriage
Refinement program	REFMAC 5.8.0405	Depositor
D D	0.173 , 0.176	Depositor
Π, Π_{free}	0.181 , 0.184	DCC
R_{free} test set	5213 reflections (5.10%)	wwPDB-VP
Wilson B-factor $(Å^2)$	28.7	Xtriage
Anisotropy	0.044	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.38 , 37.9	EDS
L-test for twinning ²	$< L >=0.50, < L^2>=0.33$	Xtriage
Estimated twinning fraction	0.025 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	9012	wwPDB-VP
Average B, all atoms $(Å^2)$	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 2.70% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: BGC, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond lengths		Bond angles	
1VIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.48	0/4146	0.82	1/5634~(0.0%)
1	В	0.50	0/4050	0.83	2/5504~(0.0%)
All	All	0.49	0/8196	0.82	3/11138~(0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	А	0	6
1	В	0	6
All	All	0	12

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	А	477	ARG	NE-CZ-NH2	-5.43	117.58	120.30
1	В	468	ARG	NE-CZ-NH2	5.25	122.93	120.30
1	В	326	ARG	NE-CZ-NH1	-5.19	117.70	120.30

There are no chirality outliers.

All (12) planarity outliers are listed belo

Mol	Chain	Res	Type	Group
1	А	237	ARG	Sidechain
1	А	326	ARG	Sidechain
1	А	36	ARG	Sidechain



Mol	Chain	Res	Type	Group
1	А	468	ARG	Sidechain
1	А	508	ARG	Sidechain
1	А	53	ARG	Sidechain
1	В	279	ARG	Sidechain
1	В	350	ARG	Sidechain
1	В	36	ARG	Sidechain
1	В	402	ARG	Sidechain
1	В	53	ARG	Sidechain
1	В	85	ARG	Sidechain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	4020	0	3870	21	0
1	В	3927	0	3774	34	0
2	С	243	0	201	3	0
3	D	122	0	102	0	0
4	Ε	111	0	93	0	0
5	А	36	0	48	9	0
5	В	12	0	16	2	0
6	А	298	0	0	0	0
6	B	243	0	0	3	0
All	All	9012	0	8104	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:433:ARG:HH11	1:B:433:ARG:HG3	1.41	0.86
1:A:104:HIS:HD2	1:A:105:LEU:O	1.70	0.74
5:A:605:GOL:O1	1:B:433:ARG:NH2	2.22	0.72
1:B:237:ARG:HD3	6:B:910:HOH:O	1.88	0.72
1:A:470:THR:HA	5:A:604:GOL:H2	1.74	0.69



	jour page	Interatomic	Clash	
Atom-1	Atom-2	distance (Å)	overlap (Å)	
1:A:104:HIS:HE1	1:A:128:TYR:OH	1.77	0.68	
1:A:523:THR:H	5:A:603:GOL:H12	1.60	0.67	
1:B:433:ARG:HH11	1:B:433:ARG:CG	2.11	0.63	
1:B:347:TYR:HA	1:B:350:ARG:HD2	1.81	0.63	
1:B:44:ASP:HB2	5:B:602:GOL:H31	1.79	0.62	
1:A:36:ARG:HG2	1:A:124:GLN:CG	2.32	0.60	
1:A:466:VAL:HG22	5:A:604:GOL:H11	1.84	0.58	
1:B:379:ASN:ND2	2:C:7:BGC:H6C1	2.19	0.57	
1:B:130:PRO:HB3	1:B:135:TYR:OH	2.06	0.55	
1:A:392:GLN:N	1:A:395:GLN:OE1	2.39	0.52	
1:A:379:ASN:O	5:A:601:GOL:H12	2.10	0.51	
1:B:508:ARG:HD3	6:B:704:HOH:O	2.09	0.51	
1:B:279:ARG:HD3	6:B:906:HOH:O	2.11	0.50	
1:A:312:PRO:HA	5:A:603:GOL:C3	2.42	0.50	
1:B:379:ASN:HD22	2:C:7:BGC:H6C1	1.76	0.49	
1:A:36:ARG:HG2	1:A:124:GLN:HG2	1.94	0.49	
1:B:266:GLY:N	1:B:267:PRO:CD	2.76	0.48	
1:B:75:ASN:H	1:B:78:GLN:NE2	2.13	0.47	
1:B:80:GLN:C	1:B:82:ILE:N	2.66	0.46	
1:B:45:TYR:HB3	5:B:602:GOL:H2	1.98	0.46	
1:A:284:ASP:HB2	5:A:605:GOL:O3	2.16	0.46	
1:A:463:VAL:O	1:A:505:LEU:HA	2.16	0.46	
1:B:32:VAL:HG23	1:B:33:GLY:O	2.16	0.46	
1:B:263:LEU:HD22	1:B:391:PRO:HG3	1.98	0.46	
1:B:30:LYS:HD2	1:B:31:ALA:H	1.82	0.45	
1:B:74:LEU:N	1:B:78:GLN:CD	2.70	0.45	
1:B:211:LEU:HD12	1:B:211:LEU:N	2.32	0.44	
1:A:260:ILE:HB	1:A:391:PRO:HG2	1.99	0.44	
1:A:471:THR:HA	1:A:490:ASP:O	2.18	0.44	
1:B:436:PHE:CD2	1:B:530:ARG:HD3	2.52	0.44	
1:A:312:PRO:HA	5:A:603:GOL:H32	2.00	0.44	
1:B:355:VAL:HG22	1:B:403:LEU:CD2	2.47	0.44	
1:B:107:LEU:HD23	1:B:108:TYR:H	1.83	0.44	
1:A:237:ARG:HH12	1:A:239:ALA:HB2	1.82	0.43	
1:B:375:ASP:OD1	1:B:377:THR:OG1	2.25	0.43	
5:A:605:GOL:C1	1:B:433:ARG:NH2	2.82	0.43	
1:B:224:TYR:CE2	1:B:237:ARG:HG3	2.53	0.42	
1:B:80:GLN:N	1:B:80:GLN:OE1	2.52	0.42	
1:B:468:ARG:CZ	1:B:499:GLN:HB2	2.49	0.42	
2:C:4:BGC:H3	2:C:6:BGC:H6C2	2.02	0.42	
1:A:237:ARG:NH1	1:A:239:ALA:HB2	2.35	0.42	



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:82:ILE:H	1:B:82:ILE:HG13	1.75	0.42
1:B:281:MET:O	1:B:281:MET:HG3	2.19	0.42
1:B:35:ARG:HG2	1:B:127:ALA:HB3	2.02	0.41
1:B:426:GLY:HA2	1:B:439:ARG:HB2	2.03	0.41
1:A:436:PHE:CD2	1:A:530:ARG:HD3	2.56	0.41
1:B:88:HIS:CE1	1:B:131:ALA:O	2.74	0.41
1:A:315:LEU:HD12	1:A:371:ILE:HG12	2.03	0.40
1:A:355:VAL:HG22	1:A:403:LEU:CD2	2.52	0.40
1:B:433:ARG:CG	1:B:433:ARG:NH1	2.76	0.40
1:A:455:ASP:HA	1:A:456:PRO:HD3	1.93	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Analysed Favoured Allowed Outliers		Outliers	Percentiles		
1	А	502/541~(93%)	486 (97%)	16 (3%)	0	100 100		
1	В	484/541~(90%)	468 (97%)	16 (3%)	0	100 100		
All	All	986/1082~(91%)	954 (97%)	32 (3%)	0	100 100		

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	А	410/436~(94%)	403~(98%)	7 (2%)	56 66		
1	В	401/436~(92%)	397~(99%)	4 (1%)	73 80		
All	All	811/872 (93%)	800 (99%)	11 (1%)	62 72		

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	36	ARG
1	А	237	ARG
1	А	315	LEU
1	А	423	THR
1	А	457	LYS
1	А	468	ARG
1	А	508	ARG
1	В	82	ILE
1	В	107	LEU
1	В	157	ASN
1	В	164	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	104	HIS
1	А	379	ASN
1	В	78	GLN
1	В	183	GLN
1	В	379	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

43 monosaccharides are modelled in this entry.



8X18

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Turne	Chain	Dec	Tink	Bo	ond leng	$_{\rm sths}$	В	ond ang	les
	туре	Chain	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	BGC	С	1	2	12,12,12	0.22	0	$17,\!17,\!17$	0.50	0
2	BGC	С	10	2	11,11,12	0.27	0	$15,\!15,\!17$	0.98	1 (6%)
2	BGC	С	11	2	11,11,12	0.32	0	$15,\!15,\!17$	0.68	1 (6%)
2	BGC	С	12	2	11,11,12	0.20	0	$15,\!15,\!17$	0.74	1 (6%)
2	BGC	С	13	2	11,11,12	0.25	0	$15,\!15,\!17$	1.01	1 (6%)
2	BGC	С	14	2	11,11,12	0.37	0	$15,\!15,\!17$	1.22	1 (6%)
2	BGC	С	15	2	11,11,12	0.18	0	$15,\!15,\!17$	0.82	1 (6%)
2	BGC	С	16	2	11,11,12	0.32	0	$15,\!15,\!17$	1.25	1 (6%)
2	BGC	С	17	2	11,11,12	0.27	0	$15,\!15,\!17$	0.96	1 (6%)
2	BGC	С	18	2	11,11,12	0.25	0	$15,\!15,\!17$	0.59	0
2	BGC	С	19	2	11,11,12	0.28	0	15,15,17	0.64	0
2	BGC	С	2	2	11,11,12	0.32	0	15,15,17	0.83	1 (6%)
2	BGC	С	20	2	11,11,12	0.30	0	15,15,17	0.63	0
2	BGC	С	21	2	11,11,12	0.88	0	$15,\!15,\!17$	0.87	1 (6%)
2	BGC	С	22	2	11,11,12	0.69	0	$15,\!15,\!17$	1.20	2 (13%)
2	BGC	С	3	2	11,11,12	0.28	0	$15,\!15,\!17$	0.74	1 (6%)
2	BGC	С	4	2	11,11,12	0.22	0	15,15,17	0.54	0
2	BGC	С	5	2	11,11,12	0.26	0	$15,\!15,\!17$	0.98	1 (6%)
2	BGC	С	6	2	11,11,12	0.24	0	$15,\!15,\!17$	0.66	0
2	BGC	С	7	2	11,11,12	0.72	0	$15,\!15,\!17$	1.55	4 (26%)
2	BGC	С	8	2	11,11,12	0.34	0	15,15,17	0.47	0
2	BGC	С	9	2	11,11,12	0.31	0	$15,\!15,\!17$	1.59	2 (13%)
3	BGC	D	1	3	12,12,12	0.22	0	17,17,17	0.47	0
3	BGC	D	10	3	11,11,12	0.41	0	15,15,17	0.72	0
3	BGC	D	11	3	11,11,12	0.28	0	$15,\!15,\!17$	0.84	1 (6%)
3	BGC	D	2	3	11,11,12	0.31	0	15,15,17	0.52	0
3	BGC	D	3	3	11,11,12	0.30	0	$15,\!15,\!17$	0.93	1 (6%)
3	BGC	D	4	3	11,11,12	0.46	0	15,15,17	1.06	1 (6%)
3	BGC	D	5	3	11,11,12	0.36	0	15,15,17	0.75	1 (6%)
3	BGC	D	6	3	11,11,12	0.35	0	$15,\!15,\!17$	0.63	0
3	BGC	D	7	3	11,11,12	0.31	0	$15,\!15,\!17$	0.92	1 (6%)



Mal	Turne	Chain	Dec	Tink	Bond lengt	$_{\rm sths}$	B	ond ang	gles	
10101	туре	Unain	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	BGC	D	8	3	11,11,12	0.35	0	15,15,17	0.71	0
3	BGC	D	9	3	11,11,12	0.47	0	15,15,17	0.75	0
4	BGC	Е	1	4	12,12,12	0.17	0	17,17,17	0.27	0
4	BGC	Е	10	4	11,11,12	0.23	0	15,15,17	0.68	0
4	BGC	Е	2	4	11,11,12	0.27	0	$15,\!15,\!17$	1.07	1 (6%)
4	BGC	E	3	4	11,11,12	0.37	0	15,15,17	0.97	1 (6%)
4	BGC	E	4	4	11,11,12	0.36	0	$15,\!15,\!17$	0.81	1 (6%)
4	BGC	Е	5	4	11,11,12	0.38	0	15,15,17	0.67	0
4	BGC	E	6	4	11,11,12	0.36	0	15,15,17	0.65	0
4	BGC	E	7	4	$11,\!11,\!12$	0.29	0	$15,\!15,\!17$	0.66	0
4	BGC	Е	8	4	$11,\!11,\!12$	0.37	0	$15,\!15,\!17$	0.78	1 (6%)
4	BGC	Е	9	4	11,11,12	0.37	0	15,15,17	0.76	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BGC	С	1	2	-	0/2/22/22	0/1/1/1
2	BGC	С	10	2	-	0/2/19/22	0/1/1/1
2	BGC	С	11	2	-	0/2/19/22	0/1/1/1
2	BGC	С	12	2	-	0/2/19/22	0/1/1/1
2	BGC	С	13	2	-	0/2/19/22	0/1/1/1
2	BGC	С	14	2	-	0/2/19/22	0/1/1/1
2	BGC	С	15	2	-	0/2/19/22	0/1/1/1
2	BGC	С	16	2	-	0/2/19/22	0/1/1/1
2	BGC	С	17	2	-	0/2/19/22	0/1/1/1
2	BGC	С	18	2	-	0/2/19/22	0/1/1/1
2	BGC	С	19	2	-	0/2/19/22	0/1/1/1
2	BGC	С	2	2	-	2/2/19/22	0/1/1/1
2	BGC	С	20	2	-	0/2/19/22	0/1/1/1
2	BGC	С	21	2	-	0/2/19/22	0/1/1/1
2	BGC	С	22	2	-	0/2/19/22	0/1/1/1
2	BGC	С	3	2	-	0/2/19/22	0/1/1/1
2	BGC	С	4	2	-	0/2/19/22	0/1/1/1
2	BGC	С	5	2	-	0/2/19/22	0/1/1/1
2	BGC	С	6	2	-	0/2/19/22	0/1/1/1
2	BGC	С	7	2	-	0/2/19/22	0/1/1/1
2	BGC	С	8	2	-	0/2/19/22	0/1/1/1
2	BGC	С	9	2	-	0/2/19/22	0/1/1/1



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	BGC	D	1	3	-	0/2/22/22	0/1/1/1
3	BGC	D	10	3	-	0/2/19/22	0/1/1/1
3	BGC	D	11	3	-	0/2/19/22	0/1/1/1
3	BGC	D	2	3	-	0/2/19/22	0/1/1/1
3	BGC	D	3	3	-	0/2/19/22	0/1/1/1
3	BGC	D	4	3	-	0/2/19/22	0/1/1/1
3	BGC	D	5	3	-	0/2/19/22	0/1/1/1
3	BGC	D	6	3	-	0/2/19/22	0/1/1/1
3	BGC	D	7	3	-	0/2/19/22	0/1/1/1
3	BGC	D	8	3	-	0/2/19/22	0/1/1/1
3	BGC	D	9	3	-	2/2/19/22	0/1/1/1
4	BGC	Е	1	4	-	0/2/22/22	0/1/1/1
4	BGC	Е	10	4	-	0/2/19/22	0/1/1/1
4	BGC	Е	2	4	-	0/2/19/22	0/1/1/1
4	BGC	Е	3	4	-	0/2/19/22	0/1/1/1
4	BGC	Е	4	4	-	0/2/19/22	0/1/1/1
4	BGC	Е	5	4	-	0/2/19/22	0/1/1/1
4	BGC	Е	6	4	-	2/2/19/22	0/1/1/1
4	BGC	Е	7	4	-	0/2/19/22	0/1/1/1
4	BGC	Е	8	4	_	1/2/19/22	0/1/1/1
4	BGC	Е	9	4	_	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (30) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	С	9	BGC	C1-C2-C3	4.92	115.72	109.67
2	С	7	BGC	C1-O5-C5	4.08	117.72	112.19
2	С	16	BGC	C1-C2-C3	3.81	114.35	109.67
4	Е	2	BGC	C1-C2-C3	3.56	114.04	109.67
2	С	14	BGC	C1-C2-C3	3.47	113.93	109.67
3	D	4	BGC	C1-C2-C3	3.28	113.69	109.67
3	D	3	BGC	C1-C2-C3	3.15	113.54	109.67
2	С	13	BGC	C1-C2-C3	3.01	113.36	109.67
2	С	22	BGC	C1-O5-C5	2.84	116.04	112.19
2	С	10	BGC	C1-C2-C3	2.83	113.15	109.67
4	Е	3	BGC	C1-C2-C3	2.79	113.10	109.67
3	D	7	BGC	C1-C2-C3	2.73	113.02	109.67
2	С	17	BGC	C1-C2-C3	2.63	112.90	109.67
2	С	15	BGC	C1-C2-C3	2.62	112.89	109.67
2	С	9	BGC	O5-C1-C2	2.57	114.74	110.77



Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	С	7	BGC	C1-C2-C3	2.53	112.78	109.67
4	Е	4	BGC	C1-C2-C3	2.40	112.62	109.67
2	С	21	BGC	C1-C2-C3	2.40	112.61	109.67
4	Е	9	BGC	C1-C2-C3	2.34	112.54	109.67
2	С	11	BGC	C1-O5-C5	2.27	115.26	112.19
2	С	7	BGC	O5-C1-C2	2.22	114.20	110.77
2	С	5	BGC	C1-C2-C3	2.18	112.34	109.67
3	D	5	BGC	C1-C2-C3	2.12	112.28	109.67
4	Е	8	BGC	C1-C2-C3	2.08	112.22	109.67
2	С	7	BGC	O2-C2-C3	-2.04	106.06	110.14
2	С	12	BGC	C1-C2-C3	2.03	112.16	109.67
2	С	2	BGC	C1-O5-C5	2.03	114.94	112.19
2	С	22	BGC	C3-C4-C5	2.03	113.85	110.24
2	С	3	BGC	C1-O5-C5	2.02	114.93	112.19
3	D	11	BGC	C1-C2-C3	2.00	112.13	109.67

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	С	2	BGC	O5-C5-C6-O6
2	С	2	BGC	C4-C5-C6-O6
3	D	9	BGC	C4-C5-C6-O6
3	D	9	BGC	O5-C5-C6-O6
4	Е	6	BGC	C4-C5-C6-O6
4	Е	6	BGC	O5-C5-C6-O6
4	Е	8	BGC	C4-C5-C6-O6

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	С	4	BGC	1	0
2	С	6	BGC	1	0
2	С	7	BGC	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.











5.6 Ligand geometry (i)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	T	c Chain Dar I		T : 1-	B	Bond lengths			Bond angles		
	Type	Chain	Res	LINK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2	
5	GOL	А	606	-	5,5,5	0.15	0	$5,\!5,\!5$	0.45	0	
5	GOL	А	605	-	5,5,5	0.13	0	$5,\!5,\!5$	0.41	0	
5	GOL	А	601	-	5,5,5	0.09	0	$5,\!5,\!5$	0.35	0	
5	GOL	A	602	-	5,5,5	0.21	0	$5,\!5,\!5$	0.64	0	
5	GOL	В	601	-	5,5,5	0.12	0	$5,\!5,\!5$	0.42	0	
5	GOL	В	602	-	5,5,5	0.34	0	$5,\!5,\!5$	0.84	0	
5	GOL	А	603	-	5,5,5	0.23	0	$5,\!5,\!5$	0.51	0	
5	GOL	A	604	-	5,5,5	0.24	0	$5,\!5,\!5$	1.06	0	



In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	А	606	-	-	4/4/4/4	-
5	GOL	А	605	-	-	2/4/4/4	-
5	GOL	А	601	-	-	4/4/4/4	-
5	GOL	А	602	-	-	0/4/4/4	-
5	GOL	В	601	-	-	2/4/4/4	-
5	GOL	В	602	-	-	2/4/4/4	-
5	GOL	А	603	-	-	2/4/4/4	-
5	GOL	А	604	-	-	4/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (20) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	А	601	GOL	O1-C1-C2-C3
5	А	603	GOL	O1-C1-C2-C3
5	А	604	GOL	O1-C1-C2-C3
5	А	605	GOL	C1-C2-C3-O3
5	А	605	GOL	O2-C2-C3-O3
5	А	606	GOL	O1-C1-C2-C3
5	В	601	GOL	O1-C1-C2-C3
5	В	602	GOL	C1-C2-C3-O3
5	А	601	GOL	O2-C2-C3-O3
5	А	604	GOL	O1-C1-C2-O2
5	А	601	GOL	C1-C2-C3-O3
5	А	603	GOL	O1-C1-C2-O2
5	А	606	GOL	O1-C1-C2-O2
5	А	604	GOL	O2-C2-C3-O3
5	В	602	GOL	O2-C2-C3-O3
5	А	601	GOL	O1-C1-C2-O2
5	А	606	GOL	O2-C2-C3-O3
5	В	601	GOL	O1-C1-C2-O2
5	А	604	GOL	C1-C2-C3-O3
5	А	606	GOL	C1-C2-C3-O3



There are no ring outliers.

5 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	А	605	GOL	3	0
5	А	601	GOL	1	0
5	В	602	GOL	2	0
5	А	603	GOL	3	0
5	А	604	GOL	2	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	504/541~(93%)	-0.39	17 (3%) 48 49	19, 27, 54, 81	0
1	В	490/541~(90%)	-0.07	43 (8%) 17 17	18, 28, 73, 107	0
All	All	994/1082~(91%)	-0.23	60 (6%) 29 28	18, 27, 65, 107	0

All (60) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	74	LEU	16.0
1	В	107	LEU	8.0
1	В	135	TYR	5.9
1	В	68	PRO	5.8
1	В	30	LYS	5.6
1	В	108	TYR	5.0
1	В	81	SER	4.3
1	А	509	ALA	4.1
1	В	75	ASN	4.1
1	В	36	ARG	4.0
1	А	36	ARG	3.9
1	А	459	LYS	3.9
1	В	78	GLN	3.9
1	В	110	HIS	3.9
1	А	432	LYS	3.8
1	А	30	LYS	3.6
1	В	79	TYR	3.6
1	В	83	ARG	3.6
1	В	80	GLN	3.4
1	В	430	GLY	3.4
1	В	39	GLN	3.3
1	В	76	TRP	3.3
1	В	160	LYS	3.3
1	В	457	LYS	3.3



8X18	3
------	---

Mol	Chain	Res	Type	RSRZ
1	В	67	LEU	3.2
1	В	454	LYS	3.2
1	А	457	LYS	3.1
1	В	105	LEU	3.1
1	В	66	VAL	3.1
1	А	160	LYS	3.1
1	В	432	LYS	3.1
1	А	454	LYS	3.0
1	А	512	LYS	3.0
1	В	82	ILE	3.0
1	В	145	LEU	3.0
1	В	95	ASN	3.0
1	В	31	ALA	3.0
1	В	131	ALA	2.9
1	А	531	LYS	2.9
1	В	64	LYS	2.9
1	В	531	LYS	2.7
1	В	456	PRO	2.7
1	А	39	GLN	2.7
1	В	33	GLY	2.6
1	В	164	ARG	2.6
1	В	459	LYS	2.6
1	А	456	PRO	2.6
1	В	77	ASP	2.5
1	А	468	ARG	2.4
1	A	395	GLN	2.3
1	В	32	VAL	2.3
1	А	143	LYS	2.3
1	В	279	ARG	2.3
1	В	496	GLU	2.2
1	В	87	ASP	2.2
1	А	511	GLY	2.2
1	В	468	ARG	2.2
1	А	142	GLY	2.1
1	В	147	LYS	2.1
1	В	281	MET	2.1

Continued from previous page...

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



6.3 Carbohydrates (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(Å^2)$	Q<0.9
3	BGC	D	1	12/12	0.82	0.17	$45,\!54,\!65,\!74$	0
2	BGC	С	2	11/12	0.84	0.15	41,53,67,70	0
4	BGC	Е	3	11/12	0.84	0.14	35,49,51,53	0
2	BGC	С	13	11/12	0.85	0.13	42,48,53,56	0
2	BGC	С	1	12/12	0.87	0.11	37,45,48,48	0
3	BGC	D	4	11/12	0.88	0.13	39,47,51,53	0
2	BGC	С	14	11/12	0.88	0.12	43,46,55,59	0
3	BGC	D	6	11/12	0.89	0.12	37,41,46,46	0
4	BGC	Е	6	11/12	0.89	0.12	33,36,43,49	0
2	BGC	С	19	11/12	0.90	0.13	40,48,52,52	0
3	BGC	D	7	11/12	0.90	0.12	40,46,55,58	0
2	BGC	С	16	11/12	0.91	0.11	39,44,49,50	0
4	BGC	Е	1	12/12	0.91	0.11	$35,\!41,\!50,\!56$	0
3	BGC	D	5	11/12	0.92	0.10	29,32,35,39	0
2	BGC	С	18	11/12	0.92	0.10	31,37,41,43	0
4	BGC	Е	5	11/12	0.92	0.10	$35,\!37,\!43,\!46$	0
2	BGC	С	21	11/12	0.92	0.10	45,48,52,54	0
2	BGC	С	15	11/12	0.93	0.09	41,44,45,46	0
2	BGC	С	20	11/12	0.93	0.09	30,35,40,40	0
2	BGC	С	5	11/12	0.93	0.09	$29,\!32,\!37,\!37$	0
3	BGC	D	8	11/12	0.93	0.09	26,32,35,36	0
3	BGC	D	9	11/12	0.93	0.09	25,29,34,34	0
2	BGC	С	17	11/12	0.93	0.10	34,41,42,43	0
3	BGC	D	2	11/12	0.93	0.09	34,36,44,45	0
4	BGC	Е	4	11/12	0.93	0.10	33,36,38,40	0
3	BGC	D	3	11/12	0.93	0.09	32,36,41,42	0
2	BGC	С	12	11/12	0.93	0.09	41,44,47,49	0
3	BGC	D	11	11/12	0.94	0.10	28,32,36,37	0
2	BGC	С	11	11/12	0.94	0.08	$29,\!36,\!40,\!42$	0
2	BGC	С	4	11/12	0.94	0.08	30,34,35,37	0
2	BGC	С	7	11/12	0.94	0.09	$26,\!29,\!33,\!36$	0
2	BGC	С	8	11/12	0.94	0.08	31,37,38,39	0
2	BGC	C	10	11/12	0.94	0.08	$27,\!33,\!34,\!34$	0
4	BGC	Е	10	11/12	0.94	0.08	26,29,32,33	0
4	BGC	Е	2	11/12	0.95	0.07	28,30,32,34	0
4	BGC	Е	7	11/12	0.95	0.08	26,31,34,39	0
4	BGC	Е	8	11/12	0.95	0.08	24,28,33,36	0



Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	BGC	Е	9	11/12	0.95	0.07	$26,\!29,\!31,\!32$	0
2	BGC	С	22	11/12	0.95	0.08	41,46,49,49	0
3	BGC	D	10	11/12	0.96	0.07	29,30,33,33	0
2	BGC	С	6	11/12	0.96	0.07	24,30,33,33	0
2	BGC	С	9	11/12	0.96	0.07	$25,\!31,\!32,\!33$	0
2	BGC	С	3	11/12	0.97	0.06	21,24,30,30	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.









6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathrm{\AA}^2)$	Q<0.9
5	GOL	А	601	6/6	0.77	0.21	$30,\!34,\!41,\!42$	0
5	GOL	В	602	6/6	0.78	0.21	41,46,47,56	0
5	GOL	А	603	6/6	0.88	0.20	42,46,63,64	0
5	GOL	А	606	6/6	0.91	0.16	55,64,68,71	0
5	GOL	А	605	6/6	0.92	0.13	39,43,49,50	0
5	GOL	В	601	6/6	0.94	0.11	$26,\!35,\!42,\!51$	0
5	GOL	А	604	6/6	0.94	0.10	30,37,38,40	0
5	GOL	А	602	6/6	0.95	0.08	26,30,30,34	0

6.5 Other polymers (i)

There are no such residues in this entry.

