



wwPDB X-ray Structure Validation Summary Report ⓘ

Jul 10, 2024 – 10:11 AM JST

PDB ID : 8X22
Title : HIV-1 reverse transcriptase mutant Q151M/Y115F/F116Y/L74V:DNA:dGT
P ternary complex
Authors : Yasutake, Y.; Hattori, S.I.; Mitsuya, H.
Deposited on : 2023-11-09
Resolution : 2.31 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.37.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.31 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5974 (2.34-2.30)
Clashscore	141614	6604 (2.34-2.30)
Ramachandran outliers	138981	6523 (2.34-2.30)
Sidechain outliers	138945	6523 (2.34-2.30)
RSRZ outliers	127900	5855 (2.34-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	557	 7% 82% 17% .
1	C	557	 11% 83% 15% ..
2	B	444	 15% 75% 15% . 9%
2	D	444	 6% 80% 11% . 9%
3	E	38	 79% 8% 5% 8%
3	F	38	 3% 71% 24% 5%

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 17744 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pol protein (Fragment).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	553	4497	2910	750	829	8	0	0	0
1	C	553	4505	2915	751	830	9	0	1	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MET	-	initiating methionine	UNP D3XFN5
A	0	VAL	-	expression tag	UNP D3XFN5
A	74	VAL	LEU	engineered mutation	UNP D3XFN5
A	115	PHE	TYR	engineered mutation	UNP D3XFN5
A	116	TYR	PHE	engineered mutation	UNP D3XFN5
A	151	MET	GLN	engineered mutation	UNP D3XFN5
A	162	SER	CYS	engineered mutation	UNP D3XFN5
A	280	SER	CYS	engineered mutation	UNP D3XFN5
C	-1	MET	-	initiating methionine	UNP D3XFN5
C	0	VAL	-	expression tag	UNP D3XFN5
C	74	VAL	LEU	engineered mutation	UNP D3XFN5
C	115	PHE	TYR	engineered mutation	UNP D3XFN5
C	116	TYR	PHE	engineered mutation	UNP D3XFN5
C	151	MET	GLN	engineered mutation	UNP D3XFN5
C	162	SER	CYS	engineered mutation	UNP D3XFN5
C	280	SER	CYS	engineered mutation	UNP D3XFN5

- Molecule 2 is a protein called HIV-1 RT p51 subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	406	3347	2178	557	606	6	0	0	0
2	D	406	3347	2178	557	606	6	0	0	0

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-15	MET	-	expression tag	UNP P12497
B	-14	ALA	-	expression tag	UNP P12497
B	-13	HIS	-	expression tag	UNP P12497
B	-12	HIS	-	expression tag	UNP P12497
B	-11	HIS	-	expression tag	UNP P12497
B	-10	HIS	-	expression tag	UNP P12497
B	-9	HIS	-	expression tag	UNP P12497
B	-8	HIS	-	expression tag	UNP P12497
B	-7	ALA	-	expression tag	UNP P12497
B	-6	LEU	-	expression tag	UNP P12497
B	-5	GLU	-	expression tag	UNP P12497
B	-4	VAL	-	expression tag	UNP P12497
B	-3	LEU	-	expression tag	UNP P12497
B	-2	PHE	-	expression tag	UNP P12497
B	-1	GLN	-	expression tag	UNP P12497
B	0	GLY	-	expression tag	UNP P12497
B	162	SER	CYS	engineered mutation	UNP P12497
B	280	SER	CYS	engineered mutation	UNP P12497
D	-15	MET	-	expression tag	UNP P12497
D	-14	ALA	-	expression tag	UNP P12497
D	-13	HIS	-	expression tag	UNP P12497
D	-12	HIS	-	expression tag	UNP P12497
D	-11	HIS	-	expression tag	UNP P12497
D	-10	HIS	-	expression tag	UNP P12497
D	-9	HIS	-	expression tag	UNP P12497
D	-8	HIS	-	expression tag	UNP P12497
D	-7	ALA	-	expression tag	UNP P12497
D	-6	LEU	-	expression tag	UNP P12497
D	-5	GLU	-	expression tag	UNP P12497
D	-4	VAL	-	expression tag	UNP P12497
D	-3	LEU	-	expression tag	UNP P12497
D	-2	PHE	-	expression tag	UNP P12497
D	-1	GLN	-	expression tag	UNP P12497
D	0	GLY	-	expression tag	UNP P12497
D	162	SER	CYS	engineered mutation	UNP P12497
D	280	SER	CYS	engineered mutation	UNP P12497

- Molecule 3 is a DNA chain called DNA/RNA (38-MER).


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
3	E	35	718	339	128	216	35	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
3	F	38	777	369	140	231	37	0	0	0

- Molecule 4 is 2'-DEOXYGUANOSINE-5'-TRIPHOSPHATE (three-letter code: DGT) (formula: $C_{10}H_{16}N_5O_{13}P_3$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
4	A	1	31	10	5	13	3	0	0
4	C	1	31	10	5	13	3	0	0

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
5	A	1	1	1	0	0
5	C	1	1	1	0	0

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 6 3 3	0	0
6	A	1	Total C O 6 3 3	0	0
6	B	1	Total C O 6 3 3	0	0
6	C	1	Total C O 6 3 3	0	0
6	D	1	Total C O 6 3 3	0	0
6	D	1	Total C O 6 3 3	0	0

- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	126	Total O 126 126	0	0
7	B	52	Total O 52 52	0	0
7	E	40	Total O 40 40	0	0
7	C	115	Total O 115 115	0	0
7	D	86	Total O 86 86	0	0
7	F	34	Total O 34 34	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Pol protein (Fragment)



- Molecule 1: Pol protein (Fragment)



- Molecule 2: HIV-1 RT p51 subunit

4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	285.23Å 285.23Å 96.08Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	48.81 – 2.31 48.81 – 2.31	Depositor EDS
% Data completeness (in resolution range)	100.0 (48.81-2.31) 100.0 (48.81-2.31)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.24 (at 2.32Å)	Xtrriage
Refinement program	PHENIX (1.14_3260: ???)	Depositor
R, R_{free}	0.182 , 0.219 0.182 , 0.219	Depositor DCC
R_{free} test set	6481 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	48.0	Xtrriage
Anisotropy	0.219	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 47.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.008 for h,-h-k,-l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	17744	wwPDB-VP
Average B, all atoms (Å ²)	65.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.31% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, DGT, OMC, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.41	0/4615	0.55	0/6267
1	C	0.44	4/4623 (0.1%)	0.55	2/6277 (0.0%)
2	B	0.47	3/3441 (0.1%)	0.60	3/4673 (0.1%)
2	D	0.38	0/3441	0.52	0/4673
3	E	0.76	0/756	0.97	0/1165
3	F	0.86	2/823 (0.2%)	0.95	0/1269
All	All	0.48	9/17699 (0.1%)	0.61	5/24324 (0.0%)

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	236	PRO	N-CA	13.07	1.69	1.47
1	C	133	PRO	N-CA	12.93	1.69	1.47
1	C	199	ARG	CZ-NH2	8.11	1.43	1.33
2	B	211	ARG	CB-CG	-5.99	1.36	1.52
3	F	13	DT	O3'-P	-5.56	1.54	1.61

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	211	ARG	NE-CZ-NH2	-10.03	115.29	120.30
1	C	133	PRO	CA-N-CD	-7.88	100.46	111.50
1	C	199	ARG	NE-CZ-NH1	-7.79	116.40	120.30
2	B	236	PRO	CA-N-CD	-5.83	103.33	111.50
2	B	211	ARG	CB-CG-CD	-5.14	98.22	111.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4497	0	4551	73	0
1	C	4505	0	4559	59	0
2	B	3347	0	3379	57	0
2	D	3347	0	3379	31	0
3	E	718	0	397	3	0
3	F	777	0	432	6	0
4	A	31	0	12	0	0
4	C	31	0	12	0	0
5	A	1	0	0	0	0
5	C	1	0	0	0	0
6	A	12	0	16	3	0
6	B	6	0	8	0	0
6	C	6	0	8	0	0
6	D	12	0	16	1	0
7	A	126	0	0	1	0
7	B	52	0	0	0	0
7	C	115	0	0	1	0
7	D	86	0	0	1	0
7	E	40	0	0	0	0
7	F	34	0	0	0	0
All	All	17744	0	16769	217	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 217 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:236:PRO:N	2:B:236:PRO:CA	1.69	1.45
1:C:133:PRO:N	1:C:133:PRO:CA	1.69	1.44
1:C:450:THR:HG23	1:C:452:LEU:H	1.37	0.88
2:D:356:ARG:HE	2:D:361:HIS:HB3	1.41	0.83
2:D:356:ARG:NE	2:D:361:HIS:HB3	1.95	0.81

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	551/557 (99%)	535 (97%)	15 (3%)	1 (0%)	47	58
1	C	552/557 (99%)	535 (97%)	17 (3%)	0	100	100
2	B	402/444 (90%)	385 (96%)	17 (4%)	0	100	100
2	D	402/444 (90%)	390 (97%)	12 (3%)	0	100	100
All	All	1907/2002 (95%)	1845 (97%)	61 (3%)	1 (0%)	51	63

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	135	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	492/494 (100%)	487 (99%)	5 (1%)	76	87
1	C	493/494 (100%)	486 (99%)	7 (1%)	67	80
2	B	365/400 (91%)	350 (96%)	15 (4%)	30	43
2	D	365/400 (91%)	353 (97%)	12 (3%)	38	52
All	All	1715/1788 (96%)	1676 (98%)	39 (2%)	50	66

5 of 39 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	D	72	ARG
2	D	356	ARG
2	D	89	GLU
2	D	311	LYS
2	D	424	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 25 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	136	ASN
1	C	278	GLN
2	D	407	GLN
1	C	221	HIS
1	C	315	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	OMC	E	4	3	19,22,23	2.71	8 (42%)	26,31,34	0.73	0
3	OMC	F	2	3	19,22,23	2.76	8 (42%)	26,31,34	0.76	0
3	OMC	F	4	3	19,22,23	2.67	8 (42%)	26,31,34	0.65	0
3	OMC	E	2	3	19,22,23	2.66	8 (42%)	26,31,34	0.71	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	OMC	E	4	3	-	0/9/27/28	0/2/2/2
3	OMC	F	2	3	-	0/9/27/28	0/2/2/2
3	OMC	F	4	3	-	0/9/27/28	0/2/2/2
3	OMC	E	2	3	-	0/9/27/28	0/2/2/2

The worst 5 of 32 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	F	2	OMC	C6-C5	6.01	1.49	1.35
3	F	4	OMC	C6-C5	5.88	1.48	1.35
3	E	4	OMC	C6-C5	5.83	1.48	1.35
3	E	4	OMC	C2-N3	5.68	1.47	1.36
3	F	2	OMC	C2-N3	5.63	1.47	1.36

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	E	4	OMC	1	0
3	F	2	OMC	1	0
3	F	4	OMC	2	0
3	E	2	OMC	1	0

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 2 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	DGT	C	601	5	26,33,33	3.67	13 (50%)	32,52,52	1.36	6 (18%)
6	GOL	A	603	-	5,5,5	0.84	0	5,5,5	1.09	1 (20%)
6	GOL	C	603	-	5,5,5	1.09	0	5,5,5	0.85	0
6	GOL	A	604	-	5,5,5	0.93	0	5,5,5	0.77	0
6	GOL	D	502	-	5,5,5	0.22	0	5,5,5	0.38	0
6	GOL	B	501	-	5,5,5	1.24	0	5,5,5	0.77	0
6	GOL	D	501	-	5,5,5	1.15	0	5,5,5	1.19	1 (20%)
4	DGT	A	601	5	26,33,33	3.63	13 (50%)	32,52,52	1.47	8 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	DGT	C	601	5	-	4/18/34/34	0/3/3/3
6	GOL	A	603	-	-	4/4/4/4	-
6	GOL	C	603	-	-	0/4/4/4	-
6	GOL	A	604	-	-	0/4/4/4	-
6	GOL	D	502	-	-	2/4/4/4	-
6	GOL	B	501	-	-	2/4/4/4	-
6	GOL	D	501	-	-	4/4/4/4	-
4	DGT	A	601	5	-	5/18/34/34	0/3/3/3

The worst 5 of 26 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	601	DGT	O4'-C1'	8.92	1.62	1.42
4	A	601	DGT	O4'-C1'	8.43	1.61	1.42
4	A	601	DGT	O4'-C4'	-7.43	1.28	1.45
4	C	601	DGT	O4'-C4'	-7.33	1.28	1.45
4	A	601	DGT	C2'-C1'	-6.65	1.33	1.52

The worst 5 of 16 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	601	DGT	C5-C6-N1	3.66	120.42	113.95

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	601	DGT	C2-N1-C6	-3.19	119.22	125.10
4	C	601	DGT	C5-C6-N1	3.12	119.47	113.95
4	C	601	DGT	C2-N1-C6	-2.88	119.79	125.10
4	A	601	DGT	C8-N7-C5	2.82	108.36	102.99

There are no chirality outliers.

5 of 21 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	601	DGT	PB-O3B-PG-O2G
4	C	601	DGT	PB-O3B-PG-O1G
6	A	603	GOL	O1-C1-C2-C3
6	D	501	GOL	C1-C2-C3-O3
6	A	603	GOL	C1-C2-C3-O3

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	604	GOL	3	0
6	D	502	GOL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	553/557 (99%)	0.47	41 (7%) 14 19	33, 59, 104, 180	0
1	C	553/557 (99%)	0.59	60 (10%) 5 8	33, 58, 109, 156	0
2	B	406/444 (91%)	1.03	67 (16%) 1 2	35, 69, 141, 176	0
2	D	406/444 (91%)	0.45	26 (6%) 19 25	32, 55, 100, 164	0
3	E	33/38 (86%)	-0.15	0 100 100	35, 55, 88, 138	0
3	F	36/38 (94%)	-0.00	1 (2%) 53 60	39, 65, 112, 157	0
All	All	1987/2078 (95%)	0.59	195 (9%) 7 10	32, 59, 119, 180	0

The worst 5 of 195 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	93	GLY	11.2
2	B	301	LEU	11.2
2	B	88	TRP	10.2
2	B	90	VAL	9.6
2	B	92	LEU	8.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	OMC	E	2	21/22	0.97	0.18	33,40,44,47	0
3	OMC	F	2	21/22	0.97	0.16	45,51,53,57	0
3	OMC	E	4	21/22	0.98	0.19	32,35,41,44	0
3	OMC	F	4	21/22	0.98	0.19	33,38,44,46	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	GOL	A	604	6/6	0.76	0.27	57,64,70,70	0
5	MG	C	602	1/1	0.85	0.21	55,55,55,55	0
5	MG	A	602	1/1	0.86	0.09	51,51,51,51	0
6	GOL	A	603	6/6	0.91	0.31	65,71,72,72	0
6	GOL	D	501	6/6	0.91	0.20	46,50,52,53	0
4	DGT	C	601	31/31	0.93	0.13	45,59,80,84	0
6	GOL	D	502	6/6	0.93	0.18	60,64,64,65	0
6	GOL	C	603	6/6	0.94	0.21	42,45,50,53	0
4	DGT	A	601	31/31	0.95	0.14	42,50,64,67	0
6	GOL	B	501	6/6	0.96	0.24	44,46,47,50	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around DGT C 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around DGT A 601:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.