



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 5, 2024 – 02:07 AM JST

PDB ID : 8X24
Title : anti-CRISPR protein-AcrIF25
Authors : Wang, J.J.
Deposited on : 2023-11-09
Resolution : 2.83 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.83 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1367 (2.86-2.82)
Clashscore	180529	1455 (2.86-2.82)
Ramachandran outliers	177936	1422 (2.86-2.82)
Sidechain outliers	177891	1423 (2.86-2.82)
RSRZ outliers	164620	1368 (2.86-2.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	153	 31% 16% . 50%
1	B	153	 37% 11% . 50%
1	C	153	 35% 14% . 50%
1	D	153	 35% 14% . 50%
1	E	153	 5% 33% 14% . 50%
1	F	153	 5% 31% 16% . 50%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3836 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called AcrIF25.



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	76	627	400	111	115	1	0	0	0
1	B	76	627	400	111	115	1	0	0	0
1	C	76	627	400	111	115	1	0	0	0
1	D	76	627	400	111	115	1	0	0	0
1	E	76	627	400	111	115	1	0	0	0
1	F	76	627	400	111	115	1	0	0	0

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	8	Total 8	O 8	0	0
2	B	19	Total 19	O 19	0	0
2	C	8	Total 8	O 8	0	0
2	D	12	Total 12	O 12	0	0
2	E	12	Total 12	O 12	0	0
2	F	15	Total 15	O 15	0	0

ALA THR ALA LYS TYR ASP GLU SER SER LYS SER GLY ASP VAL ILE THR TYR ARG SER SER GLU ASP GLY VAL LEU LEU GLU GLY ILE ILE ILE SER SER PHE ALA LYS LYS TRP GLU SER GLN ARG TYR SER ASN LEU PRO VAL VAL VAL GLU ILE VAL LYS LEU LEU HIS HIS HIS HIS HIS HIS HIS



• Molecule 1: AcrIF25



LYS ALA THR ALA LYS TYR ASP GLU SER SER LYS SER GLY ASP VAL ILE THR TYR ARG SER SER GLU ASP GLY VAL LEU LEU GLU GLY ILE ILE ILE SER SER PHE ALA LYS LYS TRP GLU SER GLN ARG TYR SER ASN LEU PRO VAL VAL VAL GLU ILE VAL LYS LEU LEU HIS HIS HIS HIS HIS HIS HIS

HIS

• Molecule 1: AcrIF25



THR LYS ASP ALA LYS TYR ASP GLU SER SER LYS SER GLY ASP VAL ILE THR TYR ARG SER SER GLU ASP GLY VAL LEU LEU GLU GLY ILE ILE ILE SER SER PHE ALA LYS LYS TRP GLU SER SER GLN ARG TYR SER ASN LEU PRO VAL VAL VAL GLU ILE VAL LYS LEU LEU HIS HIS HIS HIS HIS HIS HIS

HIS HIS HIS HIS HIS

4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	45.34Å 67.70Å 89.95Å 90.00° 90.44° 90.00°	Depositor
Resolution (Å)	44.97 – 2.83 44.97 – 2.83	Depositor EDS
% Data completeness (in resolution range)	97.0 (44.97-2.83) 97.0 (44.97-2.83)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.35 (at 2.82Å)	Xtrriage
Refinement program	PHENIX 1.19.2	Depositor
R, R_{free}	0.192 , (Not available) 0.192 , 0.241	Depositor DCC
R_{free} test set	1297 reflections (9.87%)	wwPDB-VP
Wilson B-factor (Å ²)	40.0	Xtrriage
Anisotropy	0.106	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 37.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.043 for h,-k,-l	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	3836	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.79% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.61	0/636	0.70	0/856
1	B	0.58	0/636	0.68	0/856
1	C	0.51	0/636	0.66	0/856
1	D	0.56	0/636	0.68	0/856
1	E	0.60	0/636	0.86	0/856
1	F	0.61	0/636	0.77	1/856 (0.1%)
All	All	0.58	0/3816	0.73	1/5136 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	1
1	E	0	2
All	All	0	3

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	F	72	ASN	CB-CA-C	6.19	122.78	110.40

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	34	ARG	Sidechain
1	E	43	ARG	Sidechain
1	E	62	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	627	0	641	22	0
1	B	627	0	641	16	0
1	C	627	0	641	12	0
1	D	627	0	641	14	0
1	E	627	0	641	14	0
1	F	627	0	641	18	0
2	A	8	0	0	2	0
2	B	19	0	0	0	0
2	C	8	0	0	0	0
2	D	12	0	0	1	0
2	E	12	0	0	2	0
2	F	15	0	0	0	0
All	All	3836	0	3846	85	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (85) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:18:ASN:HD22	1:E:22:LYS:HE2	1.35	0.89
1:C:12:GLY:N	1:D:62:ARG:HH12	1.73	0.87
1:E:18:ASN:ND2	1:E:22:LYS:HE2	1.90	0.85
1:B:27:LYS:H	1:B:27:LYS:CD	1.98	0.74
1:E:47:LEU:HA	2:E:201:HOH:O	1.88	0.72
1:F:1:MET:HG2	1:F:6:PHE:HE2	1.54	0.72
1:A:23:ILE:HD12	1:A:58:GLU:HG3	1.73	0.70
1:E:16:SER:N	1:E:19:GLN:OE1	2.28	0.66
1:A:2:LYS:HE3	1:A:44:ASP:OD2	1.97	0.65
1:A:18:ASN:HB2	1:A:34:ARG:HE	1.62	0.64
1:A:39:ASN:HB2	2:D:202:HOH:O	1.98	0.64
1:C:2:LYS:NZ	1:C:46:ASP:OD1	2.29	0.64
1:E:9:ALA:HB2	1:F:65:ILE:HG22	1.80	0.62
1:A:49:ASP:O	1:A:53:VAL:HG23	2.00	0.62
1:A:9:ALA:HB1	1:B:66:LEU:HD12	1.83	0.60

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:39:ASN:OD1	1:F:43:ARG:NH1	2.36	0.58
1:F:62:ARG:NH1	1:F:62:ARG:HG2	2.18	0.58
1:D:4:HIS:NE2	1:D:8:LEU:HD11	2.19	0.58
1:D:39:ASN:HB3	1:D:45:GLN:OE1	2.04	0.57
1:A:48:ASN:HD21	1:A:50:TRP:HB3	1.68	0.57
1:F:18:ASN:O	1:F:22:LYS:HG3	2.05	0.57
1:F:11:PHE:HB2	1:F:15:TYR:HB3	1.87	0.56
1:A:28:ASP:OD1	1:A:28:ASP:N	2.25	0.56
1:A:18:ASN:OD1	1:A:34:ARG:HG3	2.05	0.56
1:A:33:ASP:OD1	1:A:35:LYS:N	2.38	0.54
1:B:62:ARG:HH11	1:B:62:ARG:HG2	1.73	0.54
1:B:27:LYS:CD	1:B:27:LYS:N	2.71	0.53
1:F:67:LYS:O	1:F:71:THR:OG1	2.26	0.53
1:C:34:ARG:HD3	1:C:38:GLN:NE2	2.24	0.53
1:C:50:TRP:O	1:C:54:GLN:HG3	2.09	0.52
1:F:62:ARG:NH1	1:F:62:ARG:CG	2.73	0.52
1:B:27:LYS:H	1:B:27:LYS:HD2	1.73	0.52
1:B:28:ASP:OD1	1:B:30:SER:OG	2.27	0.51
1:D:16:SER:N	1:D:19:GLN:OE1	2.37	0.50
1:B:27:LYS:H	1:B:27:LYS:CE	2.25	0.50
1:B:62:ARG:HG2	1:B:62:ARG:NH1	2.24	0.50
1:A:37:ILE:O	1:A:41:ILE:HG13	2.12	0.49
1:E:40:TRP:HD1	1:E:45:GLN:OE1	1.94	0.49
1:A:43:ARG:NH1	2:A:203:HOH:O	2.45	0.49
1:A:48:ASN:O	1:A:52:ILE:HG13	2.13	0.48
1:B:27:LYS:N	1:B:27:LYS:HD2	2.29	0.48
1:C:1:MET:HG2	1:C:52:ILE:HD11	1.95	0.48
1:A:9:ALA:HB1	1:B:66:LEU:CD1	2.43	0.47
1:E:31:ASN:N	2:E:203:HOH:O	2.45	0.47
1:A:46:ASP:OD1	2:A:201:HOH:O	2.20	0.47
1:F:62:ARG:CG	1:F:62:ARG:HH11	2.27	0.47
1:D:48:ASN:HB3	1:D:51:VAL:HG23	1.96	0.47
1:C:5:GLU:HB3	1:D:69:LEU:HD13	1.98	0.46
1:C:74:SER:OG	1:C:75:GLN:HG2	2.16	0.46
1:C:56:LYS:HE2	1:D:73:LEU:HB3	1.96	0.46
1:E:18:ASN:O	1:E:22:LYS:HD2	2.16	0.46
1:A:48:ASN:HD22	1:A:51:VAL:HG23	1.82	0.45
1:B:62:ARG:HH11	1:B:62:ARG:CG	2.30	0.45
1:C:24:LEU:HD23	1:C:24:LEU:HA	1.85	0.44
1:A:48:ASN:ND2	1:A:51:VAL:HG23	2.32	0.44
1:E:69:LEU:HD21	1:F:5:GLU:HB3	2.00	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:70:LEU:HD23	1:F:70:LEU:HA	1.78	0.44
1:A:22:LYS:HE3	1:B:14:HIS:ND1	2.32	0.44
1:D:25:ILE:HG22	1:D:26:ASN:O	2.18	0.44
1:E:19:GLN:HG2	1:F:19:GLN:NE2	2.33	0.43
1:D:24:LEU:C	1:D:25:ILE:HD13	2.38	0.43
1:F:56:LYS:HA	1:F:59:LEU:HD12	1.99	0.43
1:C:76:ALA:HB1	1:F:56:LYS:HD2	2.01	0.43
1:A:25:ILE:HD12	1:A:26:ASN:O	2.19	0.43
1:A:60:LEU:HA	1:A:60:LEU:HD23	1.81	0.42
1:B:11:PHE:HB2	1:B:15:TYR:HB3	2.00	0.42
1:F:13:GLU:H	1:F:13:GLU:HG3	1.70	0.42
1:B:24:LEU:HD12	1:B:24:LEU:HA	1.87	0.42
1:E:19:GLN:NE2	1:F:19:GLN:HE21	2.18	0.42
1:A:56:LYS:HE2	1:A:56:LYS:HB3	1.86	0.42
1:C:25:ILE:HD13	1:C:25:ILE:HA	1.95	0.41
1:D:61:LYS:O	1:D:64:VAL:HG22	2.20	0.41
1:F:1:MET:HE3	1:F:1:MET:HB2	1.84	0.41
1:A:48:ASN:OD1	1:A:49:ASP:N	2.53	0.41
1:B:1:MET:O	1:B:46:ASP:HA	2.21	0.41
1:F:54:GLN:O	1:F:57:GLU:HB2	2.20	0.41
1:E:35:LYS:HD2	1:E:35:LYS:HA	1.67	0.41
1:D:55:LEU:HA	1:D:55:LEU:HD23	1.74	0.41
1:D:24:LEU:O	1:D:25:ILE:HD13	2.20	0.41
1:E:50:TRP:O	1:E:54:GLN:HG3	2.20	0.41
1:E:11:PHE:HB2	1:E:15:TYR:HB3	2.02	0.41
1:D:10:LEU:HA	1:D:10:LEU:HD23	1.82	0.41
1:B:35:LYS:HG3	1:B:36:THR:N	2.36	0.40
1:C:66:LEU:HG	1:C:70:LEU:HD12	2.03	0.40
1:D:70:LEU:HD23	1:D:70:LEU:HA	1.89	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	74/153 (48%)	72 (97%)	2 (3%)	0	100	100
1	B	74/153 (48%)	71 (96%)	3 (4%)	0	100	100
1	C	74/153 (48%)	71 (96%)	3 (4%)	0	100	100
1	D	74/153 (48%)	71 (96%)	3 (4%)	0	100	100
1	E	74/153 (48%)	73 (99%)	1 (1%)	0	100	100
1	F	74/153 (48%)	72 (97%)	2 (3%)	0	100	100
All	All	444/918 (48%)	430 (97%)	14 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	69/139 (50%)	61 (88%)	8 (12%)	4	9
1	B	69/139 (50%)	59 (86%)	10 (14%)	2	5
1	C	69/139 (50%)	64 (93%)	5 (7%)	12	25
1	D	69/139 (50%)	65 (94%)	4 (6%)	17	34
1	E	69/139 (50%)	58 (84%)	11 (16%)	2	3
1	F	69/139 (50%)	58 (84%)	11 (16%)	2	3
All	All	414/834 (50%)	365 (88%)	49 (12%)	4	8

All (49) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	13	GLU
1	A	28	ASP
1	A	30	SER
1	A	34	ARG
1	A	43	ARG
1	A	48	ASN
1	A	49	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	62	ARG
1	B	3	ILE
1	B	27	LYS
1	B	30	SER
1	B	34	ARG
1	B	43	ARG
1	B	49	ASP
1	B	62	ARG
1	B	67	LYS
1	B	70	LEU
1	B	75	GLN
1	C	16	SER
1	C	30	SER
1	C	34	ARG
1	C	49	ASP
1	C	58	GLU
1	D	5	GLU
1	D	33	ASP
1	D	43	ARG
1	D	62	ARG
1	E	1	MET
1	E	22	LYS
1	E	28	ASP
1	E	33	ASP
1	E	35	LYS
1	E	43	ARG
1	E	48	ASN
1	E	49	ASP
1	E	66	LEU
1	E	74	SER
1	E	75	GLN
1	F	1	MET
1	F	2	LYS
1	F	8	LEU
1	F	13	GLU
1	F	26	ASN
1	F	34	ARG
1	F	39	ASN
1	F	44	ASP
1	F	58	GLU
1	F	62	ARG
1	F	75	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	19	GLN
1	C	45	GLN
1	C	72	ASN
1	D	45	GLN
1	D	54	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	76/153 (49%)	-0.50	0 100 100	27, 34, 51, 58	0
1	B	76/153 (49%)	-0.49	0 100 100	24, 32, 49, 75	0
1	C	76/153 (49%)	-0.49	0 100 100	25, 35, 53, 60	0
1	D	76/153 (49%)	-0.43	0 100 100	26, 35, 55, 64	0
1	E	76/153 (49%)	-0.07	1 (1%) 74 71	31, 43, 70, 95	0
1	F	76/153 (49%)	-0.28	1 (1%) 74 71	30, 46, 65, 88	0
All	All	456/918 (49%)	-0.38	2 (0%) 89 88	24, 37, 64, 95	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	76	ALA	3.0
1	F	35	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.