



Full wwPDB EM Validation Report ⓘ

Feb 25, 2024 – 10:48 AM EST

PDB ID : 6X68
EMDB ID : EMD-22073
Title : Cryo-EM structure of piggyBac transposase synaptic complex with hairpin DNA (SNHP)
Authors : Chen, Q.; Hickman, A.B.; Dyda, F.
Deposited on : 2020-05-27
Resolution : 3.66 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev70
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.66 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	C	594	
1	D	594	
2	A	74	
2	B	74	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	ZN	C	603	-	-	X	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	ZN	C	604	-	-	X	-
4	ZN	D	602	-	-	X	-
4	ZN	D	603	-	-	X	-

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 10246 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transposase.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	C	478	3848	2424	676	704	44	0	0
1	D	478	3848	2424	676	704	44	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	500	LYS	GLU	variant	UNP Q283G1
D	500	LYS	GLU	variant	UNP Q283G1

- Molecule 2 is a DNA chain called hairpin DNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	P		
2	A	74	1517	725	274	444	74	0	0
2	B	50	1025	490	185	300	50	0	0

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	AltConf
3	C	2	Total Ca 2 2	0
3	D	1	Total Ca 1 1	0
3	A	1	Total Ca 1 1	0

- Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		AltConf
4	C	2	Total 2	Zn 2	0
4	D	2	Total 2	Zn 2	0

3 Residue-property plots i

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Transposase

Chain C: 

MET GLY SER SER LEU ASP ASP GLU HIS ILE ILE LEU LEU LEU ALA LEU LEU LEU GLN SER ASP ASP GLU LEU VAL VAL GLY ASP ASP SER ASP ASP GLN ARG THR THR ILE ILE SER ASP ASP HIS VAL VAL LYS LYS LYS CYS ASP ASP ASP VAL GLN SER SER ASP ASP THR THR LYS LYS THR THR ARG ARG ALA ASP ILE ASP ILE ASP VAL VAL VAL HIS HIS VAL VAL VAL VAL PRO THR THR SER SER GLY

SER GLU ILE LEU ASP GLU ASN VAL ILE ILE GLN PRO GLY SER SER LEU ALA SER ASN LYS ILE ILE THR LEU LEU PRO GLN ARG THR ILE ILE ARG ARG LYS LYS ASN HIS CYS TRP SER THR THR LYS LYS THR THR ARG ARG SER SER ARG VAL VAL SER ALA LEU ASN ILE VAL ARG ARG S117 Y129

G181 D191 M194 W212 S213 R216 D227 T231 R233 R236 D239 V240 F241 T242 P243 V244 D268 R275 G276 R277 C278 P288 S289 P311 T319 V322 R372 R376 E377 E380 V381 L382 K383 N384 S385 S405 P408 K409 P410

Y415 D421 E422 D423 A424 M435 R464 W465 P466 K492 L510 T511 S512 R516 R526 P541 G542 D545 D546 S547 T548 E549 E550 K555 C559 T560 Y561 C574 C577 C582 R583 E584 H585 N586 I587 D588 H589 C593 F594

- Molecule 1: Transposase

Chain D: 


MET GLY SER SER LEU ASP ASP GLU HIS ILE ILE LEU LEU LEU ALA LEU LEU LEU GLN SER ASP ASP GLU LEU VAL VAL GLY LEU THR LEU ASP SER ASP ASP THR THR ILE ILE SER ASP ASP HIS VAL VAL LYS LYS LYS CYS ASP ASP ASP VAL GLN SER SER ASP ASP THR THR LYS LYS THR THR ARG ARG ALA ASP ILE ASP ILE ASP VAL VAL VAL HIS HIS VAL VAL VAL VAL VAL PRO THR THR SER SER GLY

SER GLU ILE LEU ASP GLU ASN VAL ILE ILE GLN PRO GLY SER SER LEU ALA SER ASN LYS ILE ILE THR LEU LEU PRO GLN ARG THR ILE ILE ARG ARG LYS LYS ASN HIS CYS TRP SER THR THR LYS LYS THR THR ARG ARG SER SER ARG VAL VAL SER ALA LEU ASN ILE VAL ARG ARG S117 P121

C125 P131 L137 F179 D197 R216 F217 D218 L235 E269 Q270 L271 F280 K294 D300 T303 K304 R315 K334 P335 G338 C345 F349 F361 P362 Y363 K364 R376 R383 H384 P389 T392 D396 P408 D421 F422

D423 W465 P466 I484 Y485 K492 G493 E494 F502 Y507 Y527 V540 P541 G542 T543 S544 D545 D546 S547 T548 E549 E550 P551 V552 T553 K554 K555 R556 T557 Y558 C559 T560 Y561 C562 N571 A572 S573 C574 K575 K576 C577 I581 C582 H585 N586 I587 C590 C593

- Molecule 2: hairpin DNA

Chain A:  80% 16%



• Molecule 2: hairpin DNA

Chain B:  59% 8% 32%



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	35960	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	73.7	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.078	Depositor
Minimum map value	-0.058	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.003	Depositor
Recommended contour level	0.008	Depositor
Map size (Å)	205.20001, 205.20001, 205.20001	wwPDB
Map dimensions	190, 190, 190	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.08, 1.08, 1.08	Depositor

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: CA, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	C	0.65	0/3926	0.91	6/5286 (0.1%)
1	D	0.66	0/3926	0.89	6/5286 (0.1%)
2	A	1.03	3/1701 (0.2%)	1.52	26/2623 (1.0%)
2	B	0.90	0/1149	1.38	6/1771 (0.3%)
All	All	0.76	3/10702 (0.0%)	1.10	44/14966 (0.3%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	34	DT	C4'-C3'	8.85	1.62	1.53
2	A	33	DA	O3'-P	-7.62	1.52	1.61
2	A	33	DA	C3'-O3'	-5.29	1.37	1.44

All (44) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	33	DA	O4'-C4'-C3'	-11.63	99.02	106.00
1	C	384	ASN	CA-C-N	-10.82	93.39	117.20
2	B	-3	DT	P-O3'-C3'	10.13	131.85	119.70
2	A	-3	DT	P-O3'-C3'	9.78	131.44	119.70
2	A	34	DT	C3'-C2'-C1'	-9.74	90.81	102.50
2	A	34	DT	O4'-C1'-N1	8.76	114.13	108.00
2	A	34	DT	O4'-C4'-C3'	-8.23	101.06	106.00
2	B	-18	DA	O4'-C1'-N9	8.22	113.75	108.00
1	D	315	ARG	CA-C-N	-7.61	100.98	116.20
2	A	35	DG	OP1-P-OP2	-7.10	108.96	119.60
2	A	16	DC	N3-C2-O2	-6.91	117.06	121.90
2	A	34	DT	P-O3'-C3'	6.91	127.99	119.70
1	C	384	ASN	O-C-N	6.89	133.73	122.70
2	A	34	DT	OP1-P-OP2	-6.65	109.62	119.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	384	ASN	C-N-CA	6.48	137.89	121.70
2	A	33	DA	C4'-C3'-C2'	6.46	108.91	103.10
2	A	32	DC	OP1-P-OP2	-6.16	110.36	119.60
2	A	31	DG	OP1-P-OP2	-6.14	110.39	119.60
1	C	384	ASN	CA-C-O	6.09	132.88	120.10
2	A	33	DA	OP1-P-OP2	-6.08	110.49	119.60
2	A	32	DC	C6-N1-C2	5.96	122.68	120.30
2	A	33	DA	P-O3'-C3'	-5.88	112.64	119.70
2	B	-26	DT	O4'-C4'-C3'	-5.83	102.17	104.50
2	B	-9	DC	O4'-C1'-C2'	-5.83	101.23	105.90
2	A	-23	DC	O4'-C4'-C3'	-5.78	102.19	104.50
1	D	527	TYR	CB-CA-C	-5.73	98.95	110.40
2	A	26	DT	O4'-C4'-C3'	-5.73	102.21	104.50
2	A	-17	DC	P-O3'-C3'	5.56	126.37	119.70
2	A	-5	DG	P-O3'-C3'	5.53	126.33	119.70
2	A	-3	DT	O4'-C4'-C3'	-5.52	102.29	104.50
2	A	26	DT	N3-C4-O4	5.49	123.19	119.90
1	C	384	ASN	N-CA-C	5.48	125.81	111.00
1	D	507	TYR	CB-CG-CD1	5.48	124.29	121.00
2	B	-1	DA	N1-C6-N6	5.45	121.87	118.60
1	C	415	TYR	CB-CG-CD2	-5.42	117.75	121.00
2	A	-5	DG	O4'-C1'-N9	5.42	111.79	108.00
2	A	-14	DT	P-O3'-C3'	5.37	126.14	119.70
2	A	-20	DA	O4'-C1'-N9	5.31	111.72	108.00
1	D	280	PHE	CB-CA-C	-5.30	99.79	110.40
1	D	485	TYR	CB-CG-CD1	-5.27	117.84	121.00
2	A	26	DT	C5-C4-O4	-5.09	121.34	124.90
2	A	10	DG	P-O3'-C3'	5.08	125.79	119.70
1	D	561	TYR	CB-CG-CD2	-5.06	117.96	121.00
2	B	10	DG	P-O3'-C3'	5.01	125.71	119.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	3848	0	3890	34	0
1	D	3848	0	3892	37	0
2	A	1517	0	837	6	0
2	B	1025	0	566	0	0
3	A	1	0	0	1	0
3	C	2	0	0	0	0
3	D	1	0	0	0	0
4	C	2	0	0	4	0
4	D	2	0	0	5	0
All	All	10246	0	9185	75	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (75) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:131:PRO:HB3	1:D:502:PHE:CE1	1.98	0.98
1:C:577:CYS:HG	4:C:604:ZN:ZN	0.64	0.97
1:D:590:CYS:SG	1:D:593:CYS:SG	2.55	0.94
1:D:562:CYS:HG	4:D:602:ZN:ZN	0.79	0.90
1:C:129:TYR:OH	1:C:492:LYS:HE3	1.70	0.89
1:D:574:CYS:SG	1:D:593:CYS:SG	2.71	0.88
1:D:557:THR:HG22	1:D:558:TYR:N	1.89	0.85
1:D:562:CYS:SG	4:D:602:ZN:ZN	1.71	0.80
1:C:577:CYS:SG	4:C:604:ZN:ZN	1.71	0.79
1:C:577:CYS:SG	1:C:593:CYS:SG	2.83	0.77
1:D:574:CYS:HG	4:D:603:ZN:ZN	0.95	0.76
1:D:574:CYS:SG	4:D:603:ZN:ZN	1.76	0.74
1:D:557:THR:HG22	1:D:558:TYR:H	1.53	0.72
2:A:-3:DT:OP1	3:A:101:CA:CA	1.65	0.72
1:D:593:CYS:SG	4:D:603:ZN:ZN	1.77	0.71
1:D:389:PRO:HG2	1:D:392:THR:HG21	1.72	0.71
1:C:129:TYR:OH	1:C:492:LYS:CE	2.39	0.70
2:A:33:DA:H2''	2:A:34:DT:H5'	1.76	0.68
1:D:559:CYS:SG	1:D:562:CYS:SG	2.91	0.67
1:C:582:CYS:HG	4:C:603:ZN:ZN	1.05	0.66
1:D:557:THR:CG2	1:D:558:TYR:N	2.58	0.63
1:D:557:THR:CG2	1:D:558:TYR:H	2.11	0.63
1:C:582:CYS:SG	4:C:603:ZN:ZN	1.87	0.62
1:C:408:PRO:HB2	1:C:410:PRO:HD2	1.81	0.61
1:C:512:SER:O	1:C:516:ARG:HG3	2.01	0.59

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:576:LYS:HB2	1:D:593:CYS:HB3	1.84	0.59
1:D:586:ASN:HB3	1:D:590:CYS:HA	1.86	0.56
2:A:33:DA:H2''	2:A:34:DT:C5'	2.36	0.54
1:C:241:PHE:CZ	1:C:311:PRO:HG3	2.42	0.54
1:D:559:CYS:HB2	1:D:582:CYS:HB3	1.90	0.54
1:D:271:LEU:CD2	1:D:294:LYS:HG3	2.37	0.54
1:C:380:GLU:O	1:C:384:ASN:HB2	2.07	0.54
1:C:243:PRO:HG2	1:C:465:TRP:HZ3	1.73	0.54
1:D:577:CYS:HB3	1:D:593:CYS:SG	2.50	0.52
1:D:562:CYS:SG	1:D:585:HIS:CE1	3.03	0.52
1:C:191:ASP:N	1:C:191:ASP:OD1	2.42	0.51
1:D:574:CYS:SG	1:D:590:CYS:SG	3.07	0.51
1:D:492:LYS:HG2	1:D:492:LYS:O	2.10	0.51
2:A:34:DT:H2'	2:A:34:DT:O2	2.09	0.51
1:C:561:TYR:HB2	1:C:585:HIS:CE1	2.47	0.50
1:D:334:LYS:HB3	1:D:335:PRO:HD3	1.94	0.49
1:C:574:CYS:SG	1:C:577:CYS:SG	3.10	0.49
1:D:121:PRO:HB3	1:D:484:ILE:HD12	1.95	0.49
1:C:319:THR:HG22	1:C:322:VAL:O	2.13	0.49
1:C:382:LEU:HB3	1:C:405:SER:HB2	1.95	0.49
1:C:464:ARG:HG2	1:C:466:PRO:HD2	1.93	0.49
1:D:304:LYS:N	1:D:304:LYS:HD3	2.28	0.48
1:C:288:PRO:O	1:C:289:SER:C	2.50	0.48
1:C:559:CYS:HB2	1:C:582:CYS:HB3	1.96	0.48
2:A:34:DT:O2	2:A:34:DT:C2'	2.60	0.48
1:D:338:GLY:H	1:D:364:LYS:H	1.59	0.47
1:D:494:GLU:HA	1:D:494:GLU:OE1	2.15	0.47
1:C:319:THR:CG2	1:C:322:VAL:O	2.63	0.46
1:C:231:ILE:C	1:C:233:PRO:HD2	2.36	0.45
1:C:421:ASP:OD1	1:C:421:ASP:N	2.47	0.45
1:D:300:ASP:O	1:D:303:THR:O	2.33	0.45
1:D:465:TRP:N	1:D:466:PRO:CD	2.79	0.45
1:C:212:MET:HB3	1:C:213:SER:H	1.47	0.44
1:C:227:ASP:N	1:C:227:ASP:OD1	2.50	0.44
1:D:345:CYS:HB3	1:D:349:PHE:HB2	2.00	0.43
1:C:241:PHE:CE2	1:C:244:VAL:HG23	2.54	0.43
1:D:269:GLU:OE1	1:D:294:LYS:NZ	2.41	0.42
1:C:278:CYS:O	1:C:278:CYS:SG	2.77	0.42
1:C:181:GLY:HA2	1:C:510:LEU:HD23	2.01	0.42
1:D:572:ALA:HB3	1:D:581:ILE:HG13	2.02	0.42
1:D:574:CYS:SG	1:D:577:CYS:SG	3.17	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:271:LEU:HD23	1:D:294:LYS:HG3	2.02	0.42
1:D:465:TRP:CG	1:D:466:PRO:HD3	2.56	0.41
1:C:191:ASP:HB2	1:C:194:MET:HG3	2.02	0.41
1:C:587:ILE:HB	1:D:587:ILE:CG2	2.51	0.41
1:C:239:ASP:O	1:C:242:THR:OG1	2.27	0.41
1:C:577:CYS:HB3	1:C:593:CYS:SG	2.61	0.41
1:C:242:THR:HB	1:C:243:PRO:HD3	2.02	0.41
1:D:125:CYS:SG	1:D:137:LEU:HD13	2.60	0.40
1:C:275:ARG:NH1	2:A:-6:DG:N7	2.69	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	476/594 (80%)	460 (97%)	15 (3%)	1 (0%)	47	78
1	D	476/594 (80%)	465 (98%)	9 (2%)	2 (0%)	34	69
All	All	952/1188 (80%)	925 (97%)	24 (2%)	3 (0%)	44	74

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	385	SER
1	D	363	TYR
1	D	408	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	442/549 (80%)	437 (99%)	5 (1%)	73	85
1	D	442/549 (80%)	438 (99%)	4 (1%)	78	88
All	All	884/1098 (80%)	875 (99%)	9 (1%)	77	86

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	216	ARG
1	C	385	SER
1	C	526	ARG
1	C	583	ARG
1	C	589	MET
1	D	179	PHE
1	D	216	ARG
1	D	280	PHE
1	D	315	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	193	HIS
1	C	487	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry

Of 8 ligands modelled in this entry, 8 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-22073. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map




The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 83



Y Index: 109




Z Index: 101


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y



Z


The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.008. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 87 nm^3 ; this corresponds to an approximate mass of 79 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [\(i\)](#)



*Reported resolution corresponds to spatial frequency of 0.273\AA^{-1}


8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-22073 and PDB model 6X68. Per-residue inclusion information can be found in section 3 on page 6.

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.008 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.008).


9.4 Atom inclusion [i](#)



At the recommended contour level, 90% of all backbone atoms, 81% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary [i](#)

The table lists the average atom inclusion at the recommended contour level (0.008) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.8100	 0.4170
A	 0.9220	 0.3870
B	 0.9590	 0.4120
C	 0.7720	 0.4250
D	 0.7610	 0.4210

