



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2018 – 11:32 pm GMT

PDB ID : 1X7Q  
Title : Crystal structure of HLA-A\*1101 with sars nucleocapsid peptide  
Authors : Blicher, T.; Kastrup, J.S.; Buus, S.; Gajhede, M.  
Deposited on : 2004-08-16  
Resolution : 1.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.3 (157068), CSD as539be (2018)  
Xtrriage (Phenix) : 1.13  
EDS : trunk30967  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk30967


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	1761 (1.48-1.44)
Clashscore	122126	1816 (1.48-1.44)
Ramachandran outliers	120053	1793 (1.48-1.44)
Sidechain outliers	120020	1793 (1.48-1.44)
RSRZ outliers	108989	1733 (1.48-1.44)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	275	 83% 15%
2	B	99	 79% 18%
3	C	9	 78% 11% 11%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	GOL	A	5006	-	-	X	-
5	GOL	B	5004	-	-	X	-

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 3855 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HLA class I histocompatibility antigen, A-11 alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	275	2310	1431	425	444	10	0	16	0


- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	99	837	534	140	160	3	0	3	0

- Molecule 3 is a protein called SARS NUCLEOCAPSID PEPTIDE.


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	C	9	77	52	11	14	0	1	0

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O S 5 4 1	0	0
4	A	1	Total O S 5 4 1	0	0
4	B	1	Total O S 5 4 1	0	0

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	B	1	Total C O 6 3 3	0	0
5	B	1	Total C O 6 3 3	0	0


- Molecule 6 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
6	A	398	Total 398	O 398	0	0
6	B	166	Total 166	O 166	0	0
6	C	10	Total 10	O 10	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: HLA class I histocompatibility antigen, A-11 alpha chain



- Molecule 2: Beta-2-microglobulin



- Molecule 3: SARS NUCLEOCAPSID PEPTIDE



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	58.47Å 80.62Å 56.57Å 90.00° 116.79° 90.00°	Depositor
Resolution (Å)	30.00 – 1.45 31.90 – 1.45	Depositor EDS
% Data completeness (in resolution range)	99.1 (30.00-1.45) 95.8 (31.90-1.45)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	0.10	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.50 (at 1.45Å)	Xtrriage
Refinement program	SHELXL-97	Depositor
R, $R_{free}$	0.133 , 0.181 0.141 , 0.163	Depositor DCC
$R_{free}$ test set	4099 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	12.7	Xtrriage
Anisotropy	0.224	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 56.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.023 for -h-1,-k,l	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	3855	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	22.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.56% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.54	0/2440	1.36	40/3305 (1.2%)
2	B	0.53	0/872	1.18	3/1181 (0.3%)
3	C	0.63	0/85	1.12	0/114
All	All	0.54	0/3397	1.31	43/4600 (0.9%)

There are no bond length outliers.

All (43) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	35[A]	ARG	NE-CZ-NH2	-9.73	115.43	120.30
1	A	35[B]	ARG	NE-CZ-NH2	-9.73	115.43	120.30
1	A	14	ARG	NE-CZ-NH2	-9.61	115.50	120.30
1	A	170	ARG	NE-CZ-NH1	9.24	124.92	120.30
1	A	44	ARG	NE-CZ-NH2	-8.10	116.25	120.30
1	A	108	ARG	NE-CZ-NH1	8.00	124.30	120.30
1	A	114	ARG	NE-CZ-NH1	7.65	124.12	120.30
1	A	35[A]	ARG	NE-CZ-NH1	7.49	124.05	120.30
1	A	35[B]	ARG	NE-CZ-NH1	7.49	124.05	120.30
1	A	108	ARG	NE-CZ-NH2	-7.27	116.66	120.30
1	A	17	ARG	NE-CZ-NH1	7.07	123.83	120.30
1	A	48	ARG	NE-CZ-NH1	6.99	123.80	120.30
1	A	163	ARG	NE-CZ-NH1	-6.91	116.84	120.30
1	A	209	TYR	CB-CG-CD1	6.90	125.14	121.00
2	B	58	LYS	CD-CE-NZ	6.82	127.39	111.70
1	A	169	ARG	NE-CZ-NH2	-6.61	117.00	120.30
2	B	10	TYR	CB-CG-CD2	6.37	124.82	121.00
1	A	75	ARG	NE-CZ-NH1	6.27	123.43	120.30
1	A	163	ARG	NE-CZ-NH2	6.20	123.40	120.30
1	A	170	ARG	NE-CZ-NH2	-6.12	117.24	120.30
1	A	6	ARG	NE-CZ-NH1	6.10	123.35	120.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	159	TYR	CB-CG-CD2	6.05	124.63	121.00
2	B	97	ARG	NE-CZ-NH1	6.03	123.31	120.30
1	A	46	GLU	OE1-CD-OE2	-6.01	116.08	123.30
1	A	174	ASN	CA-CB-CG	5.98	126.56	113.40
1	A	173	GLU	C-N-CA	5.91	136.46	121.70
1	A	82[A]	ARG	NE-CZ-NH1	5.81	123.20	120.30
1	A	82[B]	ARG	NE-CZ-NH1	5.81	123.20	120.30
1	A	84	TYR	CB-CG-CD1	5.77	124.46	121.00
1	A	157[A]	ARG	NE-CZ-NH1	5.75	123.17	120.30
1	A	157[B]	ARG	NE-CZ-NH1	5.75	123.17	120.30
1	A	244	TRP	CA-CB-CG	5.67	124.48	113.70
1	A	17	ARG	O-C-N	-5.46	113.92	123.20
1	A	111	ARG	NE-CZ-NH2	5.41	123.01	120.30
1	A	157[A]	ARG	CD-NE-CZ	5.39	131.15	123.60
1	A	157[B]	ARG	CD-NE-CZ	5.39	131.15	123.60
1	A	84	TYR	CB-CG-CD2	-5.37	117.78	121.00
1	A	256	ARG	NE-CZ-NH2	-5.34	117.63	120.30
1	A	59	TYR	CB-CG-CD1	5.20	124.12	121.00
1	A	17	ARG	CA-C-N	5.19	126.58	116.20
1	A	110	LEU	CA-CB-CG	5.18	127.22	115.30
1	A	219	ARG	NE-CZ-NH2	-5.18	117.71	120.30
1	A	264	GLU	O-C-N	-5.11	114.51	123.20

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2310	0	2161	32	0
2	B	837	0	803	16	0
3	C	77	0	83	1	0
4	A	10	0	0	0	0
4	B	5	0	0	0	0
5	A	30	0	40	9	0
5	B	12	0	16	6	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	A	398	0	0	15	0
6	B	166	0	0	6	0
6	C	10	0	0	0	0
All	All	3855	0	3103	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:8:GLN:HG2	5:B:5004:GOL:H32	1.57	0.86
1:A:154:GLU:HG3	5:A:5006:GOL:O3	1.84	0.78
1:A:151:HIS:HA	5:A:5006:GOL:H11	1.66	0.77
2:B:4[B]:THR:HG23	2:B:86:THR:OG1	1.92	0.69
5:A:5006:GOL:H2	6:A:5262:HOH:O	1.98	0.62
1:A:231:VAL:HG12	5:B:5004:GOL:H2	1.83	0.61
5:A:5006:GOL:H12	6:A:5172:HOH:O	2.00	0.61
1:A:8:PHE:CE2	1:A:98[B]:MET:HG3	2.36	0.60
5:B:5004:GOL:H31	6:B:5014:HOH:O	2.00	0.60
1:A:231:VAL:CG1	5:B:5004:GOL:H2	2.33	0.58
2:B:8:GLN:CG	5:B:5004:GOL:H32	2.32	0.57
1:A:151:HIS:HA	5:A:5006:GOL:C1	2.35	0.56
1:A:169:ARG:HG2	6:A:5203:HOH:O	2.05	0.55
2:B:74:GLU:HA	2:B:74:GLU:OE2	2.05	0.55
1:A:174:ASN:HA	6:A:5272:HOH:O	2.07	0.55
1:A:173:GLU:O	1:A:173:GLU:HG3	2.07	0.54
1:A:275:GLU:HG2	6:A:5287:HOH:O	2.07	0.54
2:B:4[B]:THR:HG22	2:B:5:PRO:HD2	1.92	0.52
1:A:114:ARG:HD3	1:A:156[A]:GLN:NE2	2.25	0.51
1:A:117:ALA:HB2	2:B:60:TRP:CE2	2.47	0.49
1:A:230:LEU:HG	1:A:243[B]:LYS:NZ	2.27	0.48
5:A:5003:GOL:H12	6:A:5140:HOH:O	2.13	0.48
2:B:76:ASP:O	2:B:77:GLU:OE2	2.32	0.48
2:B:1:ILE:HD12	6:B:5123:HOH:O	2.14	0.47
2:B:73:THR:OG1	2:B:75:LYS:HD2	2.15	0.47
1:A:156[A]:GLN:NE2	6:A:5129:HOH:O	2.48	0.46
2:B:6:LYS:HE3	2:B:28:SER:OG	2.16	0.46
1:A:114:ARG:HD3	1:A:156[A]:GLN:HE21	1.80	0.45
1:A:19:GLU:CD	5:A:5001:GOL:H2	2.37	0.45
2:B:58:LYS:HG2	2:B:59:ASP:N	2.30	0.45

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:3:ARG:NH2	6:B:5126:HOH:O	2.49	0.45
1:A:35[B]:ARG:NH1	6:A:5338:HOH:O	2.49	0.45
2:B:3:ARG:NH1	6:B:5047:HOH:O	2.49	0.45
1:A:19:GLU:HG3	5:A:5001:GOL:H32	1.98	0.44
2:B:20[A]:SER:OG	5:B:5007:GOL:H32	2.17	0.44
1:A:131:ARG:NH2	6:A:5369:HOH:O	2.50	0.44
2:B:1:ILE:HG12	6:B:5126:HOH:O	2.17	0.44
3:C:6:THR:HG23	3:C:7:GLU:N	2.33	0.44
1:A:82[A]:ARG:NH1	6:A:5367:HOH:O	2.51	0.43
1:A:62:GLN:HG3	6:A:5357:HOH:O	2.19	0.43
1:A:154:GLU:OE1	1:A:157[A]:ARG:NH2	2.51	0.43
1:A:14:ARG:HD3	6:A:5273:HOH:O	2.19	0.42
1:A:131:ARG:NH2	1:A:157[B]:ARG:HH12	2.17	0.42
2:B:3:ARG:NE	6:B:5126:HOH:O	2.49	0.42
5:A:5002:GOL:H12	6:A:5336:HOH:O	2.19	0.42
1:A:54:GLN:HB3	6:A:5292:HOH:O	2.19	0.41
1:A:145[B]:ARG:NE	6:A:5165:HOH:O	2.50	0.41
1:A:51:TRP:CZ2	1:A:179:LEU:HD11	2.56	0.41
1:A:98[B]:MET:HE2	1:A:98[B]:MET:HB3	1.89	0.40
1:A:275:GLU:HG3	1:A:275:GLU:OXT	2.22	0.40
1:A:131:ARG:NH2	1:A:157[B]:ARG:NH1	2.69	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	289/275 (105%)	288 (100%)	1 (0%)	0	100	100
2	B	100/99 (101%)	99 (99%)	1 (1%)	0	100	100
3	C	8/9 (89%)	8 (100%)	0	0	100	100
All	All	397/383 (104%)	395 (100%)	2 (0%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	248/232 (107%)	243 (98%)	5 (2%)	58	22
2	B	97/94 (103%)	92 (95%)	5 (5%)	25	3
3	C	10/9 (111%)	9 (90%)	1 (10%)	8	0
All	All	355/335 (106%)	344 (97%)	11 (3%)	44	10

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	58	GLU
1	A	115	GLN
1	A	196	ASP
1	A	251[A]	SER
1	A	251[B]	SER
2	B	48	LYS
2	B	58	LYS
2	B	70	PHE
2	B	75	LYS
2	B	77	GLU
3	C	7	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	174	ASN
1	A	192	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	SO4	A	4002	-	4,4,4	0.64	0	6,6,6	0.39	0
4	SO4	A	4003	-	4,4,4	0.39	0	6,6,6	0.38	0
5	GOL	A	5001	-	5,5,5	0.34	0	5,5,5	0.84	0
5	GOL	A	5002	-	5,5,5	0.47	0	5,5,5	0.42	0
5	GOL	A	5003	-	5,5,5	0.28	0	5,5,5	0.54	0
5	GOL	A	5005	-	5,5,5	0.45	0	5,5,5	0.52	0
5	GOL	A	5006	-	5,5,5	0.49	0	5,5,5	1.03	0
4	SO4	B	4001	-	4,4,4	0.48	0	6,6,6	0.37	0
5	GOL	B	5004	-	5,5,5	0.26	0	5,5,5	0.58	0
5	GOL	B	5007	-	5,5,5	0.37	0	5,5,5	1.16	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	SO4	A	4002	-	-	0/0/0/0	0/0/0/0
4	SO4	A	4003	-	-	0/0/0/0	0/0/0/0
5	GOL	A	5001	-	-	0/4/4/4	0/0/0/0
5	GOL	A	5002	-	-	0/4/4/4	0/0/0/0
5	GOL	A	5003	-	-	0/4/4/4	0/0/0/0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	A	5005	-	-	0/4/4/4	0/0/0/0
5	GOL	A	5006	-	-	0/4/4/4	0/0/0/0
4	SO4	B	4001	-	-	0/0/0/0	0/0/0/0
5	GOL	B	5004	-	-	0/4/4/4	0/0/0/0
5	GOL	B	5007	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
5	B	5007	GOL	O3-C3-C2	2.42	121.84	110.11

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6 monomers are involved in 15 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	5001	GOL	2	0
5	A	5002	GOL	1	0
5	A	5003	GOL	1	0
5	A	5006	GOL	5	0
5	B	5004	GOL	5	0
5	B	5007	GOL	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	275/275 (100%)	-0.21	3 (1%) 80 80	8, 16, 37, 58	0
2	B	99/99 (100%)	-0.21	3 (3%) 50 51	9, 17, 38, 48	0
3	C	9/9 (100%)	-0.27	0 100 100	11, 18, 29, 37	0
All	All	383/383 (100%)	-0.21	6 (1%) 72 72	8, 16, 38, 58	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	1	ILE	3.4
2	B	48	LYS	3.2
2	B	75	LYS	2.6
1	A	275	GLU	2.4
1	A	90	ASP	2.2
1	A	174	ASN	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	GOL	A	5001	6/6	0.76	0.27	39,44,48,49	0
5	GOL	A	5002	6/6	0.79	0.32	49,51,53,60	0
4	SO4	A	4002	5/5	0.86	0.15	14,17,20,24	5
5	GOL	B	5004	6/6	0.87	0.29	29,49,51,54	0
5	GOL	B	5007	6/6	0.90	0.27	29,36,41,59	0
5	GOL	A	5003	6/6	0.91	0.18	39,44,46,64	0
5	GOL	A	5005	6/6	0.92	0.25	19,45,57,69	0
5	GOL	A	5006	6/6	0.93	0.23	31,32,39,53	0
4	SO4	B	4001	5/5	0.95	0.11	25,30,36,42	5
4	SO4	A	4003	5/5	0.99	0.07	25,28,36,37	5

## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.