



wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 17, 2023 – 11:53 am GMT

PDB ID : 2X7L
Title : Implications of the HIV-1 Rev dimer structure at 3.2A resolution for multi-
meric binding to the Rev response element
Authors : DiMattia, M.A.; Watts, N.R.; Stahl, S.J.; Rader, C.; Wingfield, P.T.; Stuart,
D.I.; Steven, A.C.; Grimes, J.M.
Deposited on : 2010-03-01
Resolution : 3.17 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.17 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1467 (3.20-3.16)
Clashscore	141614	1599 (3.20-3.16)
Ramachandran outliers	138981	1574 (3.20-3.16)
Sidechain outliers	138945	1573 (3.20-3.16)
RSRZ outliers	127900	1423 (3.20-3.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	231	
1	C	231	
1	E	231	
1	G	231	
1	H	231	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	J	231	<p>4% 68% 26% . .</p>
2	B	217	<p>65% 29% 5% .</p>
2	D	217	<p>2% 64% 30% 6% .</p>
2	F	217	<p>65% 29% 5% .</p>
2	I	217	<p>65% 29% 5% .</p>
2	K	217	<p>64% 30% 6% .</p>
2	L	217	<p>65% 30% 5% .</p>
3	M	115	<p>3% 36% 12% . 50%</p>
3	N	115	<p>34% 13% . 50%</p>
3	O	115	<p>3% 34% 14% . 50%</p>
3	P	115	<p>4% 33% 15% . 50%</p>
3	Q	115	<p>2% 30% 16% . 50%</p>
3	R	115	<p>36% 12% . 50%</p>

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 22554 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called FAB HEAVY CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	221	Total 1628	C 1033	N 267	O 323	S 5	0	0	1
1	C	221	Total 1628	C 1033	N 267	O 323	S 5	0	0	1
1	E	221	Total 1628	C 1033	N 267	O 323	S 5	0	0	1
1	G	221	Total 1628	C 1033	N 267	O 323	S 5	0	0	1
1	H	221	Total 1628	C 1033	N 267	O 323	S 5	0	0	1
1	J	221	Total 1628	C 1033	N 267	O 323	S 5	0	0	1

- Molecule 2 is a protein called FAB LIGHT CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	217	Total 1639	C 1026	N 269	O 337	S 7	0	0	0
2	D	217	Total 1639	C 1026	N 269	O 337	S 7	0	0	0
2	F	217	Total 1639	C 1026	N 269	O 337	S 7	0	0	0
2	I	217	Total 1639	C 1026	N 269	O 337	S 7	0	0	0
2	K	217	Total 1639	C 1026	N 269	O 337	S 7	0	0	0
2	L	217	Total 1639	C 1026	N 269	O 337	S 7	0	0	0


- Molecule 3 is a protein called PROTEIN REV.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	M	57	Total 492	C 303	N 105	O 84	0	0	0
3	N	57	Total 492	C 303	N 105	O 84	0	0	0
3	O	57	Total 492	C 303	N 105	O 84	0	0	0
3	P	57	Total 492	C 303	N 105	O 84	0	0	0
3	Q	57	Total 492	C 303	N 105	O 84	0	0	0
3	R	57	Total 492	C 303	N 105	O 84	0	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: FAB HEAVY CHAIN




- Molecule 1: FAB HEAVY CHAIN




- Molecule 1: FAB HEAVY CHAIN




• Molecule 1: FAB HEAVY CHAIN




• Molecule 1: FAB HEAVY CHAIN



• Molecule 1: FAB HEAVY CHAIN




• Molecule 2: FAB LIGHT CHAIN




• Molecule 2: FAB LIGHT CHAIN

Chain D:  2% 64% 30% 6%




• Molecule 2: FAB LIGHT CHAIN

Chain F:  65% 29% 5%




• Molecule 2: FAB LIGHT CHAIN

Chain I:  65% 29% 5%




• Molecule 2: FAB LIGHT CHAIN


Chain K:  64% 30% 6%





• Molecule 2: FAB LIGHT CHAIN

Chain L:  65% 30% 5%









- Molecule 3: PROTEIN REV






- Molecule 3: PROTEIN REV





- Molecule 3: PROTEIN REV



- Molecule 3: PROTEIN REV




- Molecule 3: PROTEIN REV



LEU
ASP
CYS
ASN
GLY
ASP
CYS
GLY
THR
SER
GLY
THR
GLN
GLY
VAL
GLY
SER
PRO
GLN
ILE
LEU
VAL
GLY
SER
PRO
THR
VAL
LEU
GLY
GLY
ALA
LYS
GLU

● Molecule 3: PROTEIN REV



ALA
GLY
ARG
SER
GLN
GLY
ASP
SER
D9
SER
E10
D11
L12
L13
V16
K20
F21
L22
P27
P28
P29
N30
T34
Q49
I55
I59
Y63
L64
G65
ARG
SER
ALA
GLU
PRO
VAL
PRO
LEU
GLN
LEU
PRO
PRO
LEU
GLY
ARG
LEU
THR
LEU
ASP
CYS
ASN
GLU
ASP
CYS
GLY

THR
SER
GLY
THR
GLN
GLY
VAL
GLY
SER
PRO
GLN
ILE
LEU
VAL
GLU
SER
PRO
THR
VAL
LEU
GLU
SER
GLY
ALA
LYS
GLU

4 Data and refinement statistics i

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	87.68Å 87.69Å 176.33Å 94.86° 95.50° 104.60°	Depositor
Resolution (Å)	48.78 – 3.17 48.77 – 3.17	Depositor EDS
% Data completeness (in resolution range)	(Not available) (48.78-3.17) 97.6 (48.77-3.17)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.74 (at 3.19Å)	Xtrriage
Refinement program	BUSTER 2.9.2	Depositor
R, R_{free}	0.234 , 0.250 0.253 , 0.265	Depositor DCC
R_{free} test set	4187 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	85.2	Xtrriage
Anisotropy	0.532	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.21 , 36.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.327 for -k,-h,-l	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	22554	wwPDB-VP
Average B, all atoms (Å ²)	143.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.38% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.62	0/1671	0.83	0/2288
1	C	0.46	0/1671	0.78	0/2288
1	E	0.59	0/1671	0.81	0/2288
1	G	0.47	0/1671	0.78	0/2288
1	H	0.56	0/1671	0.81	0/2288
1	J	0.54	0/1671	0.80	0/2288
2	B	0.60	0/1675	0.85	0/2278
2	D	0.50	0/1675	0.81	0/2278
2	F	0.59	0/1675	0.85	0/2278
2	I	0.49	0/1675	0.82	0/2278
2	K	0.57	0/1675	0.84	0/2278
2	L	0.59	0/1675	0.84	0/2278
3	M	0.53	0/501	0.69	0/672
3	N	0.55	0/501	0.68	0/672
3	O	0.60	0/501	0.72	0/672
3	P	0.52	0/501	0.68	0/672
3	Q	0.55	0/501	0.68	0/672
3	R	0.57	0/501	0.70	0/672
All	All	0.55	0/23082	0.80	0/31428

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1628	0	1594	29	0
1	C	1628	0	1594	25	0
1	E	1628	0	1594	28	0
1	G	1628	0	1594	24	0
1	H	1628	0	1594	29	0
1	J	1628	0	1594	28	0
2	B	1639	0	1581	40	0
2	D	1639	0	1581	43	0
2	F	1639	0	1581	41	0
2	I	1639	0	1581	40	0
2	K	1639	0	1581	42	0
2	L	1639	0	1581	39	0
3	M	492	0	504	7	0
3	N	492	0	504	9	0
3	O	492	0	504	7	0
3	P	492	0	504	11	0
3	Q	492	0	504	13	0
3	R	492	0	504	7	0
All	All	22554	0	22074	434	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 434 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:I:21:ILE:HG21	2:I:105:THR:HG21	1.26	1.14
2:L:21:ILE:HG21	2:L:105:THR:HG21	1.28	1.12
2:D:21:ILE:HG21	2:D:105:THR:HG21	1.24	1.11
2:K:21:ILE:HG21	2:K:105:THR:HG21	1.28	1.10
2:B:21:ILE:HG21	2:B:105:THR:HG21	1.27	1.09

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	219/231 (95%)	187 (85%)	28 (13%)	4 (2%)	8	38
1	C	219/231 (95%)	189 (86%)	26 (12%)	4 (2%)	8	38
1	E	219/231 (95%)	187 (85%)	28 (13%)	4 (2%)	8	38
1	G	219/231 (95%)	190 (87%)	25 (11%)	4 (2%)	8	38
1	H	219/231 (95%)	187 (85%)	28 (13%)	4 (2%)	8	38
1	J	219/231 (95%)	190 (87%)	25 (11%)	4 (2%)	8	38
2	B	215/217 (99%)	188 (87%)	14 (6%)	13 (6%)	1	11
2	D	215/217 (99%)	188 (87%)	14 (6%)	13 (6%)	1	11
2	F	215/217 (99%)	188 (87%)	14 (6%)	13 (6%)	1	11
2	I	215/217 (99%)	188 (87%)	14 (6%)	13 (6%)	1	11
2	K	215/217 (99%)	188 (87%)	14 (6%)	13 (6%)	1	11
2	L	215/217 (99%)	188 (87%)	15 (7%)	12 (6%)	2	12
3	M	55/115 (48%)	46 (84%)	7 (13%)	2 (4%)	3	21
3	N	55/115 (48%)	48 (87%)	5 (9%)	2 (4%)	3	21
3	O	55/115 (48%)	47 (86%)	6 (11%)	2 (4%)	3	21
3	P	55/115 (48%)	46 (84%)	7 (13%)	2 (4%)	3	21
3	Q	55/115 (48%)	46 (84%)	7 (13%)	2 (4%)	3	21
3	R	55/115 (48%)	47 (86%)	6 (11%)	2 (4%)	3	21
All	All	2934/3378 (87%)	2538 (86%)	283 (10%)	113 (4%)	3	19

5 of 113 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	174	SER
2	D	174	SER
2	F	174	SER
2	I	174	SER
2	K	174	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	182/192 (95%)	152 (84%)	30 (16%)	2	10
1	C	182/192 (95%)	154 (85%)	28 (15%)	2	12
1	E	182/192 (95%)	151 (83%)	31 (17%)	2	9
1	G	182/192 (95%)	154 (85%)	28 (15%)	2	12
1	H	182/192 (95%)	152 (84%)	30 (16%)	2	10
1	J	182/192 (95%)	152 (84%)	30 (16%)	2	10
2	B	184/185 (100%)	154 (84%)	30 (16%)	2	10
2	D	184/185 (100%)	153 (83%)	31 (17%)	2	9
2	F	184/185 (100%)	155 (84%)	29 (16%)	2	11
2	I	184/185 (100%)	154 (84%)	30 (16%)	2	10
2	K	184/185 (100%)	154 (84%)	30 (16%)	2	10
2	L	184/185 (100%)	154 (84%)	30 (16%)	2	10
3	M	53/101 (52%)	46 (87%)	7 (13%)	4	17
3	N	53/101 (52%)	45 (85%)	8 (15%)	3	13
3	O	53/101 (52%)	45 (85%)	8 (15%)	3	13
3	P	53/101 (52%)	46 (87%)	7 (13%)	4	17
3	Q	53/101 (52%)	45 (85%)	8 (15%)	3	13
3	R	53/101 (52%)	46 (87%)	7 (13%)	4	17
All	All	2514/2868 (88%)	2112 (84%)	402 (16%)	2	10

5 of 402 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	183	SER
1	J	144	CYS
3	R	34	THR
2	I	2	LEU
2	I	135	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 37 such sidechains are listed below:

Mol	Chain	Res	Type
2	L	127	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	Q	51	GLN
2	L	169	GLN
3	N	51	GLN
2	F	79	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	221/231 (95%)	-0.27	3 (1%) 75 63	75, 114, 192, 246	0
1	C	221/231 (95%)	0.11	17 (7%) 13 7	102, 173, 235, 263	0
1	E	221/231 (95%)	-0.31	4 (1%) 68 55	78, 118, 191, 251	0
1	G	221/231 (95%)	0.14	12 (5%) 25 14	100, 175, 246, 265	0
1	H	221/231 (95%)	-0.29	4 (1%) 68 55	67, 113, 195, 224	0
1	J	221/231 (95%)	-0.23	9 (4%) 37 23	74, 119, 201, 236	0
2	B	217/217 (100%)	-0.45	1 (0%) 91 86	73, 116, 189, 208	0
2	D	217/217 (100%)	-0.23	5 (2%) 60 46	98, 156, 248, 263	0
2	F	217/217 (100%)	-0.39	1 (0%) 91 86	74, 113, 200, 229	0
2	I	217/217 (100%)	-0.29	2 (0%) 84 75	88, 149, 251, 283	0
2	K	217/217 (100%)	-0.48	1 (0%) 91 86	75, 116, 188, 222	0
2	L	217/217 (100%)	-0.45	0 100 100	76, 116, 186, 211	0
3	M	57/115 (49%)	-0.10	3 (5%) 26 14	109, 164, 278, 284	0
3	N	57/115 (49%)	-0.26	1 (1%) 68 55	96, 155, 287, 298	0
3	O	57/115 (49%)	0.04	3 (5%) 26 14	97, 151, 278, 287	0
3	P	57/115 (49%)	0.01	5 (8%) 10 5	119, 165, 276, 280	0
3	Q	57/115 (49%)	0.03	2 (3%) 44 28	98, 151, 275, 277	0
3	R	57/115 (49%)	0.06	0 100 100	95, 156, 274, 280	0
All	All	2970/3378 (87%)	-0.23	73 (2%) 57 43	67, 129, 236, 298	0

The worst 5 of 73 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	J	134	SER	11.9
1	C	221	ASP	8.0
1	C	134	SER	7.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	D	123	PRO	6.8
1	H	134	SER	6.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.