



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 16, 2024 – 03:26 AM EST

PDB ID : 6X9I
Title : Human DNMT1(729-1600) Bound to Zebularine-Containing 12mer dsDNA
Authors : Pathuri, S.; Horton, J.R.; Cheng, X.
Deposited on : 2020-06-02
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.21
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.004 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.40


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	5791 (2.20-2.20)
Clashscore	180529	6634 (2.20-2.20)
Ramachandran outliers	177936	6560 (2.20-2.20)
Sidechain outliers	177891	6561 (2.20-2.20)
RSRZ outliers	164620	5791 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	874	
2	C	12	
3	D	12	

2 Entry composition i

There are 8 unique types of molecules in this entry. The entry contains 6956 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA (cytosine-5)-methyltransferase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	815	6194	3935	1088	1129	42	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	727	HIS	-	expression tag	UNP P26358
A	728	MET	-	expression tag	UNP P26358

- Molecule 2 is a DNA chain called DNA (5'-D(*GP*AP*GP*GP*CP*(5CM)P*GP*CP*CP*TP*GP*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	C	12	244	116	47	70	11	0	0	0

- Molecule 3 is a DNA chain called DNA (5'-D(*GP*CP*AP*GP*G)-R(P*(PYO))-D(P*GP*GP*CP*CP*TP*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
3	D	12	243	115	46	71	11	0	0	0

- Molecule 4 is S-ADENOSYL-L-HOMOCYSTEINE (three-letter code: SAH) (formula: C₁₄H₂₀N₆O₅S).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	S	0	0
			26	14	6	5	1		

- Molecule 5 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	2	Total	Zn	0	0
			2	2		

- Molecule 6 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			4	2	2		
6	A	1	Total	C	O	0	0
			4	2	2		
6	A	1	Total	C	O	0	0
			4	2	2		
6	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 7 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	208	Total	O	0	0
			208	208		
8	C	4	Total	O	0	0
			4	4		
8	D	13	Total	O	0	0
			13	13		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA (cytosine-5)-methyltransferase 1



- Molecule 2: DNA (5'-D(*GP*AP*GP*GP*CP*(5CM)P*GP*CP*CP*TP*GP*C)-3')



- Molecule 3: DNA (5'-D(*GP*CP*AP*GP*G)-R(P*(PYO))-D(P*GP*GP*CP*CP*TP*C)-3')



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	161.89Å 77.70Å 115.41Å 90.00° 125.67° 90.00°	Depositor
Resolution (Å)	45.21 – 2.20 45.21 – 2.20	Depositor EDS
% Data completeness (in resolution range)	93.5 (45.21-2.20) 93.5 (45.21-2.20)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.14 (at 2.20Å)	Xtrriage
Refinement program	PHENIX 1.17.1_3660	Depositor
R, R_{free}	0.194 , 0.225 0.196 , 0.229	Depositor DCC
R_{free} test set	56931 reflections (3.62%)	wwPDB-VP
Wilson B-factor (Å ²)	45.6	Xtrriage
Anisotropy	0.337	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 56.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6956	wwPDB-VP
Average B, all atoms (Å ²)	70.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.62% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, PYO, EDO, SAH, 5CM, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.29	0/6356	0.48	0/8654
2	C	0.65	0/250	0.84	0/382
3	D	0.56	0/250	0.82	0/382
All	All	0.32	0/6856	0.51	0/9418

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6194	0	5780	34	0
2	C	244	0	137	1	0
3	D	243	0	133	2	0
4	A	26	0	19	0	0
5	A	2	0	0	0	0
6	A	16	0	24	0	0
7	A	6	0	8	0	0
8	A	208	0	0	0	0
8	C	4	0	0	0	0
8	D	13	0	0	0	0
All	All	6956	0	6101	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1232:MET:HG3	3:D:17:DG:H2'	1.58	0.83
1:A:1015:LYS:NZ	1:A:1052:ASP:OD1	2.22	0.72
1:A:1415:LYS:HE2	1:A:1417:MET:HE1	1.76	0.66
1:A:995:ARG:NH2	1:A:1062:VAL:O	2.35	0.60
1:A:749:LYS:HG3	1:A:758:GLU:HA	1.85	0.57
1:A:1464:ASN:O	1:A:1473:ARG:NH1	2.36	0.57
1:A:793:HIS:ND1	1:A:822:ASP:OD1	2.39	0.56
1:A:1006:ASN:OD1	1:A:1007:GLY:N	2.38	0.56
1:A:1203:THR:HG23	1:A:1204:THR:HG23	1.90	0.54
1:A:790:GLN:HB3	1:A:825:LEU:HD12	1.90	0.52
1:A:1154:GLY:HA3	1:A:1585:ALA:HB3	1.90	0.52
1:A:1535:LYS:HE3	2:C:7:DG:O6	2.10	0.51
1:A:1455:LEU:HB2	1:A:1494:THR:OG1	2.12	0.49
1:A:866:GLN:HG3	1:A:867:LEU:HG	1.95	0.49
1:A:1514:TYR:HE2	1:A:1533:MET:HG3	1.79	0.47
1:A:1392:PRO:HB3	1:A:1401:ARG:HD3	1.97	0.47
1:A:941:LEU:HB2	1:A:992:ARG:HB2	1.97	0.46
1:A:1187:PHE:HB3	1:A:1189:GLU:HG3	1.97	0.46
1:A:1446:LEU:HD21	1:A:1452:ALA:HB2	1.99	0.45
1:A:1500:LEU:HB2	1:A:1501:PRO:HD3	1.98	0.45
1:A:971:GLU:OE1	1:A:974:ARG:HD2	2.17	0.44
1:A:1414:CYS:SG	1:A:1546:ARG:HD2	2.58	0.44
1:A:1155:PHE:HZ	1:A:1264:LEU:HD13	1.82	0.44
1:A:1476:CYS:SG	1:A:1477:SER:N	2.92	0.43
1:A:837:LYS:HG2	1:A:864:PHE:HB2	2.00	0.43
1:A:775:TYR:HB3	1:A:796:TRP:CZ3	2.54	0.42
1:A:913:GLN:HG2	1:A:923:TYR:CE2	2.54	0.42
1:A:1419:ALA:HB1	1:A:1446:LEU:HB3	2.01	0.42
1:A:777:ALA:HB2	1:A:796:TRP:CE3	2.55	0.42
1:A:898:ARG:H	1:A:898:ARG:HG2	1.68	0.41
1:A:1269:ARG:O	1:A:1272:VAL:HG22	2.21	0.41
3:D:21:DC:H2''	3:D:22:DC:C6	2.56	0.41
1:A:1263:PHE:HB3	1:A:1317:ALA:HB3	2.03	0.40
1:A:1002:PRO:HD2	1:A:1013:ASP:O	2.21	0.40
1:A:1441:ASN:O	1:A:1454:LYS:HE3	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	805/874 (92%)	782 (97%)	23 (3%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	621/753 (82%)	614 (99%)	7 (1%)	70 82

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	826	SER
1	A	920	ARG
1	A	932	LEU
1	A	1045	SER
1	A	1207	ARG
1	A	1240	TYR
1	A	1536	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	5CM	C	6	2,3	18,21,22	1.69	6 (33%)	24,30,33	1.69	7 (29%)
3	PYO	D	18	3	16,20,21	1.20	2 (12%)	21,28,31	0.71	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	5CM	C	6	2,3	-	5/7/21/22	0/2/2/2
3	PYO	D	18	3	-	4/7/25/26	0/2/2/2

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	6	5CM	O2-C2	-3.60	1.17	1.23
2	C	6	5CM	C2-N1	3.26	1.46	1.40
3	D	18	PYO	C2-N1	3.09	1.46	1.40
2	C	6	5CM	C6-C5	2.98	1.39	1.34
2	C	6	5CM	C4-N3	2.25	1.37	1.34
2	C	6	5CM	C5-C4	-2.23	1.42	1.44
3	D	18	PYO	O2-C2	-2.21	1.19	1.23
2	C	6	5CM	C6-N1	2.11	1.41	1.38

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	6	5CM	O4'-C1'-N1	3.75	114.52	107.86
2	C	6	5CM	C6-N1-C2	-3.07	116.86	120.95
2	C	6	5CM	C1'-N1-C2	2.89	122.78	117.83
2	C	6	5CM	C5A-C5-C6	-2.73	119.16	122.85
2	C	6	5CM	C4-N3-C2	-2.25	117.68	120.81
2	C	6	5CM	C5-C6-N1	-2.23	120.89	123.31
2	C	6	5CM	N1-C2-N3	2.01	122.30	118.80

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	D	18	PYO	C2'-C1'-N1-C6
3	D	18	PYO	C2'-C1'-N1-C2
2	C	6	5CM	O4'-C1'-N1-C6
2	C	6	5CM	C2'-C1'-N1-C6
3	D	18	PYO	O4'-C1'-N1-C6
2	C	6	5CM	C3'-C4'-C5'-O5'
3	D	18	PYO	O4'-C1'-N1-C2
2	C	6	5CM	C2'-C1'-N1-C2
2	C	6	5CM	O4'-C1'-N1-C2

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 2 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	EDO	A	1707	-	3,3,3	0.46	0	2,2,2	0.32	0
7	GOL	A	1705	-	5,5,5	0.92	0	5,5,5	1.05	0
6	EDO	A	1708	-	3,3,3	0.43	0	2,2,2	0.38	0
6	EDO	A	1704	-	3,3,3	0.46	0	2,2,2	0.40	0
4	SAH	A	1701	-	23,28,28	0.87	1 (4%)	22,40,40	1.07	1 (4%)
6	EDO	A	1706	-	3,3,3	0.43	0	2,2,2	0.37	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	EDO	A	1707	-	-	1/1/1/1	-
7	GOL	A	1705	-	-	2/4/4/4	-
6	EDO	A	1708	-	-	1/1/1/1	-
6	EDO	A	1704	-	-	0/1/1/1	-
4	SAH	A	1701	-	-	0/11/31/31	0/3/3/3
6	EDO	A	1706	-	-	0/1/1/1	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1701	SAH	C1'-N9	-2.16	1.44	1.49

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1701	SAH	O4'-C1'-N9	-3.56	104.03	108.75

There are no chirality outliers.


All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	A	1705	GOL	O1-C1-C2-C3
7	A	1705	GOL	O1-C1-C2-O2
6	A	1707	EDO	O1-C1-C2-O2
6	A	1708	EDO	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	815/874 (93%)	0.57	93 (11%) 11 9	30, 66, 126, 158	0
2	C	11/12 (91%)	0.48	0 100 100	68, 87, 126, 142	0
3	D	11/12 (91%)	0.13	0 100 100	62, 78, 127, 142	0
All	All	837/898 (93%)	0.57	93 (11%) 12 9	30, 66, 126, 158	0

All (93) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1477	SER	5.0
1	A	754	ALA	4.2
1	A	1456	ARG	3.9
1	A	1007	GLY	3.8
1	A	890	PHE	3.7
1	A	753	ASP	3.6
1	A	787	SER	3.6
1	A	884	PRO	3.6
1	A	1486	ASP	3.5
1	A	976	TYR	3.4
1	A	1005	SER	3.4
1	A	1441	ASN	3.3
1	A	950	ILE	3.3
1	A	963	PRO	3.3
1	A	1489	ALA	3.2
1	A	1600	ALA	3.2
1	A	1240	TYR	3.2
1	A	889	LYS	3.2
1	A	1450	THR	3.2
1	A	1492	PHE	3.2
1	A	731	ILE	3.2
1	A	1453	ARG	3.1
1	A	1481	ALA	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	1457	TYR	3.1
1	A	1465	GLY	3.0
1	A	738	VAL	3.0
1	A	1502	HIS	3.0
1	A	1448	ASP	3.0
1	A	1239	THR	3.0
1	A	756	THR	3.0
1	A	755	GLU	3.0
1	A	1462	ARG	2.9
1	A	732	SER	2.9
1	A	1464	ASN	2.9
1	A	1452	ALA	2.8
1	A	1472	LEU	2.8
1	A	1475	VAL	2.8
1	A	1455	LEU	2.8
1	A	1485	CYS	2.8
1	A	967	ASP	2.8
1	A	1443	GLU	2.8
1	A	751	CYS	2.8
1	A	745	SER	2.6
1	A	740	THR	2.6
1	A	1471	ALA	2.6
1	A	783	TRP	2.6
1	A	1478	CYS	2.6
1	A	730	ARG	2.6
1	A	979	TYR	2.5
1	A	1460	HIS	2.5
1	A	1498	TRP	2.5
1	A	850	ASP	2.5
1	A	892	PHE	2.5
1	A	1488	ALA	2.5
1	A	750	VAL	2.4
1	A	779	VAL	2.4
1	A	741	ASP	2.4
1	A	1466	ARG	2.4
1	A	1103	ALA	2.4
1	A	1006	ASN	2.4
1	A	1479	VAL	2.3
1	A	835	ILE	2.3
1	A	918	ASP	2.3
1	A	789	GLY	2.3
1	A	769	ASP	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	968	LEU	2.3
1	A	964	VAL	2.2
1	A	1458	THR	2.2
1	A	1442	ILE	2.2
1	A	737	ALA	2.2
1	A	917	LEU	2.2
1	A	916	ASP	2.2
1	A	978	ASP	2.2
1	A	1491	GLN	2.2
1	A	1104	ARG	2.1
1	A	1490	ARG	2.1
1	A	1497	PRO	2.1
1	A	775	TYR	2.1
1	A	948	PHE	2.1
1	A	1531	GLU	2.1
1	A	1467	SER	2.1
1	A	1496	ILE	2.1
1	A	788	ASN	2.1
1	A	888	ASN	2.1
1	A	965	ASP	2.1
1	A	1459	HIS	2.1
1	A	734	VAL	2.0
1	A	1444	VAL	2.0
1	A	786	SER	2.0
1	A	772	LYS	2.0
1	A	891	LYS	2.0
1	A	1136	LEU	2.0
1	A	1487	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	5CM	C	6	20/21	0.94	0.12	78,81,92,103	0
3	PYO	D	18	19/20	0.96	0.09	39,50,59,60	0

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
6	EDO	A	1704	4/4	0.82	0.16	63,71,71,76	0
7	GOL	A	1705	6/6	0.82	0.22	65,68,79,80	0
6	EDO	A	1708	4/4	0.84	0.28	51,56,57,63	0
6	EDO	A	1706	4/4	0.88	0.14	78,80,80,81	0
6	EDO	A	1707	4/4	0.92	0.13	61,64,68,71	0
5	ZN	A	1702	1/1	0.95	0.14	89,89,89,89	0
4	SAH	A	1701	26/26	0.96	0.07	33,39,48,50	0
5	ZN	A	1703	1/1	0.99	0.06	78,78,78,78	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.