

# wwPDB X-ray Structure Validation Summary Report (i)

#### Nov 26, 2024 – 06:10 PM JST

PDB ID	:	8XAA
Title	:	Structure of NAP1 in complex with H2A-H2B
Authors	:	Li, X.
Deposited on	:	2023-12-03
Resolution	:	3.35  Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.21
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.004 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.40

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\text{-}RAY \, DIFFRACTION$ 

The reported resolution of this entry is 3.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
$R_{free}$	164625	1012 (3.40-3.32)
Clashscore	180529	1035 (3.40-3.32)
Ramachandran outliers	177936	1037 (3.40-3.32)
Sidechain outliers	177891	1037 (3.40-3.32)
RSRZ outliers	164620	1012 (3.40-3.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain					
1	А	308	69%	22%	9%			
1	В	308	66%	21%	13%			
1	С	308	62%	27%	11%			
1	D	308	59%	25%	16%			
1	Е	308	59%	22%	19%			
1	F	308	60%	22%	18%			



Mol	Chain	Length	Quality of chain					
1	G	308	60%	24%	16%			
1	Н	308	61%	23%	16%			
2	Ι	92	70%	23%	8%			
2	K	92	50%	39%	11%			
3	J	99	71%	19%	10%			
3	L	99	% 60%	31%	9%			



# 2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 19338 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	Δ	200	Total	С	Ν	0	S	0	0	0
1	A	280	2217	1420	359	434	4	0		0
1	р	260	Total	С	Ν	0	S	0	0	0
1	D	209	2136	1365	344	422	5	0	0	0
1	F	240	Total	С	Ν	0	S	0	0	0
1	Ľ	249	1962	1260	310	388	4	0		
1	F	254	Total	С	Ν	0	S	0	0	0
1	I.	204	2018	1301	326	387	4		0	0
1	С	973	Total	С	Ν	0	S	0	0	0
1	U	210	2177	1387	352	435	3	0	0	0
1	П	260	Total	С	Ν	Ο	$\mathbf{S}$	0	0	0
1	D	200	2048	1315	329	400	4	0	0	0
1	C	250	Total	С	Ν	Ο	$\mathbf{S}$	0	0	0
1	G	239	2054	1319	326	405	4	0	0	0
1	Ц	258	Total	С	Ν	0	S	0	0	0
1	11	200	2032	1307	323	398	4		U	

• Molecule 1 is a protein called Nucleosome Assembly Protein.

There are 168 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	-11	MET	-	initiating methionine	UNP Q19007
А	-10	GLY	-	expression tag	UNP Q19007
А	-9	SER	-	expression tag	UNP Q19007
А	-8	SER	-	expression tag	UNP Q19007
А	-7	HIS	-	expression tag	UNP Q19007
А	-6	HIS	-	expression tag	UNP Q19007
А	-5	HIS	-	expression tag	UNP Q19007
А	-4	HIS	-	expression tag	UNP Q19007
А	-3	HIS	-	expression tag	UNP Q19007
А	-2	HIS	-	expression tag	UNP Q19007
A	-1	SER	-	expression tag	UNP Q19007
А	0	SER	-	expression tag	UNP Q19007
А	1	GLY	-	expression tag	UNP Q19007



8XAA	
------	--

Chain	Residue	Modelled	Actual	Comment	Reference
A	2	GLU	-	expression tag	UNP Q19007
A	3	ASN	-	expression tag	UNP Q19007
A	4	LEU	-	expression tag	UNP Q19007
A	5	TYR	-	expression tag	UNP Q19007
A	6	PHE	-	expression tag	UNP Q19007
A	7	GLN	_	expression tag	UNP Q19007
А	8	HIS	_	expression tag	UNP Q19007
А	9	MET	-	expression tag	UNP Q19007
В	-11	MET	-	initiating methionine	UNP Q19007
В	-10	GLY	-	expression tag	UNP Q19007
В	-9	SER	-	expression tag	UNP Q19007
В	-8	SER	-	expression tag	UNP Q19007
В	-7	HIS	-	expression tag	UNP Q19007
В	-6	HIS	-	expression tag	UNP Q19007
В	-5	HIS	-	expression tag	UNP Q19007
В	-4	HIS	-	expression tag	UNP Q19007
В	-3	HIS	-	expression tag	UNP Q19007
В	-2	HIS	-	expression tag	UNP Q19007
В	-1	SER	-	expression tag	UNP Q19007
В	0	SER	-	expression tag	UNP Q19007
В	1	GLY	-	expression tag	UNP Q19007
В	2	GLU	-	expression tag	UNP Q19007
В	3	ASN	-	expression tag	UNP Q19007
В	4	LEU	-	expression tag	UNP Q19007
В	5	TYR	-	expression tag	UNP Q19007
В	6	PHE	-	expression tag	UNP Q19007
В	7	GLN	-	expression tag	UNP Q19007
В	8	HIS	-	expression tag	UNP Q19007
В	9	MET	-	expression tag	UNP Q19007
E	-11	MET	-	initiating methionine	UNP Q19007
E	-10	GLY	-	expression tag	UNP Q19007
E	-9	SER	-	expression tag	UNP Q19007
Е	-8	SER	-	expression tag	UNP Q19007
E	-7	HIS	-	expression tag	UNP Q19007
E	-6	HIS	-	expression tag	UNP Q19007
E	-5	HIS	-	expression tag	UNP Q19007
E	-4	HIS	-	expression tag	UNP Q19007
E	-3	HIS	-	expression tag	UNP Q19007
E	-2	HIS	-	expression tag	UNP Q19007
E	-1	SER	-	expression tag	UNP Q19007
E	0	SER	-	expression tag	UNP Q19007
Е	1	GLY	-	expression tag	UNP Q19007



Chain	Residue	Modelled	Actual	Comment	Reference
Е	2	GLU	-	expression tag	UNP Q19007
E	3	ASN	-	expression tag	UNP Q19007
Е	4	LEU	-	expression tag	UNP Q19007
E	5	TYR	-	expression tag	UNP Q19007
Е	6	PHE	_	expression tag	UNP Q19007
E	7	GLN	-	expression tag	UNP Q19007
Е	8	HIS	-	expression tag	UNP Q19007
Е	9	MET	-	expression tag	UNP Q19007
F	-11	MET	-	initiating methionine	UNP Q19007
F	-10	GLY	-	expression tag	UNP Q19007
F	-9	SER	-	expression tag	UNP Q19007
F	-8	SER	-	expression tag	UNP Q19007
F	-7	HIS	-	expression tag	UNP Q19007
F	-6	HIS	-	expression tag	UNP Q19007
F	-5	HIS	-	expression tag	UNP Q19007
F	-4	HIS	-	expression tag	UNP Q19007
F	-3	HIS	-	expression tag	UNP Q19007
F	-2	HIS	-	expression tag	UNP Q19007
F	-1	SER	-	expression tag	UNP Q19007
F	0	SER	-	expression tag	UNP Q19007
F	1	GLY	-	expression tag	UNP Q19007
F	2	GLU	-	expression tag	UNP Q19007
F	3	ASN	-	expression tag	UNP Q19007
F	4	LEU	-	expression tag	UNP Q19007
F	5	TYR	-	expression tag	UNP Q19007
F	6	PHE	-	expression tag	UNP Q19007
F	7	GLN	-	expression tag	UNP Q19007
F	8	HIS	-	expression tag	UNP Q19007
F	9	MET	-	expression tag	UNP Q19007
C	-11	MET	-	initiating methionine	UNP Q19007
C	-10	GLY	-	expression tag	UNP Q19007
C	-9	SER	-	expression tag	UNP Q19007
C	-8	SER	-	expression tag	UNP Q19007
C	-7	HIS	-	expression tag	UNP Q19007
C	-6	HIS	-	expression tag	UNP Q19007
C	-5	HIS	-	expression tag	UNP Q19007
C	-4	HIS	-	expression tag	UNP Q19007
C	-3	HIS	-	expression tag	UNP Q19007
C	-2	HIS	-	expression tag	UNP Q19007
C	-1	SER	-	expression tag	UNP Q19007
C	0	SER	-	expression tag	UNP Q19007
C	1	GLY	-	expression tag	UNP Q19007



8XAA

Chain	Residue	Modelled	Actual	Comment	Reference
C	2	GLU	_	expression tag	UNP Q19007
C	3	ASN	_	expression tag	UNP Q19007
C	4	LEU	_	expression tag	UNP Q19007
C	5	TYR	_	expression tag	UNP Q19007
C	6	PHE	-	expression tag	UNP Q19007
С	7	GLN	-	expression tag	UNP Q19007
С	8	HIS	-	expression tag	UNP Q19007
С	9	MET	-	expression tag	UNP Q19007
D	-11	MET	-	initiating methionine	UNP Q19007
D	-10	GLY	_	expression tag	UNP Q19007
D	-9	SER	-	expression tag	UNP Q19007
D	-8	SER	-	expression tag	UNP Q19007
D	-7	HIS	-	expression tag	UNP Q19007
D	-6	HIS	-	expression tag	UNP Q19007
D	-5	HIS	-	expression tag	UNP Q19007
D	-4	HIS	-	expression tag	UNP Q19007
D	-3	HIS	-	expression tag	UNP Q19007
D	-2	HIS	-	expression tag	UNP Q19007
D	-1	SER	-	expression tag	UNP Q19007
D	0	SER	-	expression tag	UNP Q19007
D	1	GLY	-	expression tag	UNP Q19007
D	2	GLU	-	expression tag	UNP Q19007
D	3	ASN	-	expression tag	UNP Q19007
D	4	LEU	-	expression tag	UNP Q19007
D	5	TYR	-	expression tag	UNP Q19007
D	6	PHE	-	expression tag	UNP Q19007
D	7	GLN	-	expression tag	UNP Q19007
D	8	HIS	-	expression tag	UNP Q19007
D	9	MET	-	expression tag	UNP Q19007
G	-11	MET	-	initiating methionine	UNP Q19007
G	-10	GLY	-	expression tag	UNP Q19007
G	-9	SER	-	expression tag	UNP Q19007
G	-8	SER	-	expression tag	UNP Q19007
G	-'7	HIS	-	expression tag	UNP Q19007
G	-6	HIS	-	expression tag	UNP Q19007
G	-5	HIS	-	expression tag	UNP Q19007
G	-4	HIS	-	expression tag	UNP Q19007
G	-3	HIS	-	expression tag	UNP Q19007
G	-2	HIS	-	expression tag	UNP Q19007
G	-1	SER	-	expression tag	UNP Q19007
G		SER	-	expression tag	UNP Q19007
G	1	GLY	-	expression tag	UNP Q19007



Chain	Residue	Modelled	Actual	Comment	Reference
G	2	GLU	-	expression tag	UNP Q19007
G	3	ASN	-	expression tag	UNP Q19007
G	4	LEU	-	expression tag	UNP Q19007
G	5	TYR	-	expression tag	UNP Q19007
G	6	PHE	-	expression tag	UNP Q19007
G	7	GLN	-	expression tag	UNP Q19007
G	8	HIS	-	expression tag	UNP Q19007
G	9	MET	-	expression tag	UNP Q19007
Н	-11	MET	-	initiating methionine	UNP Q19007
Н	-10	GLY	-	expression tag	UNP Q19007
Н	-9	SER	-	expression tag	UNP Q19007
Н	-8	SER	-	expression tag	UNP Q19007
Н	-7	HIS	-	expression tag	UNP Q19007
Н	-6	HIS	-	expression tag	UNP Q19007
Н	-5	HIS	-	expression tag	UNP Q19007
Н	-4	HIS	-	expression tag	UNP Q19007
Н	-3	HIS	-	expression tag	UNP Q19007
Н	-2	HIS	-	expression tag	UNP Q19007
Н	-1	SER	-	expression tag	UNP Q19007
Н	0	SER	-	expression tag	UNP Q19007
Н	1	GLY	-	expression tag	UNP Q19007
Н	2	GLU	-	expression tag	UNP Q19007
Н	3	ASN	-	expression tag	UNP Q19007
Н	4	LEU	-	expression tag	UNP Q19007
Н	5	TYR	-	expression tag	UNP Q19007
Н	6	PHE	-	expression tag	UNP Q19007
Н	7	GLN	-	expression tag	UNP Q19007
Н	8	HIS	-	expression tag	UNP Q19007
Н	9	MET	-	expression tag	UNP Q19007

• Molecule 2 is a protein called Histone H2A.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace	
2	2 I	85	Total	С	Ν	Ο	0	0	0
			662	415	131	116			
2	K	80	Total	С	Ν	Ο	0	0	0
	IX	N 02	641	402	127	112		U	U

• Molecule 3 is a protein called Histone H2B 1.1.



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	т	80	Total	С	Ν	0	$\mathbf{S}$	0	0	0
9 1	J	09	694	438	122	132	2	0		
2	т	00	Total	С	Ν	0	S	0	0	0
э L	90	697	441	121	133	2	0		U	



## 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Chain A: 69% 22% 9% • Molecule 1: Nucleosome Assembly Protein Chain B: 66% 21% 13% MET GLY SER HIS SER HIS HIS HIS HIS HIS HIS SER HIS SER ASN ASN CUU CUU CUU CUU LYS LYS LYS GLY GLY GLY ALA ALA ALA ALA ALA • Molecule 1: Nucleosome Assembly Protein Chain E: 59% 22% 19% MET GLY SER HIS HIS HIS HIS HIS HIS
- Molecule 1: Nucleosome Assembly Protein





Chain G:

60%

24%

16%







## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1	Depositor
Cell constants	79.49Å 92.93Å 119.35Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$104.10^{\circ}$ $106.24^{\circ}$ $94.18^{\circ}$	Depositor
Bosolution (Å)	45.93 - 3.35	Depositor
Resolution (A)	45.93 - 3.35	EDS
% Data completeness	78.3(45.93-3.35)	Depositor
(in resolution range)	78.3(45.93-3.35)	EDS
R <sub>merge</sub>	0.13	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	$1.87 (at 3.32 \text{\AA})$	Xtriage
Refinement program	PHENIX (1.19.2_4158: ???)	Depositor
R R.	0.204 , $0.261$	Depositor
$n, n_{free}$	0.203 , $0.260$	DCC
$R_{free}$ test set	2171 reflections $(4.86%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	58.5	Xtriage
Anisotropy	0.360	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.32 , $66.3$	EDS
L-test for $twinning^2$	$ < L >=0.48, < L^2>=0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	19338	wwPDB-VP
Average B, all atoms $(Å^2)$	73.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.85% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

## 5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond	lengths	Bond	angles
IVIOI	Cham	RMSZ	# Z  > 5	RMSZ	# Z  > 5
1	А	0.28	0/2263	0.48	0/3061
1	В	0.27	0/2181	0.48	0/2955
1	С	0.27	0/2221	0.47	0/3006
1	D	0.26	0/2092	0.44	0/2836
1	Е	0.26	0/2005	0.46	0/2723
1	F	0.26	0/2058	0.46	0/2783
1	G	0.26	0/2098	0.47	0/2844
1	Н	0.26	0/2076	0.45	0/2816
2	Ι	0.26	0/670	0.55	0/902
2	Κ	0.25	0/649	0.58	0/874
3	J	0.26	0/705	0.47	0/951
3	L	0.25	0/708	0.48	0/955
All	All	0.26	0/19726	0.47	0/26706

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	2217	0	2163	49	0
1	В	2136	0	2057	47	0
1	С	2177	0	2113	55	0
1	D	2048	0	1978	53	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	Ε	1962	0	1872	52	0
1	F	2018	0	1976	51	0
1	G	2054	0	1987	56	0
1	Н	2032	0	1960	47	0
2	Ι	662	0	697	17	0
2	Κ	641	0	672	33	0
3	J	694	0	709	14	0
3	L	697	0	711	25	0
All	All	19338	0	18895	421	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 421 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)	
2:K:66:LEU:HB3	2:K:87:ALA:HB1	1.55	0.89	
1:G:118:LEU:HD21	1:G:136:VAL:HG13	1.64	0.79	
2:K:25:GLN:HB3	3:L:41:LYS:HE3	1.64	0.77	
1:E:22:LEU:HD23	1:E:24:LEU:H	1.51	0.75	
1:H:11:LEU:HG	1:H:12:LEU:HG	1.69	0.74	

There are no symmetry-related clashes.

### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	276/308~(90%)	265~(96%)	11 (4%)	0	100	100
1	В	263/308~(85%)	251 (95%)	12 (5%)	0	100	100
1	С	269/308~(87%)	251 (93%)	18 (7%)	0	100	100



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	D	254/308~(82%)	243~(96%)	11 (4%)	0	100	100
1	Е	243/308~(79%)	233~(96%)	10 (4%)	0	100	100
1	F	244/308~(79%)	233~(96%)	11 (4%)	0	100	100
1	G	253/308~(82%)	237~(94%)	16 (6%)	0	100	100
1	Н	252/308~(82%)	239~(95%)	13~(5%)	0	100	100
2	Ι	83/92~(90%)	82 (99%)	1 (1%)	0	100	100
2	K	80/92~(87%)	70~(88%)	10 (12%)	0	100	100
3	J	87/99~(88%)	85~(98%)	2(2%)	0	100	100
3	L	88/99~(89%)	86~(98%)	2(2%)	0	100	100
All	All	2392/2846~(84%)	2275 (95%)	117 (5%)	0	100	100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	А	235/265~(89%)	230 (98%)	5 (2%)	48	69
1	В	228/265~(86%)	225~(99%)	3 (1%)	65	79
1	С	234/265~(88%)	232 (99%)	2(1%)	75	87
1	D	216/265~(82%)	214 (99%)	2(1%)	75	87
1	Е	207/265~(78%)	204 (99%)	3 (1%)	62	78
1	F	215/265~(81%)	210 (98%)	5 (2%)	45	68
1	G	219/265~(83%)	218 (100%)	1 (0%)	86	92
1	Н	214/265~(81%)	211 (99%)	3 (1%)	62	78
2	Ι	66/71~(93%)	64 (97%)	2(3%)	36	61
2	K	64/71~(90%)	63~(98%)	1 (2%)	58	75
3	J	76/85~(89%)	75 (99%)	1 (1%)	65	79
3	L	$7\overline{6/85}~(89\%)$	76 (100%)	0	100	100



Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	2050/2432~(84%)	2022~(99%)	28 (1%)	62 78

5 of 28 residues with a non-rotameric side chain are listed below:

Mol	Chain	Res	Type
1	F	251	ASP
2	Κ	26	PHE
2	Ι	95	ASN
1	Н	18	MET
2	Ι	89	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	Е	37	GLN
3	J	47	HIS
1	Н	52	HIS

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

#### 5.6 Ligand geometry (i)

There are no ligands in this entry.

#### 5.7 Other polymers (i)

There are no such residues in this entry.



## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSRZ>2	$\mathbf{OWAB}(\mathrm{\AA}^2)$	Q<0.9
1	А	280/308~(90%)	-0.56	0 100 100	16,  35,  94,  123	0
1	В	269/308~(87%)	-0.53	0 100 100	22,  43,  93,  117	0
1	С	273/308~(88%)	-0.47	0 100 100	24, 54, 120, 138	0
1	D	260/308~(84%)	-0.28	0 100 100	36, 82, 130, 142	0
1	Ε	249/308~(80%)	-0.30	0 100 100	41, 81, 129, 157	0
1	F	254/308~(82%)	-0.33	1 (0%) 89 86	40, 81, 135, 166	0
1	G	259/308~(84%)	-0.29	1 (0%) 89 86	59, 92, 112, 128	0
1	Η	258/308~(83%)	-0.21	0 100 100	67, 95, 119, 147	0
2	Ι	85/92~(92%)	-0.57	0 100 100	32, 56, 87, 100	0
2	Κ	82/92~(89%)	-0.07	2 (2%) 59 50	57, 87, 148, 173	0
3	J	89/99~(89%)	-0.51	0 100 100	37, 58, 82, 113	0
3	L	90/99~(90%)	-0.31	1 (1%) 77 70	50, 83, 133, 149	0
All	All	2448/2846~(86%)	-0.38	5 (0%) 92 92	16, 75, 122, 173	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	L	40	TYR	3.9
2	Κ	80	ILE	3.2
2	Κ	67	ALA	2.8
1	F	10	GLY	2.6
1	G	126	LEU	2.4

### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

### 6.4 Ligands (i)

There are no ligands in this entry.

### 6.5 Other polymers (i)

There are no such residues in this entry.

