



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 10, 2023 – 05:13 AM EDT

PDB ID : 6XC7  
Title : Crystal structure of SARS-CoV-2 receptor binding domain in complex with antibodies CC12.3 and CR3022  
Authors : Yuan, M.; Liu, H.; Wu, N.C.; Zhu, X.; Wilson, I.A.  
Deposited on : 2020-06-08  
Resolution : 2.88 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.88 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2691 (2.90-2.86)
Clashscore	141614	2947 (2.90-2.86)
Ramachandran outliers	138981	2868 (2.90-2.86)
Sidechain outliers	138945	2871 (2.90-2.86)
RSRZ outliers	127900	2629 (2.90-2.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	231	
2	H	222	
3	L	221	
4	C	220	
5	D	215	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
6	B	3	 100%

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 8191 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Spike protein S1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	193	1537	987	256	286	8	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	542	SER	-	expression tag	UNP P0DTC2
A	543	GLY	-	expression tag	UNP P0DTC2
A	544	HIS	-	expression tag	UNP P0DTC2
A	545	HIS	-	expression tag	UNP P0DTC2
A	546	HIS	-	expression tag	UNP P0DTC2
A	547	HIS	-	expression tag	UNP P0DTC2
A	548	HIS	-	expression tag	UNP P0DTC2
A	549	HIS	-	expression tag	UNP P0DTC2

- Molecule 2 is a protein called CR3022 heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	H	218	1619	1027	263	320	9	0	0	0

- Molecule 3 is a protein called CR3022 light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	L	221	1716	1076	284	351	5	0	0	0


- Molecule 4 is a protein called CC12.3 heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	C	220	1645	1040	272	326	7	0	0	0

- Molecule 5 is a protein called CC12.3 light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
5	D	213	1635	1021	279	331	4	0	0	0

- Molecule 6 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
6	B	3	39	22	2	15	0	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Spike protein S1




- Molecule 2: CR3022 heavy chain



- Molecule 3: CR3022 light chain



- Molecule 4: CC12.3 heavy chain





- Molecule 5: CC12.3 light chain



- Molecule 6: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	110.87Å 110.87Å 228.49Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.25 – 2.88 46.17 – 2.88	Depositor EDS
% Data completeness (in resolution range)	99.7 (42.25-2.88) 99.7 (46.17-2.88)	Depositor EDS
$R_{merge}$	0.13	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.40 (at 2.91Å)	Xtrriage
Refinement program	PHENIX 1.16_3549	Depositor
R, $R_{free}$	0.219 , 0.259 0.219 , 0.260	Depositor DCC
$R_{free}$ test set	1654 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	79.5	Xtrriage
Anisotropy	0.084	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.28 , 27.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	8191	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	76.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.10% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.27	0/1580	0.47	0/2149
2	H	0.26	0/1660	0.49	0/2259
3	L	0.27	0/1754	0.50	0/2383
4	C	0.30	0/1685	0.56	1/2292 (0.0%)
5	D	0.29	0/1670	0.56	0/2266
All	All	0.28	0/8349	0.52	1/11349 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
2	H	0	1
4	C	0	2
5	D	0	1
All	All	0	5

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	131	THR	N-CA-C	-6.09	94.56	111.00

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	449	TYR	Peptide
4	C	127	SER	Peptide
4	C	129	LYS	Peptide
5	D	30	SER	Peptide
2	H	135	THR	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1537	0	1458	19	0
2	H	1619	0	1591	11	0
3	L	1716	0	1658	20	0
4	C	1645	0	1604	30	0
5	D	1635	0	1592	38	0
6	B	39	0	34	0	0
All	All	8191	0	7937	109	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (109) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:D:77:ARG:O	5:D:79:GLU:OE1	1.86	0.94
1:A:383:SER:HB2	2:H:96:SER:HB3	1.58	0.85
4:C:188:SER:HG	4:C:194:TYR:HH	1.22	0.84
3:L:3:GLN:H	3:L:26:SER:HB3	1.50	0.76
5:D:80:PRO:HA	5:D:83:PHE:HE2	1.51	0.75
5:D:78:LEU:C	5:D:79:GLU:OE2	2.28	0.73
5:D:143:GLU:C	5:D:143:GLU:OE2	2.28	0.71
4:C:66:ARG:NH2	4:C:86:ASP:OD2	2.24	0.70
4:C:187:SER:HA	4:C:189:LEU:HD22	1.72	0.69
4:C:186:SER:O	4:C:189:LEU:HB3	1.92	0.69
5:D:77:ARG:O	5:D:79:GLU:CD	2.30	0.69
2:H:201:LYS:HG2	2:H:202:PRO:HD3	1.78	0.66
1:A:455:LEU:HD12	1:A:493:GLN:HG3	1.79	0.65

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:D:122:ASP:HA	5:D:125:LEU:HB2	1.79	0.65
1:A:381:GLY:HA3	1:A:430:THR:HG22	1.80	0.63
5:D:78:LEU:O	5:D:79:GLU:OE2	2.17	0.63
5:D:113:PRO:HD2	5:D:201:LEU:HD11	1.80	0.63
3:L:54:ARG:HG2	3:L:58:VAL:HB	1.82	0.62
4:C:11:LEU:HB2	4:C:147:PRO:HG3	1.81	0.62
5:D:185:ASP:HA	5:D:188:LYS:HD2	1.82	0.62
4:C:152:VAL:HG22	4:C:198:VAL:HG13	1.82	0.61
4:C:29:VAL:HG21	4:C:76:SER:HA	1.83	0.60
4:C:119:PRO:HB3	4:C:145:TYR:HB3	1.83	0.60
5:D:182:SER:HG	5:D:185:ASP:H	1.51	0.59
4:C:163:VAL:HG22	4:C:182:VAL:HG12	1.85	0.59
4:C:184:VAL:HG11	4:C:194:TYR:OH	2.03	0.58
4:C:189:LEU:HD23	4:C:190:GLY:H	1.69	0.58
5:D:106:ILE:H	5:D:166:GLN:HE22	1.52	0.58
1:A:393:THR:O	1:A:394:ASN:ND2	2.38	0.57
5:D:8:PRO:HD2	5:D:11:LEU:HD11	1.87	0.57
3:L:80:ALA:HA	3:L:106:ILE:HD11	1.87	0.56
4:C:82:MET:HB3	4:C:82(C):LEU:HD21	1.88	0.56
5:D:3:VAL:H	5:D:26:SER:HB3	1.71	0.55
5:D:77:ARG:O	5:D:79:GLU:OE2	2.25	0.54
3:L:25:SER:OG	3:L:27:GLN:O	2.26	0.54
1:A:489:TYR:OH	4:C:94:ARG:NH2	2.32	0.54
4:C:195:ILE:HG22	4:C:210:LYS:HA	1.92	0.52
5:D:38:GLN:O	5:D:84:ALA:HB1	2.10	0.52
3:L:13:VAL:HG21	3:L:19:ALA:HB2	1.90	0.52
5:D:198:HIS:HB3	5:D:201:LEU:CD1	2.40	0.52
4:C:123:PRO:HB3	4:C:211:VAL:HG12	1.92	0.51
1:A:501:ASN:HB3	1:A:505:TYR:HB2	1.93	0.50
3:L:76:SER:OG	3:L:77:SER:N	2.44	0.50
5:D:29:VAL:HG11	5:D:90:GLN:HB2	1.93	0.50
5:D:85:VAL:HG22	5:D:103:LYS:HG3	1.94	0.50
1:A:403:ARG:NH2	1:A:405:ASP:OD2	2.45	0.49
5:D:21:LEU:HD13	5:D:73:LEU:HD23	1.95	0.49
4:C:188:SER:HA	4:C:191:THR:OG1	2.13	0.48
4:C:195:ILE:HD12	4:C:208:ASP:HB3	1.95	0.48
2:H:97:GLY:O	2:H:100:THR:OG1	2.32	0.48
4:C:187:SER:C	4:C:189:LEU:H	2.16	0.48
1:A:354:ASN:O	1:A:398:ASP:HA	2.14	0.48
3:L:27(D):TYR:HB3	3:L:28:ILE:HG22	1.94	0.48
4:C:189:LEU:HD23	4:C:190:GLY:N	2.28	0.48

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:D:198:HIS:HB3	5:D:201:LEU:HD12	1.95	0.47
1:A:425:LEU:HD21	1:A:512:VAL:HG11	1.96	0.47
3:L:13:VAL:HG12	3:L:14:SER:O	2.15	0.47
5:D:113:PRO:HB3	5:D:139:PHE:CD1	2.50	0.47
1:A:449:TYR:HD2	1:A:450:ASN:HA	1.80	0.47
5:D:80:PRO:HA	5:D:83:PHE:CE2	2.40	0.47
2:H:136:ALA:HB2	2:H:186:SER:HB3	1.97	0.47
5:D:147:GLN:HG2	5:D:154:LEU:HD13	1.96	0.46
3:L:31:ASN:O	3:L:50:TRP:HA	2.15	0.46
3:L:183:LYS:O	3:L:187:GLU:HG3	2.16	0.46
1:A:428:ASP:O	3:L:27(D):TYR:OH	2.29	0.46
4:C:211:VAL:HG23	4:C:211:VAL:O	2.16	0.46
4:C:138:LEU:HD13	4:C:211:VAL:HG21	1.97	0.46
2:H:100(A):PRO:HD3	3:L:91:TYR:CZ	2.51	0.45
5:D:167:ASP:O	5:D:171:SER:HA	2.16	0.45
1:A:377:PHE:CG	2:H:31:THR:HG22	2.52	0.45
5:D:120:PRO:HD3	5:D:132:VAL:HG22	1.99	0.45
3:L:184:ALA:O	3:L:188:LYS:HG3	2.16	0.45
2:H:136:ALA:HB3	2:H:184:VAL:O	2.16	0.45
5:D:37:GLN:HB2	5:D:47:LEU:HD11	1.99	0.45
4:C:69:ILE:HD11	4:C:78:LEU:HD11	1.98	0.44
4:C:130:SER:O	4:C:131:THR:HG23	2.18	0.44
5:D:115:VAL:HG12	5:D:207:LYS:HG3	2.01	0.43
5:D:167:ASP:CG	5:D:170:ASP:OD2	2.57	0.43
4:C:101:ASP:N	4:C:101:ASP:OD1	2.47	0.43
4:C:33:TYR:CE1	4:C:97:GLY:HA2	2.55	0.42
1:A:401:VAL:HG22	1:A:509:ARG:HG2	2.02	0.42
5:D:106:ILE:HG22	5:D:107:LYS:N	2.34	0.42
5:D:138:ASN:HA	5:D:172:THR:HB	2.01	0.42
2:H:122:PHE:CE2	3:L:124:GLN:HG3	2.54	0.42
4:C:127:SER:HG	4:C:130:SER:HA	1.85	0.41
2:H:181:VAL:HG11	3:L:135:LEU:HD13	2.01	0.41
4:C:163:VAL:HG22	4:C:182:VAL:CG1	2.50	0.41
4:C:127:SER:OG	4:C:130:SER:HA	2.21	0.41
1:A:475:ALA:HB1	4:C:32:ASN:HD21	1.86	0.41
3:L:122:ASP:HA	3:L:125:LEU:HB2	2.03	0.41
5:D:140:TYR:CG	5:D:141:PRO:HA	2.56	0.41
5:D:143:GLU:OE2	5:D:144:ALA:N	2.54	0.41
5:D:81:GLU:C	5:D:83:PHE:H	2.22	0.41
1:A:366:SER:O	1:A:370:ASN:HB2	2.21	0.41
1:A:398:ASP:O	1:A:511:VAL:HA	2.20	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:528:LYS:HE2	1:A:528:LYS:HB3	1.87	0.41
2:H:99:SER:N	3:L:96:TYR:OH	2.45	0.41
2:H:119:PRO:HB3	2:H:145:TYR:HB3	2.02	0.41
3:L:124:GLN:O	3:L:127:SER:OG	2.32	0.41
5:D:19:ALA:O	5:D:74:THR:HA	2.21	0.41
3:L:33:LEU:HD22	3:L:89:GLN:O	2.21	0.40
4:C:195:ILE:O	4:C:195:ILE:HG13	2.19	0.40
1:A:380:TYR:O	1:A:430:THR:HA	2.21	0.40
5:D:107:LYS:HA	5:D:140:TYR:OH	2.21	0.40
3:L:140:TYR:CG	3:L:141:PRO:HA	2.57	0.40
5:D:24:ARG:HA	5:D:69:THR:O	2.21	0.40
1:A:392:PHE:HB3	1:A:516:GLU:O	2.22	0.40
5:D:18:ARG:HB2	5:D:75:ILE:O	2.22	0.40
5:D:82:ASP:O	5:D:86:TYR:OH	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	189/231 (82%)	177 (94%)	12 (6%)	0	100	100
2	H	214/222 (96%)	208 (97%)	6 (3%)	0	100	100
3	L	219/221 (99%)	211 (96%)	8 (4%)	0	100	100
4	C	218/220 (99%)	212 (97%)	5 (2%)	1 (0%)	29	59
5	D	211/215 (98%)	201 (95%)	9 (4%)	1 (0%)	29	59
All	All	1051/1109 (95%)	1009 (96%)	40 (4%)	2 (0%)	47	76

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	C	130	SER
5	D	31	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	168/203 (83%)	161 (96%)	7 (4%)	30	61
2	H	183/187 (98%)	175 (96%)	8 (4%)	28	59
3	L	196/196 (100%)	191 (97%)	5 (3%)	46	76
4	C	186/186 (100%)	181 (97%)	5 (3%)	44	75
5	D	184/186 (99%)	176 (96%)	8 (4%)	29	60
All	All	917/958 (96%)	884 (96%)	33 (4%)	35	67

All (33) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	354	ASN
1	A	357	ARG
1	A	377	PHE
1	A	386	LYS
1	A	394	ASN
1	A	449	TYR
1	A	469	SER
2	H	22	CYS
2	H	43	LYS
2	H	73	LYS
2	H	96	SER
2	H	127	SER
2	H	138	LEU
2	H	210	LYS
2	H	216	CYS
3	L	26	SER
3	L	27(D)	TYR
3	L	60	ASP

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
3	L	203	SER
3	L	207	LYS
4	C	75	LYS
4	C	92	CYS
4	C	132	SER
4	C	197	ASN
4	C	214	LYS
5	D	18	ARG
5	D	22	SER
5	D	60	ASP
5	D	63	SER
5	D	65	SER
5	D	67	SER
5	D	79	GLU
5	D	183	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	394	ASN
5	D	166	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

3 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
6	NAG	B	1	6,1	14,14,15	0.37	0	17,19,21	0.43	0
6	NAG	B	2	6	14,14,15	0.20	0	17,19,21	0.46	0
6	BMA	B	3	6	11,11,12	0.64	0	15,15,17	0.76	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	NAG	B	1	6,1	-	2/6/23/26	0/1/1/1
6	NAG	B	2	6	-	2/6/23/26	0/1/1/1
6	BMA	B	3	6	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
6	B	2	NAG	O5-C5-C6-O6
6	B	2	NAG	C4-C5-C6-O6
6	B	1	NAG	C4-C5-C6-O6
6	B	1	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	193/231 (83%)	0.13	1 (0%) 91 90	50, 70, 113, 141	0
2	H	218/222 (98%)	0.01	2 (0%) 84 84	53, 65, 104, 193	0
3	L	221/221 (100%)	-0.08	3 (1%) 75 75	53, 67, 87, 152	0
4	C	220/220 (100%)	0.17	10 (4%) 33 29	51, 72, 139, 164	0
5	D	213/215 (99%)	0.20	8 (3%) 40 36	61, 83, 115, 146	0
All	All	1065/1109 (96%)	0.08	24 (2%) 60 59	50, 71, 116, 193	0

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	L	215	SER	8.2
2	H	215	SER	5.9
4	C	194	TYR	5.5
4	C	159	LEU	4.4
4	C	131	THR	4.4
2	H	216	CYS	4.2
4	C	158	ALA	3.4
4	C	215	SER	3.0
5	D	169	LYS	2.9
3	L	214	CYS	2.8
4	C	216	CYS	2.7
1	A	519	HIS	2.5
4	C	154	TRP	2.4
5	D	13	LEU	2.4
4	C	132	SER	2.4
5	D	12	SER	2.3
4	C	130	SER	2.3
5	D	82	ASP	2.3
5	D	83	PHE	2.2
4	C	195	ILE	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
5	D	105	GLU	2.2
5	D	135	LEU	2.1
5	D	107	LYS	2.1
3	L	104	VAL	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
6	NAG	B	1	14/15	0.93	0.16	69,75,79,83	0
6	BMA	B	3	11/12	0.93	0.11	104,110,114,114	0
6	NAG	B	2	14/15	0.94	0.12	85,94,98,103	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.