



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 2, 2024 – 02:09 PM JST

PDB ID : 8XC9
Title : Amylase A in complex with acarbose
Authors : Zhao, F.; Xu, T.T.
Deposited on : 2023-12-08
Resolution : 1.89 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.21
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.004 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.40


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.89 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	7293 (1.90-1.90)
Clashscore	180529	8090 (1.90-1.90)
Ramachandran outliers	177936	8022 (1.90-1.90)
Sidechain outliers	177891	8022 (1.90-1.90)
RSRZ outliers	164620	7292 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	498	 % 88% 5% 8%
1	B	498	 5% 84% 8% 8%
2	E	2	 100%
2	F	2	 50% 50%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 8054 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called AmyA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	460	3718	2381	632	692	13	0	0	0
1	B	460	3718	2381	632	692	13	0	0	0

- Molecule 2 is an oligosaccharide called 4,6-dideoxy-4-[[(1S,4R,5S,6S)]-4,5,6-trihydroxy-3-(hydroxymethyl)cyclohex-2-en-1-yl]amino}-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	E	2	33	19	1	13	0	0	0
2	F	2	33	19	1	13	0	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	328	Total	O	0	0
			328	328		
3	B	224	Total	O	0	0
			224	224		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: AmyA



- Molecule 1: AmyA



- Molecule 2: 4,6-dideoxy-4- $\{[(1S,4R,5S,6S)-4,5,6\text{-trihydroxy-3-(hydroxymethyl)cyclohex-2-en-1-yl]amino}\}$ - α -D-glucopyranose-(1-4)- α -D-glucopyranose



- Molecule 2: 4,6-dideoxy-4- $\{[(1S,4R,5S,6S)-4,5,6\text{-trihydroxy-3-(hydroxymethyl)cyclohex-2-en-1-yl]amino}\}$ - α -D-glucopyranose-(1-4)- α -D-glucopyranose



GLC1
ACT2

4 Data and refinement statistics i

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	140.00Å 140.00Å 84.18Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	36.43 – 1.89 36.43 – 1.89	Depositor EDS
% Data completeness (in resolution range)	99.5 (36.43-1.89) 92.1 (36.43-1.89)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.74 (at 1.89Å)	Xtrriage
Refinement program	PHENIX (1.17.1_3660: ???)	Depositor
R, R_{free}	0.221 , 0.266 0.218 , 0.266	Depositor DCC
R_{free} test set	73743 reflections (2.65%)	wwPDB-VP
Wilson B-factor (Å ²)	21.2	Xtrriage
Anisotropy	0.937	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 37.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.039 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8054	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.96% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: AC1, GLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.42	0/3847	0.56	0/5257
1	B	0.38	0/3847	0.54	0/5257
All	All	0.40	0/7694	0.55	0/10514

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3718	0	3405	11	0
1	B	3718	0	3405	23	0
2	E	33	0	21	0	0
2	F	33	0	21	2	0
3	A	328	0	0	0	0
3	B	224	0	0	0	0
All	All	8054	0	6852	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:462:VAL:HG11	1:B:477:PRO:HG2	1.80	0.62
1:B:52:TRP:HB2	1:B:343:ARG:HD2	1.88	0.55
1:B:52:TRP:CB	1:B:343:ARG:HD2	2.37	0.55
1:B:394:ALA:HA	1:B:398:PHE:HD2	1.72	0.54
1:A:477:PRO:HD2	1:A:489:TRP:CZ2	2.42	0.54
1:B:329:ARG:HD3	1:B:369:HIS:CG	2.44	0.52
1:A:357:LYS:NZ	1:A:378:ASP:OD1	2.37	0.50
1:A:153:ILE:HB	1:A:156:PHE:HB3	1.94	0.50
1:A:329:ARG:HD3	1:A:369:HIS:CG	2.47	0.49
1:B:249:ALA:HB1	1:B:252:HIS:HD2	1.77	0.49
1:B:367:ARG:HA	1:B:416:TYR:CE2	2.48	0.48
1:A:51:TYR:CD2	1:A:377:ARG:HG3	2.48	0.48
1:B:291:VAL:HG12	1:B:293:THR:HG22	1.96	0.48
1:B:70:ALA:HB2	1:B:133:ILE:HD11	1.95	0.47
1:A:418:ARG:HB2	1:A:429:LEU:HG	1.96	0.47
1:B:184:PHE:CE1	1:B:224:GLU:HB3	2.50	0.47
1:A:481:THR:O	1:A:485:GLY:HA3	2.16	0.46
1:A:266:GLN:HG2	1:A:295:HIS:CE1	2.49	0.46
1:B:280:TRP:CD2	2:F:1:GLC:H5	2.50	0.46
1:B:118:ARG:O	1:B:122:GLN:HG2	2.16	0.46
1:A:181:TRP:CH2	1:A:183:ASP:HB2	2.50	0.46
1:A:97:TYR:CD2	1:A:98:ASP:HB2	2.51	0.45
1:B:77:MET:HG3	1:B:133:ILE:HG21	1.99	0.44
1:A:256:ASP:OD1	1:A:259:ARG:NH2	2.49	0.43
1:B:134:GLN:HB3	1:B:243:SER:HB2	1.99	0.43
1:B:124:ALA:O	1:B:128:LEU:HG	2.19	0.43
1:B:204:TRP:HB3	1:B:211:PRO:CB	2.49	0.42
1:B:339:HIS:O	1:B:343:ARG:HD3	2.20	0.42
1:B:191:PHE:O	1:B:202:LYS:HE3	2.21	0.41
1:B:230:LYS:HB3	1:B:264:HIS:CD2	2.55	0.41
1:B:48:GLN:O	1:B:376:ALA:HB2	2.20	0.41
1:B:105:PHE:CE1	1:B:167:GLY:HA2	2.56	0.40
1:B:220:TYR:HE1	1:B:229:MET:HG3	1.85	0.40
1:B:280:TRP:CE3	2:F:1:GLC:H5	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	458/498 (92%)	444 (97%)	14 (3%)	0	100	100
1	B	458/498 (92%)	439 (96%)	19 (4%)	0	100	100
All	All	916/996 (92%)	883 (96%)	33 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	379/408 (93%)	378 (100%)	1 (0%)	91	92
1	B	379/408 (93%)	375 (99%)	4 (1%)	70	71
All	All	758/816 (93%)	753 (99%)	5 (1%)	81	83

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	209	HIS
1	B	145	MET
1	B	343	ARG
1	B	434	SER
1	B	471	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GLC	E	1	2	12,12,12	1.03	1 (8%)	17,17,17	1.81	3 (17%)
2	AC1	E	2	2	21,22,23	0.72	1 (4%)	22,32,34	0.74	1 (4%)
2	GLC	F	1	2	12,12,12	1.19	1 (8%)	17,17,17	1.92	6 (35%)
2	AC1	F	2	2	21,22,23	0.68	1 (4%)	22,32,34	0.52	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	E	1	2	-	0/2/22/22	0/1/1/1
2	AC1	E	2	2	-	3/6/43/46	0/2/2/2
2	GLC	F	1	2	-	0/2/22/22	0/1/1/1
2	AC1	F	2	2	-	2/6/43/46	0/2/2/2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	1	GLC	O4-C4	-3.70	1.34	1.43

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	1	GLC	O4-C4	-2.85	1.36	1.43
2	E	2	AC1	C1B-N4A	2.29	1.51	1.47
2	F	2	AC1	C1B-N4A	2.20	1.51	1.47

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	1	GLC	O5-C1-C2	4.81	118.87	110.28
2	F	1	GLC	C1-O5-C5	4.27	121.71	113.66
2	E	1	GLC	C1-O5-C5	4.11	121.42	113.66
2	F	1	GLC	O5-C1-C2	4.01	117.44	110.28
2	F	1	GLC	O4-C4-C5	-2.72	102.55	109.30
2	F	1	GLC	O5-C5-C4	2.41	114.07	109.69
2	E	1	GLC	C1-C2-C3	2.26	115.00	110.31
2	F	1	GLC	C1-C2-C3	2.20	114.89	110.31
2	E	2	AC1	C2-C3-C4	2.19	112.56	110.63
2	F	1	GLC	O4-C4-C3	-2.05	105.61	110.35

There are no chirality outliers.

All (5) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
2	E	2	AC1	C7B-C1B-N4A-C4
2	E	2	AC1	C7B-C5B-C6B-O6B
2	F	2	AC1	C7B-C1B-N4A-C4
2	F	2	AC1	C7B-C5B-C6B-O6B
2	E	2	AC1	C4A-C5B-C6B-O6B

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	F	1	GLC	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	460/498 (92%)	0.30	4 (0%) 81 82	14, 22, 33, 41	0
1	B	460/498 (92%)	0.73	27 (5%) 29 30	17, 29, 46, 63	0
All	All	920/996 (92%)	0.51	31 (3%) 48 50	14, 25, 42, 63	0

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	35	PRO	3.4
1	B	36	GLN	3.4
1	B	64	ALA	3.0
1	B	67	ALA	2.8
1	A	203	TRP	2.7
1	A	271	GLU	2.7
1	B	491	PRO	2.6
1	B	124	ALA	2.6
1	B	466	VAL	2.6
1	B	493	SER	2.5
1	B	62	LEU	2.4
1	B	39	PRO	2.4
1	B	462	VAL	2.4
1	B	127	ALA	2.3
1	B	430	VAL	2.3
1	B	490	VAL	2.3
1	B	458	TYR	2.3
1	B	472	GLY	2.2
1	B	117	THR	2.2
1	B	34	LEU	2.2
1	A	422	SER	2.1
1	B	398	PHE	2.1
1	B	125	LEU	2.1
1	B	470	ALA	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	489	TRP	2.1
1	B	483	SER	2.1
1	B	54	ALA	2.1
1	B	422	SER	2.1
1	A	37	ILE	2.0
1	B	456	TYR	2.0
1	B	59	TRP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

SUGAR-RSR INFOmissingINFO

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.