



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 15, 2023 – 06:14 PM EDT

PDB ID : 1XEY  
Title : Crystal structure of the complex of Escherichia coli GADA with glutarate at 2.05 Å resolution  
Authors : Dutyshev, D.I.; Darii, E.L.; Fomenkova, N.P.; Pechik, I.V.; Polyakov, K.M.; Nikonov, S.V.; Andreeva, N.S.; Sukhareva, B.S.  
Deposited on : 2004-09-13  
Resolution : 2.05 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1692 (2.04-2.04)
Clashscore	141614	1773 (2.04-2.04)
Ramachandran outliers	138981	1752 (2.04-2.04)
Sidechain outliers	138945	1752 (2.04-2.04)
RSRZ outliers	127900	1672 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	466	<div style="display: flex; align-items: center;"> <div style="width: 2%; height: 10px; background-color: red; margin-right: 2px;"></div> <div style="width: 84%; height: 10px; background-color: green; margin-right: 2px;"></div> <div style="width: 11%; height: 10px; background-color: yellow; margin-right: 2px;"></div> <div style="width: 3%; height: 10px; background-color: grey;"></div> </div> <p style="margin-left: 20px;">84% 11% . .</p>
1	B	466	<div style="display: flex; align-items: center;"> <div style="width: 2%; height: 10px; background-color: red; margin-right: 2px;"></div> <div style="width: 86%; height: 10px; background-color: green; margin-right: 2px;"></div> <div style="width: 9%; height: 10px; background-color: yellow; margin-right: 2px;"></div> <div style="width: 3%; height: 10px; background-color: grey;"></div> </div> <p style="margin-left: 20px;">86% 9% . .</p>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	ACT	A	600	-	X	X	-
4	GUA	A	506[B]	-	X	-	-

## 2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 7612 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glutamate decarboxylase alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	448	Total 3568	C 2274	N 606	O 662	S 26	0	7	0
1	B	449	Total 3586	C 2289	N 611	O 660	S 26	0	7	0

- Molecule 2 is ACETATE ION (three-letter code: ACT) (formula: C<sub>2</sub>H<sub>3</sub>O<sub>2</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	Total 4	C 2	O 2	0	0
2	A	1	Total 4	C 2	O 2	0	0

- Molecule 3 is PYRIDOXAL-5'-PHOSPHATE (three-letter code: PLP) (formula: C<sub>8</sub>H<sub>10</sub>NO<sub>6</sub>P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	N	O	P			
3	A	1	Total	15	8	1	5	1	0	0
3	B	1	Total	15	8	1	5	1	0	0

- Molecule 4 is GLUTARIC ACID (three-letter code: GUA) (formula:  $C_5H_8O_4$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
			Total	C	O			
4	A	1	Total	15	9	6	0	1
4	B	1	Total	15	9	6	0	1


- Molecule 5 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
5	A	209	Total 209	O 209	0	0
5	B	181	Total 181	O 181	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Glutamate decarboxylase alpha



- Molecule 1: Glutamate decarboxylase alpha



## 4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	117.10Å 117.10Å 196.40Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.00 – 2.05 30.04 – 2.05	Depositor EDS
% Data completeness (in resolution range)	89.3 (30.00-2.05) 89.3 (30.04-2.05)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.09	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.98 (at 2.05Å)	Xtrriage
Refinement program	REFMAC 5.1.24	Depositor
R, $R_{free}$	0.148 , 0.199 0.148 , 0.199	Depositor DCC
$R_{free}$ test set	2857 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	21.9	Xtrriage
Anisotropy	0.410	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 48.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.039 for h,-h-k,-l	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	7612	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	23.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.84% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACT, GUA, PLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.80	0/3694	0.85	13/5008 (0.3%)
1	B	0.80	0/3715	0.82	9/5035 (0.2%)
All	All	0.80	0/7409	0.84	22/10043 (0.2%)

There are no bond length outliers.

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	440	ASP	CB-CG-OD2	6.72	124.35	118.30
1	B	39	ASP	CB-CG-OD2	6.34	124.00	118.30
1	B	106	ASP	CB-CG-OD2	6.20	123.88	118.30
1	B	243	ASP	CB-CG-OD2	5.96	123.67	118.30
1	A	243	ASP	CB-CG-OD2	5.86	123.57	118.30
1	A	39	ASP	CB-CG-OD2	5.75	123.48	118.30
1	A	432	ASP	CB-CG-OD2	5.69	123.42	118.30
1	A	34	LEU	CA-CB-CG	5.65	128.30	115.30
1	A	40	ASP	CB-CG-OD2	5.61	123.35	118.30
1	B	97	ASP	CB-CG-OD1	5.59	123.33	118.30
1	A	417	ASP	CB-CG-OD2	5.54	123.29	118.30
1	A	450	ASP	CB-CG-OD2	5.54	123.29	118.30
1	B	40	ASP	CB-CG-OD2	5.44	123.20	118.30
1	B	372	ASP	CB-CG-OD2	5.35	123.11	118.30
1	B	440	ASP	CB-CG-OD2	5.35	123.11	118.30
1	A	291	ASP	CB-CG-OD2	5.27	123.05	118.30
1	B	190	ASP	CB-CG-OD2	5.20	122.98	118.30
1	A	189	MET	CG-SD-CE	5.15	108.43	100.20
1	A	372	ASP	CB-CG-OD2	5.14	122.93	118.30
1	B	355	ASP	CB-CG-OD1	5.13	122.92	118.30
1	A	97	ASP	CB-CG-OD1	5.06	122.86	118.30
1	A	199	ASP	CB-CG-OD2	5.04	122.83	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3568	0	3454	23	0
1	B	3586	0	3477	15	0
2	A	8	0	6	4	0
3	A	15	0	6	2	0
3	B	15	0	6	2	0
4	A	15	0	0	0	0
4	B	15	0	0	0	0
5	A	209	0	0	1	0
5	B	181	0	0	0	0
All	All	7612	0	6949	36	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:334:LEU:HD11	2:A:600:ACT:H2	1.63	0.78
1:A:342:VAL:HG21	2:A:600:ACT:H1	1.73	0.70
1:A:334:LEU:HD11	2:A:600:ACT:CH3	2.23	0.68
1:B:229:LYS:NZ	1:B:233:ASP:OD2	2.31	0.63
1:A:103:MET:HE3	1:B:33:PRO:HD3	1.83	0.60
1:B:37:MET:CE	1:B:41:VAL:HG12	2.31	0.60
1:A:127:SER:O	1:A:131[A]:MET:HG2	2.03	0.58
1:A:126:SER:HB2	3:A:500:PLP:O4P	2.05	0.57
1:A:352:TYR:OH	1:A:435[A]:GLU:OE1	2.23	0.56
1:B:131[B]:MET:HG2	1:B:173:TRP:CZ3	2.40	0.56
1:A:310:ILE:HD13	1:A:311:GLY:O	2.06	0.55
1:A:395:LEU:HD21	1:A:441:TYR:CZ	2.42	0.54
1:A:16:SER:HB2	2:A:700:ACT:CH3	2.40	0.51
1:A:339:TYR:O	1:A:343:GLN:HG2	2.10	0.50

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:370:ARG:HG2	1:A:370:ARG:HH11	1.76	0.50
1:B:126:SER:HB2	3:B:500:PLP:O4P	2.12	0.49
1:B:395:LEU:HD21	1:B:441:TYR:CZ	2.49	0.48
1:A:245:ALA:HA	1:A:272:ALA:HA	1.96	0.47
1:A:315:ILE:HG23	1:B:131[B]:MET:SD	2.53	0.47
1:A:435[B]:GLU:OE2	1:A:439:GLU:OE2	2.32	0.47
1:B:339:TYR:O	1:B:343:GLN:HG2	2.15	0.47
1:B:127:SER:O	1:B:131[A]:MET:HG2	2.14	0.47
1:B:129:ALA:HB1	1:B:287:VAL:HB	1.97	0.45
1:A:446:LYS:HE3	1:A:446:LYS:HB3	1.47	0.45
1:A:275:HIS:HA	1:A:280:ALA:O	2.17	0.44
1:A:321:ALA:O	1:A:324[B]:VAL:HG12	2.18	0.43
1:B:276:LYS:NZ	3:B:500:PLP:O3	2.51	0.43
1:B:284:CYS:HB2	1:B:323:GLN:HB3	2.01	0.43
1:A:144:ARG:NH1	1:A:237:ASP:O	2.53	0.42
1:A:114:LYS:H	1:A:114:LYS:HG2	1.63	0.41
1:A:195:ILE:HD13	1:A:195:ILE:HA	1.90	0.41
1:A:284:CYS:HB2	1:A:323:GLN:HB3	2.03	0.40
1:B:59:ASN:HA	1:B:405:GLN:HB2	2.02	0.40
1:B:157[B]:LEU:HD21	1:B:206:VAL:HG23	2.02	0.40
1:A:276:LYS:NZ	3:A:500:PLP:O3	2.55	0.40
5:A:1072:HOH:O	1:B:400:ARG:NH2	2.53	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	453/466 (97%)	439 (97%)	14 (3%)	0	100	100
1	B	454/466 (97%)	441 (97%)	13 (3%)	0	100	100
All	All	907/932 (97%)	880 (97%)	27 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	376/389 (97%)	360 (96%)	16 (4%)	29	22
1	B	378/389 (97%)	359 (95%)	19 (5%)	24	16
All	All	754/778 (97%)	719 (95%)	35 (5%)	27	19

All (35) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	30	LYS
1	A	114	LYS
1	A	143	LYS
1	A	144	ARG
1	A	183	ARG
1	A	186	GLN
1	A	229	LYS
1	A	290	ARG
1	A	297	GLN
1	A	310	ILE
1	A	333	ARG
1	A	355	ASP
1	A	433	PHE
1	A	436	LEU
1	A	446	LYS
1	A	449	SER
1	B	8	ASP
1	B	21	LYS
1	B	37	MET
1	B	78	LEU
1	B	157[A]	LEU
1	B	157[B]	LEU
1	B	290	ARG
1	B	293	GLU
1	B	333	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	348[A]	GLN
1	B	348[B]	GLN
1	B	355	ASP
1	B	381	LYS
1	B	383	LYS
1	B	417	ASP
1	B	433	PHE
1	B	436	LEU
1	B	446	LYS
1	B	449	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	81	ASN
1	A	344	ASN
1	B	81	ASN
1	B	344	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	PLP	A	500	1	15,15,16	1.71	3 (20%)	20,22,23	1.70	5 (25%)
4	GUA	B	505[A]	-	8,8,8	1.43	1 (12%)	9,9,9	1.51	2 (22%)
4	GUA	A	506[A]	-	8,8,8	1.49	2 (25%)	9,9,9	2.52	2 (22%)
3	PLP	B	500	1	15,15,16	1.91	3 (20%)	20,22,23	2.10	7 (35%)
2	ACT	A	700	-	3,3,3	1.18	0	3,3,3	2.96	2 (66%)
4	GUA	A	506[B]	-	8,8,8	1.37	2 (25%)	9,9,9	1.43	3 (33%)
4	GUA	B	505[B]	-	8,8,8	1.34	1 (12%)	9,9,9	1.28	0
2	ACT	A	600	-	3,3,3	1.42	1 (33%)	3,3,3	2.38	2 (66%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PLP	A	500	1	-	0/6/6/8	0/1/1/1
4	GUA	B	505[A]	-	-	3/6/6/6	-
4	GUA	A	506[A]	-	-	4/6/6/6	-
3	PLP	B	500	1	-	2/6/6/8	0/1/1/1
4	GUA	A	506[B]	-	-	5/6/6/6	-
4	GUA	B	505[B]	-	-	5/6/6/6	-

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	500	PLP	O3-C3	-4.90	1.25	1.37
3	B	500	PLP	O3-C3	-4.76	1.25	1.37
3	B	500	PLP	C2-N1	3.10	1.39	1.33
3	B	500	PLP	C6-N1	2.81	1.40	1.34
4	A	506[A]	GUA	O2-C1	-2.24	1.23	1.30
3	A	500	PLP	C6-N1	2.21	1.39	1.34
4	A	506[B]	GUA	O2-C1	-2.21	1.23	1.30
4	A	506[A]	GUA	O4-C5	-2.20	1.23	1.30
4	A	506[B]	GUA	O4-C5	-2.20	1.23	1.30
3	A	500	PLP	C2-N1	2.20	1.38	1.33
4	B	505[A]	GUA	O4-C5	-2.17	1.23	1.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	505[B]	GUA	O4-C5	-2.17	1.23	1.30
2	A	600	ACT	OXT-C	-2.07	1.20	1.30

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	506[A]	GUA	O2-C1-O1	-5.96	108.44	123.30
3	B	500	PLP	C4A-C4-C5	-4.69	116.11	120.94
2	A	700	ACT	OXT-C-CH3	3.78	130.82	115.18
4	A	506[A]	GUA	O2-C1-C2	3.64	125.71	114.03
3	A	500	PLP	C5-C6-N1	-3.40	118.15	123.82
2	A	700	ACT	OXT-C-O	-3.40	109.53	122.05
3	B	500	PLP	O4P-P-O1P	-3.37	97.01	106.47
2	A	600	ACT	OXT-C-O	-3.22	110.19	122.05
3	B	500	PLP	C5-C6-N1	-3.20	118.50	123.82
3	B	500	PLP	O3P-P-O2P	3.11	119.51	107.64
3	A	500	PLP	O4P-P-O1P	-3.05	97.92	106.47
4	B	505[A]	GUA	O2-C1-O1	-2.87	116.15	123.30
3	A	500	PLP	C6-C5-C4	2.83	120.39	118.16
3	A	500	PLP	O4P-C5A-C5	2.73	114.56	109.35
2	A	600	ACT	OXT-C-CH3	2.54	125.67	115.18
4	B	505[A]	GUA	O2-C1-C2	2.52	122.14	114.03
4	A	506[B]	GUA	O2-C1-O1	-2.45	117.18	123.30
3	B	500	PLP	C3-C4-C5	2.42	121.35	118.74
3	B	500	PLP	O4P-C5A-C5	2.40	113.92	109.35
3	A	500	PLP	C4A-C4-C5	-2.39	118.48	120.94
3	B	500	PLP	C6-N1-C2	2.37	123.57	119.17
4	A	506[B]	GUA	O2-C1-C2	2.37	121.66	114.03
4	A	506[B]	GUA	C3-C2-C1	-2.26	108.77	114.47

There are no chirality outliers.

All (19) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	500	PLP	C5A-O4P-P-O3P
4	A	506[B]	GUA	C1-C2-C3-C4
4	B	505[B]	GUA	C1-C2-C3-C4
4	B	505[A]	GUA	C2-C3-C4-C5
4	A	506[A]	GUA	C1-C2-C3-C4
4	A	506[A]	GUA	O1-C1-C2-C3
4	A	506[A]	GUA	O2-C1-C2-C3
4	A	506[A]	GUA	C2-C3-C4-C5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
4	A	506[B]	GUA	C3-C4-C5-O3
4	B	505[A]	GUA	O1-C1-C2-C3
4	B	505[B]	GUA	O2-C1-C2-C3
4	B	505[B]	GUA	O1-C1-C2-C3
4	B	505[B]	GUA	C3-C4-C5-O4
4	A	506[B]	GUA	C3-C4-C5-O4
4	B	505[A]	GUA	O2-C1-C2-C3
4	A	506[B]	GUA	O1-C1-C2-C3
4	A	506[B]	GUA	O2-C1-C2-C3
3	B	500	PLP	C5A-O4P-P-O1P
4	B	505[B]	GUA	C3-C4-C5-O3

There are no ring outliers.

4 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	500	PLP	2	0
3	B	500	PLP	2	0
2	A	700	ACT	1	0
2	A	600	ACT	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	448/466 (96%)	-0.61	5 (1%) 80 82	11, 19, 40, 59	0
1	B	449/466 (96%)	-0.50	8 (1%) 68 71	13, 21, 42, 63	1 (0%)
All	All	897/932 (96%)	-0.56	13 (1%) 75 78	11, 20, 42, 63	1 (0%)

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	114	LYS	3.7
1	B	385	GLY	3.6
1	A	307	GLY	3.3
1	B	307	GLY	3.2
1	B	452	PRO	3.0
1	B	114	LYS	3.0
1	A	5	LEU	2.7
1	B	9	PHE	2.6
1	B	308	GLY	2.5
1	A	306	LEU	2.4
1	B	112	ALA	2.2
1	B	6	LEU	2.1
1	A	115	ASN	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	ACT	A	700	4/4	0.90	0.21	24,27,29,30	0
4	GUA	A	506[A]	9/9	0.93	0.20	10,15,20,20	6
4	GUA	A	506[B]	9/9	0.93	0.20	19,21,22,22	6
4	GUA	B	505[A]	9/9	0.95	0.13	9,14,17,19	6
4	GUA	B	505[B]	9/9	0.95	0.13	16,19,21,22	6
2	ACT	A	600	4/4	0.96	0.17	22,25,25,26	0
3	PLP	A	500	15/16	0.99	0.06	11,12,14,14	0
3	PLP	B	500	15/16	0.99	0.09	10,13,15,17	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.