



wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 17, 2023 – 11:40 PM EDT

PDB ID : 4XEV
Title : Fusion of Pyk2-FAT domain with Leupaxin LD1 motif, complexed with Leupaxin LD4 peptide
Authors : Miller, D.J.
Deposited on : 2014-12-24
Resolution : 2.01 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.01 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	167	 6% (poor fit), 81% (0-3 outliers), 7% (1 outlier), 11% (2-3 outliers), 11% (not modelled)
1	D	167	 3% (poor fit), 83% (0-3 outliers), 5% (1 outlier), 12% (2-3 outliers), 12% (not modelled)
2	C	19	 26% (poor fit), 89% (0-3 outliers), 11% (1 outlier), 11% (2-3 outliers), 11% (not modelled)
2	F	19	 26% (poor fit), 79% (0-3 outliers), 16% (1 outlier), 5% (2-3 outliers), 5% (not modelled)

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2579 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fusion protein of Protein-tyrosine kinase 2-beta FAT domain and Leupaxin LD1 motif.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	148	1111	696	193	219	3	0	0	0
1	D	147	1136	715	190	226	5	0	4	0

There are 28 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	867	GLY	-	expression tag	UNP Q14289
A	868	SER	-	expression tag	UNP Q14289
A	869	HIS	-	expression tag	UNP Q14289
A	870	MET	-	expression tag	UNP Q14289
A	899	SER	CYS	engineered mutation	UNP Q14289
A	972	ALA	CYS	engineered mutation	UNP Q14289
A	1006	GLY	-	linker	UNP Q14289
A	1007	GLY	-	linker	UNP Q14289
A	1995	SER	-	linker	UNP Q14289
A	1996	GLY	-	linker	UNP Q14289
A	1997	GLY	-	linker	UNP Q14289
A	1998	GLY	-	linker	UNP Q14289
A	1999	GLY	-	linker	UNP Q14289
A	2000	GLY	-	linker	UNP Q14289
D	867	GLY	-	expression tag	UNP Q14289
D	868	SER	-	expression tag	UNP Q14289
D	869	HIS	-	expression tag	UNP Q14289
D	870	MET	-	expression tag	UNP Q14289
D	899	SER	CYS	engineered mutation	UNP Q14289
D	972	ALA	CYS	engineered mutation	UNP Q14289
D	1006	GLY	-	linker	UNP Q14289
D	1994	GLY	-	linker	UNP Q14289
D	1995	SER	-	linker	UNP Q14289
D	1996	GLY	-	linker	UNP Q14289

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	1997	GLY	-	linker	UNP Q14289
D	1998	GLY	-	linker	UNP Q14289
D	1999	GLY	-	linker	UNP Q14289
D	2000	GLY	-	linker	UNP Q14289

- Molecule 2 is a protein called 19-mer peptide containing Leupaxin LD4 motif.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
			Total	C	N	O				S
2	C	19	135	82	22	29	2	0	0	0
2	F	18	126	78	22	24	2	0	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	32	Total	O	0	0
			32	32		
3	D	38	Total	O	0	0
			38	38		
3	F	1	Total	O	0	0
			1	1		

4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	52.08Å 79.23Å 53.23Å 90.00° 117.62° 90.00°	Depositor
Resolution (Å)	27.26 – 2.01 27.26 – 2.01	Depositor EDS
% Data completeness (in resolution range)	95.8 (27.26-2.01) 95.9 (27.26-2.01)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.04	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.77 (at 2.01Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, R_{free}	0.204 , 0.250 0.206 , 0.252	Depositor DCC
R_{free} test set	1277 reflections (5.19%)	wwPDB-VP
Wilson B-factor (Å ²)	41.5	Xtrriage
Anisotropy	0.659	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 55.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.016 for l,-k,h	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2579	wwPDB-VP
Average B, all atoms (Å ²)	60.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.06% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.35	0/1117	0.48	0/1514
1	D	0.39	0/1153	0.53	0/1560
2	C	0.22	0/135	0.41	0/182
2	F	0.26	0/126	0.45	0/169
All	All	0.36	0/2531	0.50	0/3425

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1111	0	1142	6	0
1	D	1136	0	1192	5	0
2	C	135	0	124	1	0
2	F	126	0	125	2	0
3	A	32	0	0	1	0
3	D	38	0	0	0	0
3	F	1	0	0	0	0
All	All	2579	0	2583	13	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 13 close contacts within the same asymmetric unit are listed below, sorted by their

clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:921:ILE:HD12	2:C:93:LEU:HD22	1.80	0.64
1:A:942:THR:OG1	1:A:990:LEU:HD13	1.99	0.63
1:D:2008:LEU:O	1:D:2012:GLU:HG2	2.05	0.57
1:A:935:SER:O	1:A:939:ILE:HG12	2.08	0.53
1:D:897:GLU:O	1:D:901:LEU:HG	2.12	0.49

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	144/167 (86%)	143 (99%)	1 (1%)	0	100	100
1	D	147/167 (88%)	147 (100%)	0	0	100	100
2	C	17/19 (90%)	16 (94%)	1 (6%)	0	100	100
2	F	16/19 (84%)	16 (100%)	0	0	100	100
All	All	324/372 (87%)	322 (99%)	2 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	121/141 (86%)	116 (96%)	5 (4%)	30	28
1	D	129/141 (92%)	128 (99%)	1 (1%)	81	86
2	C	13/15 (87%)	12 (92%)	1 (8%)	13	8
2	F	12/15 (80%)	12 (100%)	0	100	100
All	All	275/312 (88%)	268 (98%)	7 (2%)	46	49

5 of 7 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	973	LYS
1	A	1003	LEU
2	C	104	GLN
1	D	898	LEU
1	A	938	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	996	GLN
1	D	900	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	148/167 (88%)	0.38	10 (6%) 17 16	38, 58, 101, 119	0
1	D	147/167 (88%)	0.19	5 (3%) 45 44	36, 53, 88, 102	0
2	C	19/19 (100%)	1.27	5 (26%) 0 0	59, 75, 100, 104	0
2	F	18/19 (94%)	0.86	5 (27%) 0 0	59, 71, 98, 106	0
All	All	332/372 (89%)	0.37	25 (7%) 14 13	36, 57, 97, 119	0

The worst 5 of 25 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	876	THR	4.9
2	F	86	LYS	3.4
1	D	876	THR	3.3
2	F	101	THR	3.3
1	A	2006	ALA	3.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.