

# Full wwPDB X-ray Structure Validation Report (i)

#### Jun 11, 2024 – 05:55 PM EDT

PDB ID	:	1XGF
Title	:	Backbone Structure of COCOSIN, an 11S storage protein from cocos nucifera
Authors	:	Balasundaresan, D.; Ponnuswamy, M.N.
Deposited on	:	2004-09-17
Resolution	:	2.61  Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.20.1
EDS	:	2.36.2
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36.2

Clashscore

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\text{-}RAY\;DIFFRACTION$ 

The reported resolution of this entry is 2.61 Å.

141614

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.

Metric	Percent	tile Ranks	Value			
Clashscore			7			
Wor.	те Ге	-	Better			
Pe	centile relative to all X-ray structures					
Percentile relative to X-ray structures of similar resolution						
Motrio	Whole archive	Simi	lar resolution			
wietric	(# Entries)	(# Entries, )	$\operatorname{resolution} \operatorname{range}(\operatorname{\AA}))$			

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

4168(2.64-2.60)

Mol	Chain	Length	Quality of chain		
1	А	382	81%	19%	_
1	В	382	76%	23%	•



#### $1 \mathrm{XGF}$

## 2 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 3056 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	Λ	200	Total	С	Ν	Ο	0	0	0
I A	362	1528	764	382	382	0	0	0	
1	1 D	P 200	Total	С	Ν	Ο	0	0	0
	382	1528	764	382	382	0	0	0	

• Molecule 1 is a protein called cocosin.



## 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: cocosin



## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	Н 3	Depositor
Cell constants	93.66Å $93.66$ Å $216.45$ Å	Densite
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $120.00^{\circ}$	Depositor
$\mathbf{P}_{\text{acclution}}(\hat{\mathbf{A}})$	2.80 - 2.61	Depositor
Resolution (A)	29.60 - 2.61	EDS
% Data completeness	(Not available) $(2.80-2.61)$	Depositor
(in resolution range)	94.7(29.60-2.61)	EDS
R <sub>merge</sub>	(Not available)	Depositor
R <sub>sym</sub>	(Not available)	Depositor
$< I/\sigma(I) > 1$	$0.96 (at 2.61 \text{\AA})$	Xtriage
Refinement program	SHELXL-97	Depositor
D D.	0.240 , $0.294$	Depositor
$\Pi, \Pi_{free}$	0.405 , (Not available)	DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor $(Å^2)$	34.5	Xtriage
Anisotropy	0.020	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.07, 159.0	EDS
L-test for twinning <sup>2</sup>	$< L >=0.49, < L^2>=0.32$	Xtriage
Estimated twinning fraction	0.045 for h,-h-k,-l	Xtriage
$F_o, F_c$ correlation	0.67	EDS
Total number of atoms	3056	wwPDB-VP
Average B, all atoms $(Å^2)$	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.17% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

## 5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

There are no protein, RNA or DNA chains available to summarize Z scores of covalent bonds and angles.

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	А	0	69
1	В	0	80
All	All	0	149

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (149) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	А	108	UNK	Peptide
1	А	111	UNK	Mainchain
1	А	118	UNK	Mainchain
1	А	126	UNK	Mainchain
1	А	134	UNK	Mainchain
1	А	141	UNK	Peptide,Mainchain
1	А	150	UNK	Mainchain
1	А	152	UNK	Mainchain
1	А	153	UNK	Mainchain
1	А	158	UNK	Mainchain
1	А	219	UNK	Mainchain
1	А	220	UNK	Mainchain
1	А	221	UNK	Mainchain
1	А	223	UNK	Mainchain
1	А	235	UNK	Mainchain
1	А	27	UNK	Mainchain

Continued on next page...



Mol	Chain	Res	Type	Group
1	A	32	UNK	Mainchain
1	A	321	UNK	Mainchain
1	A	326	UNK	Mainchain
1	A	333	UNK	Mainchain
1	A	338	UNK	Mainchain
1	A	342	UNK	Mainchain
1	Δ	348	UNK	Mainchain
1	Δ	349	UNK	Mainchain
1	Δ	355	UNK	Mainchain
1		357	UNK	Mainchain
1		358	UNK	Mainchain
1	A	300	UNK	Poptido Mainchain
1	A	261	UNK	Teptue, Mainchain
1	A	301		Mainchain
1	A	300		Mainchain
1	A	309		Mainchain
1	A	377		Mainchain
1	A	382		Mainchain
1	A	384	UNK	Mainchain
1	A	387	UNK	Mainchain
1	A	395	UNK	Mainchain
1	A	397	UNK	Mainchain
1	A	398	UNK	Mainchain
1	A	399	UNK	Mainchain
1	А	404	UNK	Mainchain
1	A	407	UNK	Mainchain
1	А	416	UNK	Peptide,Mainchain
1	А	429	UNK	Mainchain
1	А	435	UNK	Mainchain
1	А	436	UNK	Mainchain
1	А	440	UNK	Mainchain
1	А	441	UNK	Mainchain
1	А	447	UNK	Mainchain
1	А	448	UNK	Mainchain
1	А	449	UNK	Mainchain
1	А	461	UNK	Mainchain
1	А	468	UNK	Mainchain
1	А	472	UNK	Mainchain
1	А	473	UNK	Mainchain
1	А	478	UNK	Peptide, Mainchain
1	А	485	UNK	Mainchain
1	А	72	UNK	Mainchain
1	А	8	UNK	Mainchain

Continued from previous page...

Continued on next page...



Mol	Chain	Res		Group
1	A	86	UNK	Mainchain
1	A	88	UNK	Mainchain
1	A	89	UNK	Mainchain
1	A	9	UNK	Mainchain
1	A	91	UNK	Mainchain
1	A	92	UNK	Peptide Mainchain
1	B	10	UNK	Mainchain
1	B	111	UNK	Mainchain
1	B	113	UNK	Mainchain
1	B	115	UNK	Mainchain
1	B	118	UNK	Mainchain
1	B	120	UNK	Mainchain
1	B	138	UNK	Mainchain
1	B	140	UNK	Mainchain
1	B	140	UNK	Mainchain
1	B	140	UNK	Mainchain
1	B	15 157	UNK	Mainchain
1	B	107 167	UNK	Mainchain
1	D	160	UNK	Mainchain
1	D	109	UNK	Mainchain
1	D	170	UNK	Mainchain
1	D	19	UNK	Mainchain
1	D	200		Mainchain
1	D D	212		Mainchain
1	D	219	UNK	Mainchain
1	B	220		Mainchain
1	B	222		Mainchain
1	B	232		Mainchain
1	B	233		
1	B	321		Peptide, Mainchain
1	B	331	UNK	Mainchain
1	B	332	UNK	Mainchain
1	В	333	UNK	Mainchain
1	B	336	UNK	Mainchain
1	B	339	UNK	Mainchain
1	B	340	UNK	Mainchain
1	B	348	UNK	Peptide,Mainchain
1	B	349	UNK	Mainchain
1	B	351	UNK	Mainchain
1	В	36	UNK	Mainchain
1	В	360	UNK	Mainchain
1	В	369	UNK	Mainchain
1	В	377	UNK	Mainchain

Continued from previous page...

Continued on next page...



Mol Chain Deg Tung Choun					
	D	nes	Type	Group	
1	В	379	UNK	Mainchain	
1	B	38	UNK	Mainchain	
1	В	382	UNK	Mainchain	
1	В	383	UNK	Mainchain	
1	В	384	UNK	Mainchain	
1	В	385	UNK	Mainchain	
1	В	388	UNK	Mainchain	
1	В	390	UNK	Mainchain	
1	В	395	UNK	Mainchain	
1	В	396	UNK	Mainchain	
1	В	397	UNK	Mainchain	
1	В	417	UNK	Mainchain	
1	В	43	UNK	Mainchain	
1	В	433	UNK	Mainchain	
1	В	438	UNK	Mainchain	
1	В	440	UNK	Mainchain	
1	В	443	UNK	Mainchain	
1	В	448	UNK	Mainchain	
1	В	454	UNK	Mainchain	
1	В	458	UNK	Mainchain	
1	В	460	UNK	Mainchain	
1	В	461	UNK	Mainchain	
1	В	464	UNK	Mainchain	
1	В	482	UNK	Mainchain	
1	В	484	UNK	Mainchain	
1	В	485	UNK	Peptide.Mainchain	
1	В	487	UNK	Mainchain	
1	В	50	UNK	Mainchain	
1	B	52	UNK	Mainchain	
1	В	58	UNK	Mainchain	
1	B	61	UNK	Mainchain	
1	B	63	UNK	Mainchain	
1	B	65	UNK	Mainchain	
1	B	69	UNK	Mainchain	
1	B	7	UNK	Mainchain	
1	B	73	UNK	Mainchain	
1	B	8	UNK	Mainchain	
1	R	84	UNK	Mainchain	
1	R	85	UNK	Pentide	
1	R	80	INK	Mainchain	
1	B	09	UNK	Mainchain	
L L	Ы	90	UNK	Ivianichani	

Continued from previous page...



### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	1528	0	54	9	0
1	В	1528	0	47	13	0
All	All	3056	0	101	22	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (22) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom 1	Atom 2	Interatomic	Clash	
Atom-1	Atom-2	distance (Å)	overlap (Å)	
1:A:492:UNK:C	1:A:493:UNK:H	1.27	1.48	
1:A:492:UNK:C	1:A:493:UNK:N	1.93	1.30	
1:B:416:UNK:C	1:B:417:UNK:N	1.96	1.27	
1:B:416:UNK:C	1:B:417:UNK:H2	1.57	1.11	
1:A:492:UNK:CA	1:A:493:UNK:H	1.72	1.02	
1:B:416:UNK:CA	1:B:417:UNK:H2	1.83	0.90	
1:A:492:UNK:C	1:A:493:UNK:C	2.57	0.82	
1:A:492:UNK:CA	1:A:493:UNK:N	2.39	0.79	
1:B:416:UNK:CA	1:B:417:UNK:N	2.43	0.76	
1:A:492:UNK:C	1:A:493:UNK:CA	2.73	0.65	
1:A:328:UNK:O	1:A:329:UNK:C	2.49	0.60	
1:B:416:UNK:C	1:B:417:UNK:H	2.11	0.59	
1:B:479:UNK:O	1:B:482:UNK:O	2.26	0.53	
1:B:492:UNK:O	1:B:493:UNK:C	2.60	0.49	
1:B:79:UNK:O	1:B:132:UNK:N	2.50	0.45	
1:B:418:UNK:O	1:B:419:UNK:C	2.65	0.43	
1:A:359:UNK:O	1:A:360:UNK:C	2.66	0.43	
1:B:321:UNK:O	1:B:322:UNK:C	2.67	0.43	
1:B:78:UNK:O	1:B:80:UNK:N	2.53	0.41	
1:B:74:UNK:O	1:B:119:UNK:N	2.53	0.41	
1:B:322:UNK:O	1:B:323:UNK:C	2.67	0.40	
1:A:73:UNK:O	1:A:74:UNK:C	2.69	0.40	

There are no symmetry-related clashes.



### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

There are no protein backbone outliers to report in this entry.

#### 5.3.2 Protein sidechains (i)

There are no protein residues with a non-rotameric sidechain to report in this entry.

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

#### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry (i)

There are no ligands in this entry.

### 5.7 Other polymers (i)

There are no such residues in this entry.

#### 5.8 Polymer linkage issues (i)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	В	7
1	А	5

All chain breaks are listed below:



Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	А	251:UNK	С	321:UNK	N	28.56
1	В	251:UNK	С	321:UNK	N	28.17
1	В	178:UNK	С	200:UNK	N	14.82
1	А	178:UNK	С	200:UNK	N	13.68
1	В	92:UNK	С	108:UNK	N	5.16
1	В	416:UNK	С	417:UNK	N	1.96
1	A	492:UNK	С	493:UNK	N	1.93
1	В	38:UNK	С	39:UNK	N	1.19
1	В	458:UNK	С	459:UNK	N	1.19
1	А	86:UNK	С	87:UNK	N	1.18
1	А	491:UNK	С	492:UNK	N	1.12
1	В	8:UNK	С	9:UNK	N	1.09



## 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

Unable to reproduce the depositors R factor - this section is therefore empty.

## 6.2 Non-standard residues in protein, DNA, RNA chains (i)

Unable to reproduce the depositors R factor - this section is therefore empty.

## 6.3 Carbohydrates (i)

Unable to reproduce the depositors R factor - this section is therefore empty.

## 6.4 Ligands (i)

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.5 Other polymers (i)

Unable to reproduce the depositors R factor - this section is therefore empty.

