



# Full wwPDB X-ray Structure Validation Report i

Nov 23, 2023 – 12:05 AM JST

PDB ID : 7XIK  
Title : SARS-CoV-2-Omicron-RBD and B38-GWP/P-VK antibody complex  
Authors : Shi, R.; Wang, Y.; Wu, Z.; Han, X.; Yan, J.  
Deposited on : 2022-04-13  
Resolution : 2.89 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>  
with specific help available everywhere you see the i symbol.

The types of validation reports are described at  
<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.36
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*


The reported resolution of this entry is 2.89 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1957 (2.90-2.90)
Clashscore	141614	2172 (2.90-2.90)
Ramachandran outliers	138981	2115 (2.90-2.90)
Sidechain outliers	138945	2117 (2.90-2.90)
RSRZ outliers	127900	1906 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



## 2 Entry composition [\(i\)](#)

There are 3 unique types of molecules in this entry. The entry contains 4851 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Spike protein S1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	195	Total	C 1563	N 1008	O 264	S 283	8	0	0

There are 21 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	339	ASP	GLY	variant	UNP P0DTC2
A	371	LEU	SER	variant	UNP P0DTC2
A	373	PRO	SER	variant	UNP P0DTC2
A	375	PHE	SER	variant	UNP P0DTC2
A	417	ASN	LYS	variant	UNP P0DTC2
A	440	LYS	ASN	variant	UNP P0DTC2
A	446	SER	GLY	variant	UNP P0DTC2
A	477	ASN	SER	variant	UNP P0DTC2
A	478	LYS	THR	variant	UNP P0DTC2
A	484	ALA	GLU	variant	UNP P0DTC2
A	493	ARG	GLN	variant	UNP P0DTC2
A	496	SER	GLY	variant	UNP P0DTC2
A	498	ARG	GLN	variant	UNP P0DTC2
A	501	TYR	ASN	variant	UNP P0DTC2
A	505	HIS	TYR	variant	UNP P0DTC2
A	538	HIS	-	expression tag	UNP P0DTC2
A	539	HIS	-	expression tag	UNP P0DTC2
A	540	HIS	-	expression tag	UNP P0DTC2
A	541	HIS	-	expression tag	UNP P0DTC2
A	542	HIS	-	expression tag	UNP P0DTC2
A	543	HIS	-	expression tag	UNP P0DTC2

- Molecule 2 is a protein called B38 Fab heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	218	Total	C	N	O	S	0	1	0
			1628	1026	275	320	7			


- Molecule 3 is a protein called B38 Fab light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	L	216	Total	C	N	O	S	3	1	0
			1660	1046	274	334	6			


### 3 Residue-property plots [\(i\)](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Spike protein S1



- Molecule 2: B38 Fab heavy chain



- Molecule 3: B38 Fab light chain



## 4 Data and refinement statistics i

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	84.96 Å    145.86 Å    144.15 Å 90.00°    90.00°    90.00°	Depositor
Resolution (Å)	73.41 – 2.89 73.41 – 2.89	Depositor EDS
% Data completeness (in resolution range)	99.7 (73.41-2.89) 99.7 (73.41-2.89)	Depositor EDS
$R_{merge}$	0.26	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) >$ <sup>1</sup>	2.21 (at 2.91 Å)	Xtriage
Refinement program	PHENIX 1.20.1-4487	Depositor
$R$ , $R_{free}$	0.211 , 0.263 0.211 , 0.263	Depositor DCC
$R_{free}$ test set	2009 reflections (9.89%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	54.0	Xtriage
Anisotropy	0.420	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 44.3	EDS
L-test for twinning <sup>2</sup>	$<  L  > = 0.47$ , $< L^2 > = 0.30$	Xtriage
Estimated twinning fraction	0.025 for 1/2*h-1/2*k,-3/2*h-1/2*k,-l 0.032 for 1/2*h+1/2*k,3/2*h-1/2*k,-l	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	4851	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.41% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $< |L| >$ ,  $< L^2 >$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality (i)

### 5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.29	0/1610	0.53	0/2192
2	H	0.30	0/1671	0.53	0/2278
3	L	0.41	0/1701	0.53	0/2312
All	All	0.34	0/4982	0.53	0/6782

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1563	0	1489	23	0
2	H	1628	0	1594	29	0
3	L	1660	0	1618	24	0
All	All	4851	0	4701	71	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (71) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:388:ASN:HB3	1:A:527:PRO:HD2	1.80	0.63

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:L:40:LYS:NZ	3:L:82:GLU:O	2.31	0.62
1:A:403:ARG:HG3	1:A:505:HIS:HA	1.82	0.61
3:L:91:GLN:HE21	3:L:99:THR:H	1.47	0.61
1:A:408:ARG:HH12	1:A:409:GLN:HE21	1.48	0.60
1:A:393:THR:HG21	1:A:518:LEU:H	1.67	0.60
2:H:35:SER:OG	2:H:98:GLU:OE2	2.21	0.59
2:H:128:PRO:HG3	2:H:140:LEU:HB3	1.85	0.58
3:L:24:ARG:HG2	3:L:71:GLU:HG2	1.87	0.57
2:H:190:SER:HB2	2:H:194:GLN:HB2	1.86	0.57
2:H:11:LEU:HD12	2:H:112:THR:HB	1.86	0.57
2:H:124:PHE:CE2	3:L:126:GLN:HG3	2.41	0.56
1:A:358:ILE:HB	1:A:395:VAL:O	2.05	0.56
3:L:195:ALA:HB2	3:L:210:SER:HB3	1.90	0.53
2:H:2:VAL:HA	2:H:25:SER:O	2.07	0.53
3:L:34:LEU:HB2	3:L:52:ALA:HB2	1.90	0.52
1:A:409:GLN:OE1	1:A:418:ILE:N	2.43	0.51
2:H:167:THR:HG23	2:H:180:LEU:HD21	1.92	0.51
1:A:365:TYR:CG	1:A:387:LEU:HD23	2.46	0.51
3:L:90:GLN:HG2	3:L:91:GLN:N	2.26	0.50
1:A:354:ASN:O	1:A:398:ASP:HA	2.11	0.50
3:L:24:ARG:NH1	3:L:71:GLU:OE2	2.45	0.50
1:A:461:LEU:HG	1:A:465:GLU:HB3	1.95	0.49
3:L:6:GLN:O	3:L:102:GLN:NE2	2.45	0.49
3:L:90:GLN:HG2	3:L:91:GLN:H	1.78	0.49
2:H:48:VAL:HG13	2:H:63:VAL:HG11	1.95	0.49
2:H:4:LEU:HD21	2:H:27:PHE:HZ	1.78	0.49
2:H:132:SER:OG	2:H:133:THR:N	2.46	0.49
3:L:144:ARG:NH2	3:L:165:VAL:HG11	2.27	0.49
3:L:8:PRO:HG2	3:L:11:LEU:HB2	1.95	0.48
3:L:38:GLN:HB2	3:L:48:LEU:HD11	1.95	0.48
1:A:385:THR:O	1:A:386:LYS:HG2	2.14	0.48
3:L:192:LYS:HE2	3:L:212:ASN:HB3	1.96	0.47
2:H:158:SER:O	2:H:158:SER:OG	2.28	0.46
2:H:28:ILE:HD12	2:H:31:TRP:CZ3	2.50	0.46
2:H:6:GLU:OE2	2:H:108:GLY:N	2.48	0.46
3:L:139:ASN:OD1	3:L:140:ASN:ND2	2.48	0.46
2:H:38:ARG:HD3	2:H:48:VAL:HG22	1.96	0.46
2:H:212:ARG:HD3	2:H:214:GLU:OE1	2.15	0.46
2:H:17[B]:SER:HA	2:H:82:MET:O	2.16	0.45
2:H:17[A]:SER:HA	2:H:82:MET:O	2.16	0.45
2:H:217:SER:O	2:H:217:SER:OG	2.31	0.45

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:374:PHE:O	1:A:436:TRP:HA	2.17	0.45
1:A:420:ASP:OD2	2:H:56:SER:OG	2.23	0.45
1:A:376:THR:O	1:A:434:ILE:HA	2.17	0.45
1:A:497:PHE:CE2	1:A:507:PRO:HB3	2.51	0.44
2:H:47:TRP:HZ2	2:H:50:VAL:HG12	1.82	0.44
1:A:412:PRO:HG3	1:A:429:PHE:HB3	1.99	0.44
1:A:458:LYS:HE2	2:H:31:TRP:CD2	2.53	0.44
2:H:98:GLU:HB3	2:H:102:MET:HG2	2.00	0.44
1:A:388:ASN:N	1:A:388:ASN:OD1	2.51	0.43
3:L:30:PRO:HD2	3:L:33:TYR:CE2	2.53	0.43
2:H:133:THR:HG23	2:H:135:GLY:H	1.85	0.42
3:L:36:TRP:CZ3	3:L:89:CYS:HB3	2.54	0.42
1:A:403:ARG:NH1	3:L:94:SER:HB3	2.34	0.42
1:A:370:ASN:OD1	1:A:370:ASN:N	2.49	0.42
1:A:456:PHE:HB3	1:A:473:TYR:CG	2.55	0.42
2:H:202:HIS:CD2	2:H:204:PRO:HD2	2.54	0.42
3:L:115:PRO:HB3	3:L:141:PHE:HB3	2.01	0.42
3:L:33:TYR:CE2	3:L:93:ASN:HB2	2.55	0.41
1:A:380:TYR:O	1:A:430:THR:HA	2.20	0.41
2:H:85:LEU:HD22	2:H:113:VAL:HG21	2.02	0.41
2:H:72:HIS:CD2	2:H:74:SER:HB3	2.56	0.41
1:A:342:PHE:CZ	1:A:368:LEU:HD22	2.56	0.41
3:L:34:LEU:HA	3:L:90:GLN:O	2.21	0.41
3:L:203:LEU:HD22	3:L:207:VAL:HG21	2.02	0.41
1:A:478:LYS:HA	1:A:479:PRO:HD3	1.92	0.40
2:H:12:VAL:HG23	2:H:113:VAL:HG22	2.03	0.40
2:H:36:TRP:CE2	2:H:80:LEU:HB2	2.56	0.40
3:L:11:LEU:HD23	3:L:106:LEU:HD13	2.04	0.40
2:H:124:PHE:CZ	3:L:126:GLN:HG3	2.57	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	193/225 (86%)	185 (96%)	8 (4%)	0	100	100
2	H	217/222 (98%)	209 (96%)	8 (4%)	0	100	100
3	L	215/219 (98%)	212 (99%)	3 (1%)	0	100	100
All	All	625/666 (94%)	606 (97%)	19 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	169/200 (84%)	161 (95%)	8 (5%)	26	59
2	H	182/184 (99%)	180 (99%)	2 (1%)	73	92
3	L	189/191 (99%)	180 (95%)	9 (5%)	25	58
All	All	540/575 (94%)	521 (96%)	19 (4%)	36	70

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	341	VAL
1	A	359	SER
1	A	377	PHE
1	A	382	VAL
1	A	388	ASN
1	A	389	ASP
1	A	494	SER
1	A	496	SER
2	H	158	SER
2	H	199	ASN
3	L	1	ASP
3	L	7	SER
3	L	34	LEU
3	L	66	SER
3	L	91	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
3	L	94	SER
3	L	154	ASN
3	L	176	SER
3	L	187	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	417	ASN
2	H	72	HIS
3	L	91	GLN
3	L	139	ASN
3	L	140	ASN

### 5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [\(i\)](#)

There are no ligands in this entry.

### 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	195/225 (86%)	0.41	5 (2%) 56 52	34, 51, 77, 95	0
2	H	218/222 (98%)	0.40	4 (1%) 68 67	29, 41, 72, 112	0
3	L	216/219 (98%)	0.26	1 (0%) 91 91	32, 45, 61, 81	0
All	All	629/666 (94%)	0.36	10 (1%) 72 71	29, 45, 73, 112	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	H	0	ASP	5.3
2	H	132	SER	2.6
2	H	133	THR	2.6
1	A	372	ALA	2.4
3	L	79	LEU	2.3
1	A	387	LEU	2.2
2	H	1	GLU	2.1
1	A	368	LEU	2.1
1	A	524	VAL	2.0
1	A	369	TYR	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

## 6.4 Ligands [\(i\)](#)

There are no ligands in this entry.

## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.