

Full wwPDB X-ray Structure Validation Report (i)

Oct 23, 2024 – 11:22 PM EDT

:	1XLX
:	Catalytic Domain Of Human Phosphodiesterase 4B In Complex With Cilomi-
	last
:	Card, G.L.; England, B.P.; Suzuki, Y.; Fong, D.; Powell, B.; Lee, B.; Luu, C.;
	Tabrizizad, M.; Gillette, S.; Ibrahim, P.N.; Artis, D.R.; Bollag, G.; Milburn,
	M.V.; Kim, SH.; Schlessinger, J.; Zhang, K.Y.J.
:	2004-09-30
:	2.19 Å(reported)
	::

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

:	4.02b-467
:	2022.3.0, CSD as543be (2022)
:	1.20.1
:	3.0
:	1.1.7(2018)
:	20231227.v01 (using entries in the PDB archive December 27th 2023)
:	9.0.003 (Gargrove)
:	1.0.11
:	Engh & Huber (2001)
:	Parkinson et al. (1996)
:	2.39
	: : : : : : : : : : : : : : : : : : :



1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY\;DIFFRACTION$

The reported resolution of this entry is 2.19 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R_{free}	164625	5791 (2.20-2.20)
Clashscore	180529	6634 (2.20-2.20)
Ramachandran outliers	177936	6560 (2.20-2.20)
Sidechain outliers	177891	6561 (2.20-2.20)
RSRZ outliers	164620	5791 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain				
1	А	398	3% 66%	14%	·	19%	
1	В	398	2% 5 9%	21%	•	19%	



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 5356 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
1	А	323	Total 2617	C 1653	N 441	O 503	S 20	7	0	0
1	В	323	Total 2612	C 1650	N 439	O 503	S 20	4	0	0

• Molecule 1 is a protein called cAMP-specific 3',5'-cyclic phosphodiesterase 4B.

Chain	Residue	Modelled	Actual	Comment	Reference
A	131	MET	-	cloning artifact	UNP Q07343
А	132	GLY	-	cloning artifact	UNP Q07343
А	133	SER	-	cloning artifact	UNP Q07343
А	134	SER	-	cloning artifact	UNP Q07343
А	135	HIS	-	cloning artifact	UNP Q07343
А	136	HIS	-	cloning artifact	UNP Q07343
A	137	HIS	-	cloning artifact	UNP Q07343
А	138	HIS	-	cloning artifact	UNP Q07343
А	139	HIS	-	cloning artifact	UNP Q07343
А	140	HIS	-	cloning artifact	UNP Q07343
А	141	SER	-	cloning artifact	UNP Q07343
А	142	SER	-	cloning artifact	UNP Q07343
А	143	GLY	-	cloning artifact	UNP Q07343
А	144	LEU	-	cloning artifact	UNP Q07343
А	145	VAL	-	cloning artifact	UNP Q07343
A	146	PRO	-	cloning artifact	UNP Q07343
А	147	ARG	-	cloning artifact	UNP Q07343
A	148	GLY	-	cloning artifact	UNP Q07343
А	149	SER	-	cloning artifact	UNP Q07343
A	150	HIS	-	cloning artifact	UNP Q07343
А	151	MET	-	cloning artifact	UNP Q07343
A	194	CME	CYS	modified residue	UNP Q07343
В	131	MET	-	cloning artifact	UNP Q07343
В	132	GLY	-	cloning artifact	UNP Q07343
В	133	SER	-	cloning artifact	UNP Q07343

There are 44 discrepancies between the modelled and reference sequences:



Chain	Residue	Modelled	Actual	Comment	Reference
В	134	SER	-	cloning artifact	UNP Q07343
В	135	HIS	-	cloning artifact	UNP Q07343
В	136	HIS	-	cloning artifact	UNP Q07343
В	137	HIS	-	cloning artifact	UNP Q07343
В	138	HIS	-	cloning artifact	UNP Q07343
В	139	HIS	-	cloning artifact	UNP Q07343
В	140	HIS	-	cloning artifact	UNP Q07343
В	141	SER	-	cloning artifact	UNP Q07343
В	142	SER	-	cloning artifact	UNP Q07343
В	143	GLY	-	cloning artifact	UNP Q07343
В	144	LEU	-	cloning artifact	UNP Q07343
В	145	VAL	-	cloning artifact	UNP Q07343
В	146	PRO	-	cloning artifact	UNP Q07343
В	147	ARG	-	cloning artifact	UNP Q07343
В	148	GLY	-	cloning artifact	UNP Q07343
В	149	SER	-	cloning artifact	UNP Q07343
В	150	HIS	-	cloning artifact	UNP Q07343
В	151	MET	-	cloning artifact	UNP Q07343
В	194	CME	CYS	modified residue	UNP Q07343

• Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	1	Total Zn 1 1	0	0
2	В	1	Total Zn 1 1	0	0

• Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	1	Total Mg 1 1	0	0
3	В	1	Total Mg 1 1	0	0

• Molecule 4 is CILOMILAST (three-letter code: CIO) (formula: $C_{20}H_{25}NO_4$).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	Δ	1	Total	С	Ν	0	0	0
4 A	1	25	20	1	4	0	0	
4	В	1 Total C N O		1 Total C N O	0	0		
4 B	L	25	20	1	4	0	0	

• Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	А	35	$\begin{array}{cc} \text{Total} & \text{O} \\ 35 & 35 \end{array}$	0	0
5	В	38	Total O 38 38	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: cAMP-specific 3',5'-cyclic phosphodiesterase 4B





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	89.27Å 94.26Å 106.08Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Bosolution (Å)	70.71 - 2.19	Depositor
Resolution (A)	70.46 - 2.19	EDS
% Data completeness	92.8 (70.71-2.19)	Depositor
(in resolution range)	92.8 (70.46-2.19)	EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$2.06 (at 2.20 \text{\AA})$	Xtriage
Refinement program	REFMAC 5.1.25	Depositor
P. P.	0.232 , 0.282	Depositor
n, n_{free}	0.239 , 0.288	DCC
R_{free} test set	2198 reflections (5.07%)	wwPDB-VP
Wilson B-factor $(Å^2)$	41.2	Xtriage
Anisotropy	0.667	Xtriage
Bulk solvent $k_{sol}(e/A^3), B_{sol}(A^2)$	$0.35 \;,\; 53.5$	EDS
L-test for $twinning^2$	$ < L >=0.48, < L^2>=0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5356	wwPDB-VP
Average B, all atoms $(Å^2)$	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 51.88 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.2661e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ZN, CME, CIO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal Chain		Bo	nd lengths	Bond angles		
	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	0.89	0/2661	1.03	10/3608~(0.3%)	
1	В	0.99	3/2655~(0.1%)	1.06	13/3600~(0.4%)	
All	All	0.94	3/5316~(0.1%)	1.05	23/7208~(0.3%)	

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	В	224	GLU	CD-OE1	7.27	1.33	1.25
1	В	227	TYR	CD1-CE1	-5.81	1.30	1.39
1	В	183	VAL	CB-CG2	-5.02	1.42	1.52

Mol	Chain	Res	Type	Atoms	Ζ	$Observed(^{o})$	$Ideal(^{o})$
1	В	261	ASP	CB-CG-OD2	9.69	127.02	118.30
1	В	299	ASP	CB-CG-OD2	7.99	125.49	118.30
1	А	340	ASP	CB-CG-OD2	7.44	125.00	118.30
1	А	475	ASP	CB-CG-OD1	7.07	124.66	118.30
1	В	468	ASP	CB-CG-OD2	6.83	124.45	118.30
1	В	225	ASP	CB-CG-OD2	6.67	124.30	118.30
1	А	230	ASP	CB-CG-OD2	6.62	124.25	118.30
1	В	261	ASP	CB-CG-OD1	-6.58	112.38	118.30
1	А	468	ASP	CB-CG-OD2	6.45	124.10	118.30
1	В	340	ASP	CB-CG-OD2	6.38	124.04	118.30
1	А	214	ASP	CB-CG-OD2	5.96	123.66	118.30
1	В	256	ASP	CB-CG-OD2	5.81	123.53	118.30
1	А	460	ASP	CB-CG-OD1	5.68	123.42	118.30
1	А	379	ASP	CB-CG-OD2	5.45	123.21	118.30
1	В	221	MET	CG-SD-CE	5.43	108.88	100.20
1	А	321	ASP	CB-CG-OD2	5.39	123.15	118.30

All (23) bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	А	375	ASP	CB-CG-OD2	5.38	123.14	118.30
1	В	277	ASP	CB-CG-OD2	5.22	123.00	118.30
1	А	484	ILE	CB-CA-C	-5.19	101.22	111.60
1	В	318	GLU	CA-CB-CG	-5.18	102.00	113.40
1	В	375	ASP	CB-CG-OD2	5.17	122.95	118.30
1	В	465	ASP	CB-CG-OD1	5.10	122.89	118.30
1	В	392	ASP	CB-CG-OD2	5.05	122.84	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	2617	0	2546	30	0
1	В	2612	0	2541	48	0
2	А	1	0	0	0	0
2	В	1	0	0	0	0
3	А	1	0	0	0	0
3	В	1	0	0	0	0
4	А	25	0	24	4	0
4	В	25	0	24	1	0
5	А	35	0	0	0	0
5	В	38	0	0	1	0
All	All	5356	0	5135	80	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (80) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:220:MET:CE	1:B:220:MET:SD	2.02	1.48
1:B:207:LYS:NZ	1:B:207:LYS:HB3	1.75	1.00
1:B:207:LYS:HB3	1:B:207:LYS:HZ3	1.32	0.92



		Interatomic	Clash	
Atom-1	Atom-2	distance $(Å)$	overlap (Å)	
1:A:234:HIS:ND1	1:A:413:GLU:OE2	2.17	0.77	
1:A:340:ASP:OD1	1:A:385:ARG:HD3	1.87	0.75	
1:B:376:ASN:CG	1:B:377:TYB:H	1.95	0.70	
1:B:217:ILE:HG22	1:B:221:MET:CE	2.24	0.68	
1:B:424:ARG:HB2	1:B:426:MET:HE3	1.76	0.67	
1:A:367:THR:O	1:A:368:SEB:CB	2.43	0.67	
1:B:275:ASP:O	1:B:278:HIS:HB2	1.95	0.66	
1:B:207:LYS:NZ	1:B:207:LYS:CB	2.52	0.66	
1:B:207:LYS:HB3	1:B:207:LYS:HZ2	1.60	0.64	
1:B:397:THR:HB	1:B:469:ILE:HG23	1.80	0.62	
1:B:255:LEU:HD23	1:B:372:LEU:HD12	1.83	0.60	
1:B:220:MET:CE	1:B:220:MET:CG	2.79	0.60	
1:B:217:ILE:HG22	1:B:221:MET:HE2	1.83	0.60	
1:B:286:LEU:HB3	1:B:291:SEB:OG	2.03	0.58	
1:A:234:HIS:HD1	1:A:413:GLU:CD	2.06	0.58	
1:A:163:GLU:CD	1:A:163:GLU:N	2.57	0.57	
1:A:421:GLU:HA	1:A:426:MET:HE2	1.86	0.57	
1:B:340:ASP:OD1	1:B:385:ARG:HD3	2.05	0.56	
1:B:331:ARG:HG3	1:B:331:ARG:HH11	1.70	0.56	
1:A:397:THR:HB	1:A:469:ILE:HG23	1.89	0.55	
1:B:237:LEU:HD23	1:B:409:ARG:NH1	2.20	0.55	
1:B:337:MET:O	1:B:341:MET:HG3	2.07	0.54	
1:B:250:LEU:HG	1:B:387:MET:HE3	1.88	0.53	
1:B:424:ARG:HD2	1:B:426:MET:CE	2.39	0.53	
1:A:411:MET:HG3	1:A:439:VAL:HG22	1.91	0.52	
1:A:421:GLU:HA	1:A:426:MET:CE	2.39	0.52	
1:A:367:THR:O	1:A:368:SER:HB3	2.09	0.51	
1:A:411:MET:HE3	4:A:101:CIO:H181	1.92	0.51	
1:A:381:ILE:O	1:A:385:ARG:HG3	2.11	0.51	
1:B:244:GLN:O	1:B:247:HIS:HB3	2.12	0.50	
1:A:197:TYR:CD1	1:A:217:ILE:HD11	2.45	0.50	
1:A:411:MET:HA	1:A:411:MET:HE2	1.94	0.50	
1:B:255:LEU:CD2	1:B:372:LEU:HD12	2.43	0.49	
1:B:308:LEU:HG	1:B:342:VAL:HG11	1.93	0.49	
1:A:356:ASP:HB3	1:A:382:GLN:NE2	2.27	0.49	
1:A:242:VAL:HG12	1:A:271:ALA:HB1	1.96	0.48	
1:B:331:ARG:HG3	1:B:331:ARG:NH1	2.28	0.48	
1:B:207:LYS:CB	1:B:207:LYS:HZ2	2.20	0.48	
1:B:258:VAL:HG11	1:B:374:LEU:HD12	1.97	0.47	
1:A:411:MET:HE1	4:A:101:CIO:C18	2.45	0.47	
1:A:258:VAL:HG11	1:A:374:LEU:HD12	1.97	0.47	



Atom 1	Atom 2	Interatomic	Clash	
Atom-1	Atom-2	distance (Å)	overlap (Å)	
1:A:165:HIS:HB3	1:A:186:TYR:CE2	2.49	0.46	
1:B:390:CYS:HB3	1:B:455:TRP:CZ2	2.51	0.46	
1:B:416:GLN:O	1:B:417:GLN:C	2.52	0.46	
1:B:219:TYR:CD2	1:B:219:TYR:C	2.89	0.46	
1:A:340:ASP:OD1	1:A:385:ARG:CD	2.61	0.45	
1:A:323:PHE:HB3	1:A:326:LEU:HG	1.98	0.45	
1:B:333:THR:O	1:B:337:MET:HG3	2.16	0.45	
1:A:327:THR:HG22	1:A:330:GLN:H	1.80	0.45	
4:A:101:CIO:H20	4:A:101:CIO:H15	1.63	0.45	
1:B:395:ASN:HB2	1:B:396:PRO:HD3	1.98	0.44	
1:A:300:GLU:O	1:A:301:SER:C	2.56	0.44	
1:B:223:LEU:HD12	1:B:276:VAL:HG21	1.99	0.44	
1:B:411:MET:HG3	1:B:439:VAL:HG13	2.00	0.44	
1:B:250:LEU:CD2	1:B:387:MET:HE1	2.49	0.43	
1:A:411:MET:CE	4:A:101:CIO:C18	2.96	0.42	
1:B:213:SER:O	1:B:217:ILE:HG12	2.19	0.42	
1:B:250:LEU:HD21	1:B:387:MET:HE1	2.00	0.42	
1:B:274:HIS:O	1:B:307:HIS:CD2	2.72	0.42	
1:B:278:HIS:CE1	1:B:303:LEU:HD13	2.54	0.42	
1:B:336:LYS:HB3	1:B:336:LYS:HE2	1.93	0.42	
1:B:289:THR:CG2	1:B:426:MET:HE1	2.50	0.42	
1:A:165:HIS:HB3	1:A:186:TYR:CZ	2.54	0.42	
1:B:203:ARG:NH2	1:B:247:HIS:O	2.50	0.42	
1:A:269:PHE:CE2	1:A:273:ILE:HD13	2.55	0.41	
1:B:376:ASN:CG	1:B:377:TYR:N	2.68	0.41	
1:B:238:HIS:O	1:B:242:VAL:HG23	2.21	0.41	
1:A:275:ASP:HA	1:A:307:HIS:CD2	2.56	0.41	
4:B:102:CIO:H8	5:B:22:HOH:O	2.20	0.41	
1:B:313:LYS:NZ	1:B:316:GLN:OE1	2.54	0.41	
1:A:209:PHE:CD2	1:A:265:LEU:HD22	2.56	0.41	
1:B:327:THR:HG22	1:B:329:LYS:H	1.86	0.41	
1:A:209:PHE:CE2	1:A:265:LEU:HD22	2.57	0.40	
1:B:421:GLU:OE1	1:B:428:ILE:HA	2.21	0.40	
1:B:196:MET:HE1	1:B:268:ILE:HG22	2.04	0.40	
1:B:242:VAL:HG21	1:B:274:HIS:CE1	2.57	0.40	
1:A:275:ASP:O	1:A:278:HIS:HB2	2.22	0.40	

There are no symmetry-related clashes.



5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	320/398~(80%)	306 (96%)	10 (3%)	4 (1%)	10 8
1	В	320/398~(80%)	306 (96%)	13 (4%)	1 (0%)	37 42
All	All	640/796~(80%)	612 (96%)	23 (4%)	5 (1%)	16 16

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	А	376	ASN
1	А	256	ASP
1	А	368	SER
1	В	375	ASP
1	А	318	GLU

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	292/359~(81%)	278~(95%)	14~(5%)	21 28
1	В	291/359~(81%)	270~(93%)	21 (7%)	12 13
All	All	583/718~(81%)	548~(94%)	35~(6%)	16 19

All (35) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type			
1	А	163	GLU			



Mol	Chain	Res	Type
1	А	178	LEU
1	А	189	ASN
1	А	204	ASP
1	А	210	ARG
1	А	213	SER
1	А	295	LEU
1	А	313	LYS
1	А	326	LEU
1	А	327	THR
1	А	331	ARG
1	А	352	SER
1	А	442	SER
1	А	470	LEU
1	В	168	LYS
1	В	189	ASN
1	В	204	ASP
1	В	207	LYS
1	В	212	SER
1	В	213	SER
1	В	317	GLU
1	В	319	HIS
1	В	324	MET
1	В	329	LYS
1	В	352	SER
1	В	356	ASP
1	В	373	LEU
1	В	375	ASP
1	В	385	ARG
1	В	408	ASP
1	В	423	GLU
1	В	429	SER
1	В	436	THR
1	В	441	LYS
1	В	477	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	382	GLN
1	А	481	GLN



5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Turne	Chain	Dog	Bond lengths				Bond angles		
	туре	Unain	nes		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2
1	CME	В	194	1	8,9,10	1.69	2 (25%)	6,9,11	0.99	0
1	CME	А	194	1	8,9,10	1.53	1 (12%)	6,9,11	1.97	3 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CME	В	194	1	-	1/5/8/10	-
1	CME	А	194	1	-	1/5/8/10	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	В	194	CME	OH-CZ	-3.75	1.23	1.42
1	А	194	CME	OH-CZ	-3.68	1.23	1.42
1	В	194	CME	CA-N	-2.09	1.42	1.48

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	А	194	CME	CB-SG-SD	2.56	110.47	103.86
1	А	194	CME	CE-SD-SG	2.55	114.66	103.46
1	А	194	CME	CA-CB-SG	-2.33	104.91	114.45



There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	А	194	CME	CE-SD-SG-CB
1	В	194	CME	CE-SD-SG-CB

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Turne	Chain	Dec	B ea Link Bond lengths				B	ond ang	les
IVIOI	туре	Chain	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	CIO	В	102	-	27,27,27	1.60	3 (11%)	34,38,38	1.64	7 (20%)
4	CIO	А	101	-	27,27,27	1.38	3 (11%)	34,38,38	1.58	7 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	CIO	В	102	-	-	2/16/38/38	0/3/3/3
4	CIO	А	101	-	-	2/16/38/38	0/3/3/3

All (6) bond length outliers are listed below:



Mol	Chain	Res	Type	Atoms	Ζ	$Observed(\text{\AA})$	Ideal(Å)
4	В	102	CIO	C6-C1	-5.04	1.52	1.55
4	А	101	CIO	C6-C1	-4.67	1.52	1.55
4	В	102	CIO	C2-C1	-4.12	1.52	1.55
4	В	102	CIO	C20-C7	2.86	1.43	1.39
4	А	101	CIO	C2-C1	-2.05	1.54	1.55
4	А	101	CIO	O11-C10	2.05	1.40	1.37

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
4	А	101	CIO	C3-C2-C1	4.29	116.84	111.94
4	В	102	CIO	C12-O11-C10	4.13	123.58	117.51
4	В	102	CIO	C5-C6-C1	3.88	116.37	111.94
4	А	101	CIO	O11-C10-C13	2.91	119.35	115.40
4	В	102	CIO	O11-C10-C9	-2.85	119.50	124.30
4	А	101	CIO	C12-O11-C10	2.81	121.63	117.51
4	В	102	CIO	C5-C4-C23	-2.73	105.67	111.28
4	А	101	CIO	C8-C7-C1	2.70	123.91	121.03
4	А	101	CIO	C2-C1-C6	-2.61	105.75	107.74
4	В	102	CIO	O25-C23-C4	2.52	120.68	114.16
4	А	101	CIO	C2-C3-C4	-2.38	107.12	111.16
4	В	102	CIO	O11-C10-C13	2.37	118.61	115.40
4	A	101	CIO	C2-C1-C21	2.17	111.72	108.12
4	В	102	CIO	C2-C1-C6	-2.07	106.15	107.74

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	А	101	CIO	C16-C15-O14-C13
4	А	101	CIO	C19-C15-O14-C13
4	В	102	CIO	C16-C15-O14-C13
4	В	102	CIO	C19-C15-O14-C13

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	В	102	CIO	1	0
4	А	101	CIO	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,



bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle RSRZ \rangle$	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	322/398~(80%)	0.37	10 (3%) 51 48	16, 24, 31, 43	2 (0%)
1	В	322/398~(80%)	0.25	6 (1%) 66 62	14, 24, 30, 38	1 (0%)
All	All	644/796~(80%)	0.31	16 (2%) 58 55	14, 24, 31, 43	3 (0%)

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	А	318	GLU	7.0
1	А	485	PRO	3.9
1	В	430	PRO	3.4
1	А	257	ALA	2.8
1	В	485	PRO	2.7
1	В	272	ALA	2.6
1	А	296	MET	2.4
1	А	279	PRO	2.4
1	А	402	LEU	2.3
1	А	319	HIS	2.3
1	А	281	VAL	2.3
1	В	316	GLN	2.2
1	А	324	MET	2.2
1	А	399	SER	2.1
1	В	257	ALA	2.0
1	В	368	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CME	В	194	10/11	0.89	0.15	$33,\!36,\!55,\!57$	0
1	CME	А	194	10/11	0.93	0.15	32,36,59,64	0

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathbf{A}^2)$	Q<0.9
4	CIO	В	102	25/25	0.84	0.15	$39,\!51,\!56,\!62$	0
4	CIO	А	101	25/25	0.86	0.14	45,48,52,57	0
2	ZN	А	1001	1/1	0.95	0.08	37,37,37,37	0
3	MG	А	1002	1/1	0.95	0.06	29,29,29,29	0
2	ZN	В	1001	1/1	0.96	0.07	39,39,39,39	0
3	MG	В	1002	1/1	0.97	0.05	30,30,30,30	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







6.5 Other polymers (i)

There are no such residues in this entry.

