



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 27, 2024 – 03:11 PM EDT

PDB ID : 1XM6  
Title : Catalytic Domain Of Human Phosphodiesterase 4B In Complex With (R)-Mesopram  
Authors : Card, G.L.; England, B.P.; Suzuki, Y.; Fong, D.; Powell, B.; Lee, B.; Luu, C.; Tabrizizad, M.; Gillette, S.; Ibrahim, P.N.; Artis, D.R.; Bollag, G.; Milburn, M.V.; Kim, S.-H.; Schlessinger, J.; Zhang, K.Y.J.  
Deposited on : 2004-10-01  
Resolution : 1.92 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtrriage (Phenix) : 1.20.1  
EDS : 3.0  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.003 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.39


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.92 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	1028 (1.92-1.92)
Clashscore	180529	1100 (1.92-1.92)
Ramachandran outliers	177936	1087 (1.92-1.92)
Sidechain outliers	177891	1087 (1.92-1.92)
RSRZ outliers	164620	1028 (1.92-1.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	398	 7% 74% 10% • 15%
1	B	398	 5% 73% 10% • 16%

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 5733 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called cAMP-specific 3',5'-cyclic phosphodiesterase 4B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	338	2749	1737	459	529	24	8	0	0
1	B	334	2718	1719	454	522	23	8	0	0

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	131	MET	-	initiating methionine	UNP Q07343
A	132	GLY	-	cloning artifact	UNP Q07343
A	133	SER	-	cloning artifact	UNP Q07343
A	134	SER	-	cloning artifact	UNP Q07343
A	135	HIS	-	expression tag	UNP Q07343
A	136	HIS	-	expression tag	UNP Q07343
A	137	HIS	-	expression tag	UNP Q07343
A	138	HIS	-	expression tag	UNP Q07343
A	139	HIS	-	expression tag	UNP Q07343
A	140	HIS	-	expression tag	UNP Q07343
A	141	SER	-	cloning artifact	UNP Q07343
A	142	SER	-	cloning artifact	UNP Q07343
A	143	GLY	-	cloning artifact	UNP Q07343
A	144	LEU	-	cloning artifact	UNP Q07343
A	145	VAL	-	cloning artifact	UNP Q07343
A	146	PRO	-	cloning artifact	UNP Q07343
A	147	ARG	-	cloning artifact	UNP Q07343
A	148	GLY	-	cloning artifact	UNP Q07343
A	149	SER	-	cloning artifact	UNP Q07343
A	150	HIS	-	cloning artifact	UNP Q07343
A	151	MET	-	cloning artifact	UNP Q07343
B	131	MET	-	initiating methionine	UNP Q07343
B	132	GLY	-	cloning artifact	UNP Q07343
B	133	SER	-	cloning artifact	UNP Q07343
B	134	SER	-	cloning artifact	UNP Q07343

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	135	HIS	-	expression tag	UNP Q07343
B	136	HIS	-	expression tag	UNP Q07343
B	137	HIS	-	expression tag	UNP Q07343
B	138	HIS	-	expression tag	UNP Q07343
B	139	HIS	-	expression tag	UNP Q07343
B	140	HIS	-	expression tag	UNP Q07343
B	141	SER	-	cloning artifact	UNP Q07343
B	142	SER	-	cloning artifact	UNP Q07343
B	143	GLY	-	cloning artifact	UNP Q07343
B	144	LEU	-	cloning artifact	UNP Q07343
B	145	VAL	-	cloning artifact	UNP Q07343
B	146	PRO	-	cloning artifact	UNP Q07343
B	147	ARG	-	cloning artifact	UNP Q07343
B	148	GLY	-	cloning artifact	UNP Q07343
B	149	SER	-	cloning artifact	UNP Q07343
B	150	HIS	-	cloning artifact	UNP Q07343
B	151	MET	-	cloning artifact	UNP Q07343


- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Zn 1 1	0	0
2	B	1	Total Zn 1 1	0	0

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Mg 1 1	0	0
3	B	1	Total Mg 1 1	0	0

- Molecule 4 is (5R)-5-(4-methoxy-3-propoxyphenyl)-5-methyl-1,3-oxazolidin-2-one (three-letter code: 5RM) (formula: C<sub>14</sub>H<sub>19</sub>NO<sub>4</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			19	14	1	4		
4	B	1	Total	C	N	O	0	0
			19	14	1	4		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	118	Total	O	0	0
			118	118		
5	B	106	Total	O	0	0
			106	106		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: cAMP-specific 3',5'-cyclic phosphodiesterase 4B



- Molecule 1: cAMP-specific 3',5'-cyclic phosphodiesterase 4B



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	89.22Å 94.06Å 105.24Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	70.71 – 1.92 70.13 – 1.92	Depositor EDS
% Data completeness (in resolution range)	99.7 (70.71-1.92) 99.7 (70.13-1.92)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.71 (at 1.90Å)	Xtrriage
Refinement program	REFMAC 5.1.25	Depositor
R, $R_{free}$	0.201 , 0.227 0.206 , 0.235	Depositor DCC
$R_{free}$ test set	3428 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	37.8	Xtrriage
Anisotropy	0.248	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 36.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	5733	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	27.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 42.35 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.0671e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CME, ZN, MG, 5RM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.51	1/2772 (0.0%)	0.81	9/3751 (0.2%)
1	B	0.53	1/2741 (0.0%)	0.78	6/3709 (0.2%)
All	All	0.52	2/5513 (0.0%)	0.79	15/7460 (0.2%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	328	LYS	CB-CG	-6.00	1.36	1.52
1	A	328	LYS	CB-CG	5.62	1.67	1.52

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	392	ASP	CB-CG-OD2	7.25	124.83	118.30
1	A	214	ASP	CB-CG-OD2	6.77	124.40	118.30
1	A	340	ASP	CB-CG-OD2	6.65	124.28	118.30
1	B	356	ASP	CB-CG-OD2	6.63	124.27	118.30
1	B	468	ASP	CB-CG-OD2	6.44	124.10	118.30
1	A	468	ASP	CB-CG-OD2	6.41	124.07	118.30
1	B	340	ASP	CB-CG-OD2	6.19	123.87	118.30
1	A	392	ASP	CB-CG-OD2	6.03	123.73	118.30
1	A	230	ASP	CB-CG-OD2	5.86	123.58	118.30
1	A	375	ASP	CB-CG-OD2	5.69	123.42	118.30
1	A	256	ASP	CB-CG-OD2	5.54	123.28	118.30
1	A	385	ARG	NE-CZ-NH1	5.25	122.93	120.30
1	B	433	ASP	CB-CG-OD2	5.25	123.02	118.30
1	B	375	ASP	CB-CG-OD2	5.15	122.93	118.30
1	A	172	ASP	CB-CG-OD2	5.07	122.86	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2749	0	2669	18	0
1	B	2718	0	2641	20	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	19	0	19	0	0
4	B	19	0	19	2	0
5	A	118	0	0	1	0
5	B	106	0	0	1	0
All	All	5733	0	5348	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:340:ASP:OD1	1:A:385:ARG:HD3	1.54	1.07
1:B:340:ASP:OD1	1:B:385:ARG:HD3	1.73	0.89
1:A:340:ASP:OD1	1:A:385:ARG:CD	2.34	0.73
1:B:196:MET:HE1	1:B:243:ALA:HB1	1.71	0.70
1:A:252:THR:HG21	5:A:1075:HOH:O	2.01	0.61
1:B:340:ASP:OD1	1:B:385:ARG:CD	2.49	0.60
1:A:397:THR:HB	1:A:469:ILE:HG23	1.87	0.56
1:B:252:THR:HG21	5:B:2041:HOH:O	2.06	0.56
1:B:249:LEU:O	1:B:252:THR:HG23	2.06	0.56
1:B:442:SER:HB3	4:B:1003:5RM:H161	1.88	0.55
1:A:275:ASP:HA	1:A:307:HIS:CD2	2.42	0.55
1:A:376:ASN:CG	1:A:377:TYR:H	2.10	0.54
1:B:196:MET:CE	1:B:243:ALA:HB1	2.37	0.53
1:B:376:ASN:CG	1:B:377:TYR:H	2.11	0.53

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:275:ASP:HA	1:B:307:HIS:CD2	2.45	0.51
1:B:196:MET:HE2	1:B:196:MET:HA	1.91	0.51
1:A:249:LEU:O	1:A:252:THR:HG23	2.10	0.50
1:A:316:GLN:O	1:A:317:GLU:CB	2.60	0.50
1:B:410:ILE:HG23	1:B:411:MET:HE2	1.94	0.48
1:B:181:PHE:CD1	1:B:237:LEU:HD21	2.48	0.48
1:A:356:ASP:HB3	1:A:382:GLN:NE2	2.28	0.48
1:A:274:HIS:O	1:A:307:HIS:CD2	2.67	0.48
1:A:316:GLN:O	1:A:317:GLU:HB2	2.14	0.47
1:B:196:MET:CE	1:B:196:MET:HA	2.46	0.46
1:B:209:PHE:CD1	1:B:326:LEU:HD21	2.52	0.45
1:B:411:MET:HE1	4:B:1003:5RM:C16	2.46	0.45
1:A:184:ALA:HB2	1:A:237:LEU:HD13	1.99	0.44
1:A:270:ALA:O	1:A:274:HIS:HB3	2.16	0.44
1:A:432:CME:HZ3	1:A:510:LEU:O	2.17	0.44
1:B:274:HIS:O	1:B:307:HIS:CD2	2.70	0.44
1:B:309:ALA:O	1:B:313:LYS:HB2	2.17	0.44
1:B:356:ASP:HB3	1:B:382:GLN:NE2	2.33	0.44
1:B:395:ASN:HB2	1:B:396:PRO:HD3	2.00	0.44
1:A:431:MET:HE3	1:A:510:LEU:HD23	2.00	0.43
1:A:429:SER:OG	1:A:512:LEU:HD11	2.19	0.41
1:B:180:ILE:HA	1:B:183:VAL:HG13	2.01	0.41
1:A:209:PHE:CD1	1:A:326:LEU:HD22	2.56	0.40
1:A:350:HIS:HB3	1:A:507:GLN:HE21	1.86	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	331/398 (83%)	322 (97%)	8 (2%)	1 (0%)	37 26

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	327/398 (82%)	320 (98%)	7 (2%)	0	100	100
All	All	658/796 (83%)	642 (98%)	15 (2%)	1 (0%)	44	34

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	317	GLU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	304/357 (85%)	296 (97%)	8 (3%)	41	26
1	B	300/357 (84%)	292 (97%)	8 (3%)	40	24
All	All	604/714 (85%)	588 (97%)	16 (3%)	41	26

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	204	ASP
1	A	329	LYS
1	A	353	LEU
1	A	364	LYS
1	A	426	MET
1	A	470	LEU
1	A	477	ARG
1	A	503	MET
1	B	163	GLU
1	B	178	LEU
1	B	212	SER
1	B	352	SER
1	B	365	LYS
1	B	366	VAL
1	B	504	GLU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	508	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	382	GLN
1	A	507	GLN
1	B	382	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

6 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	CME	B	194	1	8,9,10	1.65	1 (12%)	6,9,11	1.79	2 (33%)
1	CME	A	194	1	8,9,10	1.71	2 (25%)	6,9,11	1.55	2 (33%)
1	CME	A	320	1	8,9,10	1.59	1 (12%)	6,9,11	1.51	2 (33%)
1	CME	B	432	1	8,9,10	1.67	1 (12%)	6,9,11	1.29	0
1	CME	B	320	1	8,9,10	1.76	1 (12%)	6,9,11	1.45	1 (16%)
1	CME	A	432	1	8,9,10	1.69	1 (12%)	6,9,11	1.21	1 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CME	B	194	1	-	0/5/8/10	-
1	CME	A	194	1	-	1/5/8/10	-
1	CME	A	320	1	-	2/5/8/10	-
1	CME	B	432	1	-	0/5/8/10	-
1	CME	B	320	1	-	1/5/8/10	-
1	CME	A	432	1	-	2/5/8/10	-

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	432	CME	OH-CZ	-4.07	1.21	1.42
1	B	432	CME	OH-CZ	-3.92	1.22	1.42
1	B	320	CME	OH-CZ	-3.74	1.23	1.42
1	A	194	CME	OH-CZ	-3.69	1.23	1.42
1	A	320	CME	OH-CZ	-3.68	1.23	1.42
1	B	194	CME	OH-CZ	-3.60	1.23	1.42
1	A	194	CME	CA-N	-2.06	1.42	1.48

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	194	CME	CB-SG-SD	2.88	111.30	103.86
1	B	320	CME	CZ-CE-SD	-2.44	105.23	113.39
1	B	194	CME	OH-CZ-CE	2.35	119.98	110.82
1	A	320	CME	CB-SG-SD	2.25	109.69	103.86
1	A	194	CME	OH-CZ-CE	2.23	119.53	110.82
1	A	194	CME	CA-CB-SG	-2.13	105.72	114.45
1	A	320	CME	OH-CZ-CE	2.11	119.06	110.82
1	A	432	CME	OH-CZ-CE	2.10	119.03	110.82

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	194	CME	CE-SD-SG-CB
1	A	432	CME	SD-CE-CZ-OH
1	B	320	CME	CA-CB-SG-SD
1	A	320	CME	CZ-CE-SD-SG
1	A	432	CME	CZ-CE-SD-SG
1	A	320	CME	CA-CB-SG-SD

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	432	CME	1	0

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	5RM	B	1003	-	18,20,20	1.16	1 (5%)	22,28,28	1.52	4 (18%)
4	5RM	A	1003	-	18,20,20	1.38	1 (5%)	22,28,28	1.60	3 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	5RM	B	1003	-	-	0/12/23/23	0/2/2/2
4	5RM	A	1003	-	-	0/12/23/23	0/2/2/2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1003	5RM	O5-C4	5.22	1.43	1.36
4	B	1003	5RM	O5-C4	3.94	1.42	1.36

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1003	5RM	C11-O10-C9	4.99	124.83	117.51
4	B	1003	5RM	O5-C4-N3	-3.69	107.55	109.67
4	B	1003	5RM	C11-O10-C9	3.41	122.51	117.51
4	A	1003	5RM	O5-C4-N3	-3.24	107.81	109.67
4	A	1003	5RM	C14-O13-C12	2.52	123.77	117.69
4	B	1003	5RM	O13-C12-C9	2.13	120.16	115.75
4	B	1003	5RM	O5-C4-O19	-2.12	118.87	121.64

There are no chirality outliers.

There are no torsion outliers.



There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	1003	5RM	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	335/398 (84%)	0.56	26 (7%) 20 25	17, 25, 35, 49	17 (5%)
1	B	331/398 (83%)	0.30	20 (6%) 29 33	15, 25, 34, 42	13 (3%)
All	All	666/796 (83%)	0.43	46 (6%) 24 30	15, 25, 35, 49	30 (4%)

All (46) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	508	PHE	7.6
1	A	508	PHE	7.5
1	A	512	LEU	6.9
1	A	510	LEU	6.3
1	B	502	LEU	6.1
1	B	510	LEU	6.1
1	A	502	LEU	5.3
1	B	505	LYS	5.3
1	B	511	THR	5.1
1	B	506	PHE	4.6
1	A	505	LYS	4.5
1	A	511	THR	4.5
1	A	319	HIS	4.1
1	A	507	GLN	4.1
1	B	507	GLN	3.5
1	B	504	GLU	3.5
1	A	506	PHE	3.5
1	A	318	GLU	3.3
1	A	513	ASP	3.3
1	B	501	GLY	3.2
1	A	504	GLU	3.1
1	A	369	SER	2.9
1	A	368	SER	2.9
1	B	509	GLU	2.9

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	501	GLY	2.9
1	A	375	ASP	2.8
1	B	214	ASP	2.8
1	A	287	ILE	2.8
1	A	509	GLU	2.8
1	A	499	CYS	2.8
1	A	296	MET	2.7
1	B	503	MET	2.6
1	A	327	THR	2.6
1	B	316	GLN	2.6
1	A	463	GLN	2.4
1	B	319	HIS	2.4
1	B	163	GLU	2.4
1	B	375	ASP	2.3
1	B	189	ASN	2.3
1	B	425	GLY	2.2
1	A	295	LEU	2.1
1	B	371	VAL	2.1
1	A	449	TYR	2.1
1	B	374	LEU	2.0
1	A	324	MET	2.0
1	A	317	GLU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	CME	B	320	10/11	0.89	0.16	38,40,58,61	0
1	CME	A	320	10/11	0.91	0.13	44,47,61,64	0
1	CME	A	194	10/11	0.93	0.12	28,30,48,50	0
1	CME	B	194	10/11	0.95	0.10	29,31,48,50	0
1	CME	A	432	10/11	0.95	0.12	31,34,51,53	0
1	CME	B	432	10/11	0.95	0.13	31,35,52,54	0

## 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.


## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	5RM	A	1003	19/19	0.95	0.08	31,33,36,37	0
4	5RM	B	1003	19/19	0.95	0.07	30,34,35,36	0
3	MG	B	1002	1/1	0.97	0.05	22,22,22,22	0
3	MG	A	1002	1/1	0.99	0.06	21,21,21,21	0
2	ZN	A	1001	1/1	1.00	0.02	31,31,31,31	0
2	ZN	B	1001	1/1	1.00	0.02	31,31,31,31	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [i](#)

There are no such residues in this entry.